

# A MINIMIZING-MOVEMENTS APPROACH TO GENERIC SYSTEMS

ANSGAR JÜNGEL, ULISSE STEFANELLI, AND LARA TRUSSARDI

ABSTRACT. We present a new time discretization scheme adapted to the structure of GENERIC systems. The scheme is variational in nature and is based on a *conditional* incremental minimization. The GENERIC structure of the scheme provides stability and convergence of the scheme. We prove that the scheme can be rigorously implemented in the case of the damped harmonic oscillator. Numerical evidence is collected, illustrating the performance of the method.

## 1. INTRODUCTION

The aim of this note is to discuss a new variational time-discretization scheme adapted to the structure of General Equations for Non-Equilibrium Reversible-Irreversible Coupling (GENERIC). Introduced by Grmela & Öttinger, this formulation provides a unified frame for describing the time evolution of physical systems out of equilibrium in presence of reversible and irreversible dynamics [36].

Let  $y$  denote the *state* of a closed, nonequilibrium physical system and let  $E(y)$  and  $S(y)$  be the corresponding total *energy* and total *entropy*, respectively. The GENERIC formulation of the time evolution of the system reads

$$y' = L(y) DE(y) + K(y) DS(y). \quad (1.1)$$

Here,  $y'$  denotes the time derivative,  $L$  is the antisymmetric *Poisson* operator, and  $K$  is the symmetric and positive definite *Onsager* operator (details in Section 2).

The gist of the GENERIC system (1.1) is that conservative and dissipative dynamics are clearly separated. Under a specific compatibility condition, see (2.4) below, this entails that a solution  $t \mapsto y(t)$  to (1.1) conserves energy and accumulates entropy in a quantifiable way, namely

$$\frac{d}{dt} E(y) = 0 \quad \text{and} \quad \frac{d}{dt} S(y) = \langle DS(y), K(y) DS \rangle \geq 0.$$

The conservation of energy and the quantified dissipative character are the distinguishing traits of the GENERIC system (1.1). To replicate these properties at the discrete level leads to so-called *structure-preserving* approximations. These have drawn interest in the last years, giving rise to different numerical solutions adapted to different applicative contexts.

---

*Date:* May 28, 2020.

*2010 Mathematics Subject Classification.* 80M30, 65L20.

*Key words and phrases.* Damped harmonic oscillator, structure-preserving time discretization, GENERIC system.

In the last years, GENERIC has attracted increasing attention and has been applied to a number of situations ranging from complex fluids [18], to dissipative quantum mechanics [32], to thermomechanics [5, 21–23, 31], to electromagnetism [24], to shape-memory alloys [3], to the Vlasov-Fokker-Planck [13, 14, 20] and the Boltzmann equation [35], and to large-deviation limits of reversible stochastic processes [26, 27].

Numerical schemes conserving energy can be found for instance in [16, 40, 42–44, 47]; see [8] for a review and [49] for a contribution explicitly focusing on GENERIC formulations. These schemes are often *discrete-gradient* methods, where gradients of functionals are specifically modified in order to fulfill a discrete chain rule and exactly replicate conservation [16, 17, 19, 30]. In the thermomechanical context, structure-preserving discretizations either in terms of the absolute temperature [9, 10, 15, 38] or of the internal energy or the entropy [5, 6] have been obtained. The reader is referred to [28] for an approach to open systems.

GENERIC integrators able to conserve energy and accurately describe entropy accumulation have been proposed in [37], where however some limitations are also mentioned. In particular, energy and Onsager operators have to be suitably modified in a time-step dependent manner in order energy conservation to hold. Integrators are actually constructed in case of single dissipation mechanism only (were  $K$  be a matrix, it would have to have rank one) and no convergence theory is provided. The discretization of the damped harmonic oscillator is addressed in [37], where nonetheless the temperature is given by a prescribed heat bath. In this case, explicit solutions have to be used in order to specify the GENERIC integrator.

To the best of our knowledge, all available numerical schemes directly target GENERIC systems in their differential form (1.1). Our focus is here on a time-discretization of variational nature instead, fitting into the general scheme of so-called *minimizing movements* [1, 2]. In particular, at all discrete steps we aim at solving a specific minimization problem. We start by defining the entropy-production potential  $\Psi(y, \xi) = \langle \xi, K(y)\xi \rangle / 2$  and denoting by  $\Psi^*(y, \cdot)$  its conjugate in the second variable. Then, the GENERIC relation (1.1) in  $[0, T]$  can be equivalently rewritten in terms of the scalar equation

$$-S(y(t)) + \int_0^t \Psi^*(y, y' - L(y) DE(y)) + \int_0^t \Psi(y, DS(y)) + S(y(0)) = 0 \quad (1.2)$$

for all  $t \in [0, T]$ . The reformulation of dissipative systems in terms of scalar equations as (1.2) is usually referred to as *De Giorgi's Energy-Dissipation Principle* [12, 29, 33]. This principle has already been applied in a variety of different contexts, including generalized gradient flows [4, 46], curves of maximal slope in metric spaces [2, 11], rate-independent systems [34, 41], and optimal control [39]. For GENERIC systems, the reformulation of (1.1) in terms of such variational principle has been already presented in [14].

Our new minimizing-movements approach to (1.1) consists in tackling a discrete version of relation (1.2). Assume to be given a time partition  $0 = t_0 < t_1 < \dots < t_N = T$  with steps  $\tau_i = t_i - t_{i-1}$ , and an initial state  $y^0$ . We define the time-discrete trajectory  $\{y_i\}_{i=0}^N$  by letting  $y_0 = y^0$  and subsequently perform the minimization

$$\min_y \left\{ -S(y) + \tau_i \Psi^* \left( y, \frac{y - y_{i-1}}{\tau_i} - L(y) DE(y) \right) + \tau_i \Psi(y, DS(y)) + S(y_{i-1}) \right\}$$

for  $i = 1, \dots, N$ . The latter is nothing but a localized and discretized version of the De Giorgi's Energy-Dissipation Principle (1.2).

The minimizing-movements scheme above has been introduced in [25]. The theory in [25] is however restricted to the case of state-independent operators  $L$  and  $K$ , which severely limits the applicability to real GENERIC systems. In addition, the convergence analysis in [25] relies on a suitable set of a priori assumptions, leaving open the discussion whether these can be met in practice.

The aim of this note is then threefold. First, we extend the reach of the numerical method to include the case of state dependent operators  $L(y)$  and  $K(y)$ , hence covering the full extent of the GENERIC theory (Section 2). Our main result is the conditional convergence of Theorem 2.1. Second, we provide the detailed analysis of a specific case, the damped harmonic oscillator, in which the above-mentioned convergence assumptions can actually be proved to hold (Section 3). Eventually, we present numerical experiments assessing the performance of the minimizing-movements scheme and compare it to the classical implicit Euler scheme (Subsection 3.5).

## 2. THE MINIMIZING-MOVEMENTS SCHEME FOR GENERIC SYSTEMS

In this section, we recall the structure of a GENERIC system [36] by specifying the assumptions on functionals and operators that will be used throughout. Moreover, we formulate our minimizing-movements scheme and present a conditional convergence result, namely Theorem 2.1.

The GENERIC system

$$y' = L(y) DE(y) + K(y) \partial S(y) \quad \text{a.e. in } [0, T] \quad (2.1)$$

is defined by specifying the quintuple  $(Y, E, S, L, K)$ . In the following, the state space  $Y$  is assumed to be a reflexive Banach space. The functionals  $E$  and  $S$  represent the total energy and the total entropy of the system, respectively. We assume  $E$  to be Fréchet differentiable, with strongly-weakly continuous differential  $DE$ , and  $-S : Y \rightarrow (-\infty, \infty]$  to be proper and lower semicontinuous with single-valued and strongly-weakly continuous Fréchet subdifferential  $\partial(-S)$ . Recall that one has  $\xi \in \partial(-S)(y)$  iff  $S(y) > -\infty$  and

$$\liminf_{x \rightarrow y} \frac{S(y) - S(x) - \langle \xi, x - y \rangle}{\|x - y\|_Y} \geq 0.$$

In the following, we also make use of the obvious notation  $-\partial S = \partial(-S)$ .

The operators  $L$  and  $K$  define a *Poisson* and an *Onsager* geometric structure on  $Y$ , respectively. In particular, for all states  $y \in Y$  we assume that  $L(y)$  and  $K(y)$  are linear and continuous from  $Y^*$  to  $Y$ . Moreover,  $L(y)$  is required to be antisymmetric,  $L^*(y) = -L(y)$ , and to fulfill the Jacobi identity  $\{\{g_1, g_2\}, g_3\} + \{\{g_2, g_3\}, g_1\} + \{\{g_3, g_1\}, g_2\} = 0$ . Here,  $g_i$  denotes any differentiable function on  $Y$  and the Poisson bracket is defined as  $\{g, \tilde{g}\}(y) = \langle Dg(y), L(y)D\tilde{g}(y) \rangle$ , where  $\langle \cdot, \cdot \rangle$  denotes the duality pairing between  $Y^*$  (dual) and  $Y$ . Moreover, we assume the strong-weak continuity

$$y_n \xrightarrow{Y} y \quad \text{and} \quad \xi_n \xrightarrow{Y^*} \xi \quad \Rightarrow \quad L(y_n)\xi_n \xrightarrow{Y} L(y)\xi. \quad (2.2)$$

The mapping  $K(y)$  is asked to be symmetric and positive definite, namely  $K(y) = K^*(y) \geq 0$ . We associate to  $K$  the so-called *entropy-production potential*  $\Psi : Y \times Y^* \rightarrow [0, \infty)$  given by  $\Psi(y, \xi) = \langle \xi, K(y)\xi \rangle / 2$  and let  $\Psi^*$  be its conjugate in the second variable, namely  $\Psi^*(y, \eta) = \sup_{\xi} (\langle \xi, \eta \rangle - \Psi(y, \xi))$ . We also assume the lower semicontinuity of the sum of the entropy-production potential and its dual, that is

$$\begin{aligned} y_n \xrightarrow{Y} y, \quad \eta_n \xrightarrow{Y^*} \eta, \quad \text{and} \quad \xi_n \xrightarrow{Y^*} \xi \\ \Rightarrow \liminf_{n \rightarrow \infty} (\Psi^*(y_n, \eta_n) + \Psi(y_n, \xi_n)) \geq \Psi^*(y, \eta) + \Psi(y, \xi). \end{aligned} \quad (2.3)$$

In addition, functionals and operators are asked to satisfy the crucial *noninteraction* condition

$$L^*(y) \partial S(y) = K^*(y) DE(y) = 0 \quad \forall y \in Y. \quad (2.4)$$

This condition ensures that the system of dissipative actions  $K(y) \partial S(y)$  does not spoil energy conservation and the system of conservative actions  $L(y) DE(y)$  does not contribute to dissipation. Indeed, by assuming sufficient smoothness one checks that

$$\begin{aligned} \frac{d}{dt} E(y) &= \langle DE(y), y' \rangle \stackrel{(2.1)}{=} \langle DE(y), L(y) DE(y) \rangle + \langle DE(y), K(y) \partial S(y) \rangle \\ &= 0 + \langle \partial S(y), K^*(y) DE(y) \rangle \stackrel{(2.4)}{=} 0, \\ \frac{d}{dt} S(y) &= \langle \partial S(y), y' \rangle \stackrel{(2.1)}{=} \langle \partial S(y), L(y) DE(y) \rangle + \langle \partial S(y), K(y) \partial S(y) \rangle \\ &= \langle DE(y), L^*(y) \partial S(y) \rangle + \langle \partial S(y), K(y) \partial S(y) \rangle \\ &\stackrel{(2.4)}{=} \langle \partial S(y), K(y) \partial S(y) \rangle \geq 0. \end{aligned} \quad (2.5)$$

The noninteraction condition (2.4) hence implies that trajectories  $y$  solving (2.1) have constant total energy and entropy rate  $\langle \partial S(y), K(y) \partial S(y) \rangle$ . In particular, the entropy is nondecreasing and entropy production results solely from irreversible processes. In computing (2.5) we have used the chain rule  $(d/dt)S(y) = \langle \partial S(y), y' \rangle$  almost everywhere. This is classical in case  $-S$  is (a regular perturbation of) a convex functional [7, Prop. 3.3, p. 73]. The reader is referred to [45] for a general discussion out of the convex case.

Before moving on, let us remark that the structure of GENERIC is geometric in nature. Indeed, it is invariant by coordinate changes. Let  $y = \phi(\tilde{y})$  for  $\tilde{y} \in \tilde{Y}$  and define  $\tilde{E}(\tilde{y}) = E(\phi(\tilde{y}))$ ,  $-\tilde{S}(\tilde{y}) = -S(\phi(\tilde{y}))$ ,  $\tilde{L}(\tilde{y}) = D\phi(\tilde{y})^{-1} L(\phi(\tilde{y})) D\phi(\tilde{y})^{-*}$ , and  $\tilde{K}(\tilde{y}) = D\phi(\tilde{y})^{-1} K(\phi(\tilde{y})) D\phi(\tilde{y})^{-*}$ , where  $D\phi(\tilde{y})^{-*} : \tilde{Y}^* \rightarrow Y^*$  is the adjoint of the inverse of  $D\phi(\tilde{y}) : \tilde{Y} \rightarrow Y$ . Then, the quintuple  $(\tilde{Y}, \tilde{E}, \tilde{S}, \tilde{L}, \tilde{K})$  satisfies the above structural assumptions and the GENERIC structure (2.1) can be rewritten as  $\tilde{y}' = \tilde{L}(\tilde{y}) D\tilde{E}(\tilde{y}) + \tilde{K}(\tilde{y}) \partial \tilde{S}(\tilde{y})$ .

We now reconsider the discussion leading to (1.2) and specify it further by remarking that relation (2.1) is actually equivalent to the *inequality*

$$-S(y(t)) + \int_0^t \Psi^*(y, y' - L(y) DE(y)) + \int_0^t \Psi(y, \partial S(y)) + S(y(0)) \leq 0 \quad (2.6)$$

for all  $t \in [0, T]$ . The equivalence between (2.1) and (2.6) follows from Fenchel's relations. Applied to the entropy-production potential  $\Psi$ , these relations read

$$\Psi^*(y, \eta) + \Psi(y, \xi) \geq \langle \xi, \eta \rangle \quad \forall y, \eta \in Y, \xi \in Y^*, \quad (2.7)$$

$$\Psi^*(y, \eta) + \Psi(y, \xi) = \langle \xi, \eta \rangle \quad \Leftrightarrow \quad \xi \in \partial\Psi(y, \eta), \quad (2.8)$$

where the subdifferential is taken in the second variable only. By noting that  $\partial\Psi(y, \partial S(y)) = K(y) \partial S(y)$  and using (2.7)-(2.8), one can prove the equivalences

$$(2.1) \quad \Leftrightarrow \quad y' - L(y) DE(y) = \partial\Psi(y, \partial S(y)) \quad \text{a.e.}$$

$$\stackrel{(2.8)}{\Leftrightarrow} \Psi^*(y, y' - L(y) DE(y)) + \Psi(y, \partial S(y)) - \langle y' - L(y) DE(y), \partial S(y) \rangle \leq 0 \quad \text{a.e.}$$

$$\stackrel{(2.4)}{\Leftrightarrow} \Psi^*(y, y' - L(y) DE(y)) + \Psi(y, \partial S(y)) - \frac{d}{dt} S(y) \leq 0 \quad \text{a.e.}$$

$$\stackrel{(2.7)}{\Leftrightarrow} (2.6).$$

In particular, the last left-to-right implication follows by integration while the right-to-left counterpart from the nonnegativity of the integrand, given (2.4) and (2.7).

The minimizing-movements scheme corresponds to a discretization of inequality (2.6). To each time partition  $0 = t_0 < t_1 < \dots < t_N = T$ , we associate the time steps  $\tau_i = t_i - t_{i-1}$  and the diameter  $\tau = \max \tau_i$ . Given the vector  $\{y_i\}_{i=0}^N \in Y^{N+1}$ , we introduce the backward piecewise constant and piecewise linear interpolations  $\bar{y} : [0, T] \rightarrow Y$  and  $\hat{y} : [0, T] \rightarrow Y$ ,

$$\begin{aligned} \bar{y}(0) &= \hat{y}(0), \quad \bar{y}(t) = y_i, \quad \hat{y}(t) = \frac{t - t_{i-1}}{\tau_i} y_{i-1} + \frac{t_i - t}{\tau_i} y_i, \\ &\forall t \in (t_{i-1}, t_i], \quad i = 1, \dots, N. \end{aligned}$$

We define the *incremental* functional by  $G : (0, \infty) \times Y \times Y \rightarrow (-\infty, \infty]$  as

$$G(\tau, \eta; y) = -S(y) + \tau \Psi^* \left( y, \frac{y - \eta}{\tau} - L(y) DE(y) \right) + \tau \Psi(y, \partial S(y)) + S(\eta).$$

By letting  $y_0 = y^0$  we find the discrete solution  $\{y_i\}_{i=0}^N$  by subsequently solving the minimization problem

$$\min_y G(\tau_i, y_{i-1}; y) \quad \text{for } i = 1, \dots, N. \quad (2.9)$$

Note that, for all  $\tau > 0$  and  $y_{i-1} \in Y$  with  $S(y_{i-1}) > -\infty$ , the map  $y \mapsto G(\tau_i, y_{i-1}; y)$  is strongly lower semicontinuous because of the lower semicontinuity of  $-S$ , the lower semicontinuity (2.3) of  $\Psi^* + \Psi$ , the weak-strong continuity of  $DE$  and  $\partial S$ , and the continuity (2.2) of  $L$ . In order to solve problem (2.9) one has hence to check that  $y \mapsto G(\tau_i, y_{i-1}; y)$  is strongly coercive.

The main result of this section is the following *conditional* convergence result.

**Theorem 2.1** (Conditional convergence). *Under the above assumptions, let a sequence of partitions  $0 = t_0^n < \dots < t_{N^n}^n = T$  be given with  $\tau^n \rightarrow 0$  as  $n \rightarrow \infty$ , and*

let  $\{y_i^n\}_{i=1}^{N^n}$  be such that  $y_0^n = y^0$ . Assume that

$$\sum_{i=1}^{N^n} (G(\tau_i^n, y_{i-1}^n; y_i^n))^+ \rightarrow 0 \quad \text{as } n \rightarrow \infty, \quad (2.10)$$

$$\{\widehat{y}^n\} \text{ is bounded in } H^1(0, T; Y) \text{ and takes values in } K \subset\subset Y, \quad (2.11)$$

$$\{L(\overline{y}^n) DE(\overline{y}^n)\} \text{ is bounded in } L^2(0, T; Y), \quad (2.12)$$

$$\{\partial S(\overline{y}^n)\} \text{ is bounded in } L^2(0, T; Y^*). \quad (2.13)$$

Then, up to a subsequence,  $\widehat{y}^n \rightharpoonup y$  in  $H^1(0, T; H)$ , where  $y$  is a solution of the GENERIC system (2.1) in the sense of inequality (2.6).

*Proof of Theorem 2.1.* We infer from the Aubin-Lions lemma [48, Cor. 7], upon extracting not relabeled subsequences, that

$$\begin{aligned} \widehat{y}^n &\rightharpoonup y \quad \text{in } H^1(0, T; Y), \\ \overline{y}^n, \widehat{y}^n &\rightarrow y \quad \text{in } C([0, T]; Y), \\ L(\overline{y}^n) DE(\overline{y}^n) &\rightharpoonup \ell \quad \text{in } L^2(0, T; Y), \end{aligned} \quad (2.14)$$

$$\partial S(\overline{y}^n) \rightharpoonup s \quad \text{in } L^2(0, T; Y^*). \quad (2.15)$$

As  $\partial S$  is strongly-weakly closed, we have that  $s = \partial S(y)$  almost everywhere. On the other hand, the strong-weak continuity of  $DE$  and the continuity (2.2) of  $L$  imply that  $L(\overline{y}^n) DE(\overline{y}^n) \rightarrow L(y) DE(y)$  pointwise in time, so that  $\ell = L(y) DE(y)$  almost everywhere.

Fix any  $t \in (0, T]$  and let  $t_m^n$  be such that  $t \in (t_{m-1}^n, t_m^n]$ . The discrete sequence of solutions  $\{y_i^n\}_{i=0}^{N^n}$  fulfills

$$\begin{aligned} &-S(\overline{y}^n(t)) + \int_0^{t_m^n} \Psi^*(\overline{y}^n, (\widehat{y}^n)' - L(\overline{y}^n) DE(\overline{y}^n)) + \int_0^{t_m^n} \Psi(\overline{y}^n, \partial S(\overline{y}^n)) + S(y^0) \\ &= \sum_{i=1}^m G(\tau_i^n, y_{i-1}^n; y_i^n). \end{aligned} \quad (2.16)$$

As  $\Psi(y, \cdot) \geq \Psi(y, 0) = 0$ , we conclude that  $\Psi^* \geq 0$  as well. By restricting integrals to the interval  $[0, t] \subset [0, t_m^n]$  equation (2.16) implies that

$$\begin{aligned} &-S(\overline{y}^n(t)) + \int_0^t \Psi^*(\overline{y}^n, (\widehat{y}^n)' - L(\overline{y}^n) DE(\overline{y}^n)) + \int_0^t \Psi(\overline{y}^n, \partial S(\overline{y}^n)) + S(y^0) \\ &\leq \sum_{i=1}^{N^n} (G(\tau_i^n, y_{i-1}^n; y_i^n))^+. \end{aligned} \quad (2.17)$$

We now pass to the limit inferior as  $n \rightarrow \infty$  in relation (2.17). By using convergence (2.10), the lower semicontinuity of  $-S$ , convergences (2.14)-(2.15), and the lower semicontinuity (2.3), we readily obtain that the limit  $y$  fulfills inequality (2.6).  $\square$

The conditional convergence result of Theorem 2.1 relies on the possibility of solving the inequality  $G(\tau_i^n, y_{i-1}^n; y_i^n) \leq 0$  up to a small, controllable error, and establishing some a priori bounds on the discrete solution. The validity of these conditions has to be checked on the specific problem at hand. In the coming Section 3 we give an example of a situation where (2.10)-(2.13) actually hold.

In case  $-S$  is convex, an example of  $\{y_i^n\}_{i=0}^{N^n}$  fulfilling (2.10) are the solutions of the implicit Euler scheme

$$\frac{y_i^n - y_{i-1}^n}{\tau_i^n} = L(y_i^n) DE(y_i^n) + K(y_i^n) \partial S(y_i^n) \quad (2.18)$$

whenever available. Indeed, such  $\{y_i^n\}_{i=0}^{N^n}$  fulfills

$$\begin{aligned} G(\tau_i^n, y_{i-1}^n; y_i^n) &\leq \tau_i^n \Psi^* \left( y_i^n, \frac{y_i^n - y_{i-1}^n}{\tau_i^n} - L(y_i^n) DE(y_i^n) \right) + \tau_i^n \Psi(y_i^n, \partial S(y_i^n)) \\ &\quad - \langle \partial S(y_i^n), y_i^n - y_{i-1}^n \rangle = 0, \end{aligned} \quad (2.19)$$

where the last equality follows from (2.8) and (2.18).

Note that the incremental minimization problem (2.9) may have no solutions, even if the Euler scheme is solvable. In the purely dissipative case  $L = 0$ , the existence of a solution to (2.9) is ensured as soon as  $\tau^* > 0$  exists, so that, for all  $\tau \in (0, \tau^*)$  and  $y_{i-1} \in Y$  with  $S(y_{i-1}) > -\infty$ , the function  $y \mapsto -S(y) + \tau \Psi(y, (y - y_{i-1})/\tau)$  is strongly coercive. The latter follows whenever  $-S$  has strongly compact sublevels and would imply in particular that the Euler scheme is also solvable. We anticipate however that the example discussed in Section 3 is not purely dissipative and  $-S$ , albeit convex, does not have strongly compact sublevels.

Let us mention that, in specific applications, the bounds (2.11)-(2.13) may follow from (2.10). For instance, this would be the case if the coercivity

$$\Psi^*(y, \eta) + \Psi(y, \xi) \geq c \|\eta\|_Y^2 + c \|\xi\|_{Y^*}^2 - \frac{1}{c} \quad (2.20)$$

holds for some  $c > 0$  and all  $y, \eta \in Y$  and  $\xi \in Y^*$ , namely in case  $K(y)$  is positive definite and bounded, uniformly with respect to  $y$ . This however does not apply to the example of the damped harmonic oscillator from Section 3, for  $K$  is singular there.

The quadratic nature of the entropy-production potential could be generalized to the case of  $p$ -growth with  $p > 1$  without any specific intricacy. In particular, one could consider the polynomial case  $\partial \Psi(y, \xi) = K(y) \|\xi\|_{Y^*}^{p-2} \xi$  and coercivity (2.20) would then read

$$\Psi^*(y, \eta) + \Psi(y, \xi) \geq c \|\eta\|_Y^{p'} + c \|\xi\|_{Y^*}^p - \frac{1}{c}$$

for  $1/p + 1/p' = 1$ . This setting would correspond to the case of doubly-nonlinear GENERIC dynamics, namely

$$y' = L(y) DE(y) + K(y) \|\partial S(y)\|_{Y^*}^{p-2} \partial S(y).$$

Again, under the noninteraction condition (2.4), suitably regular solutions conserve energy, since  $\langle DE(y), \partial \Psi(y, \partial S(y)) \rangle = \|\partial S(y)\|_{Y^*}^{p-2} \langle \partial S(y), K^*(y) DE(y) \rangle = 0$ , and have entropy rate  $(d/dt)(S(y)) = \|\partial S(y)\|_{Y^*}^{p-2} \langle \partial S(y), K(y) DS(y) \rangle \geq 0$ .

Let us further remark that the conditional convergence result of Theorem 2.1 can serve as an a-posteriori tool to check the convergence of time-discrete approximations  $\{y_i^n\}$ , regardless of the method used to generate them. In particular, relation (2.10) can be seen as a sort of a-posteriori error estimator.

We conclude this section by pointing out that the statement of Theorem 2.1 would hold also for other minimizing-movements schemes, where nonlinearities are

possibly evaluated explicitly. In particular, instead of  $G$  one could consider the functional  $\tilde{G} : (0, \infty) \times Y \times Y \rightarrow (-\infty, \infty]$  defined by

$$\tilde{G}(\tau, \eta; y) = -S(y) + \tau \Psi^* \left( y^1, \frac{y - \eta}{\tau} - L(y^2) DE(y^3) \right) + \tau \Psi(y^1, \partial S(y)) + S(\eta),$$

where  $y^j$ ,  $j = 1, 2, 3$ , are independently chosen to be either  $y_i$  or  $y_{i-1}$  (or else, for instance  $(y_i + y_{i-1})/2$ ). Under conditions (2.10)-(2.13), convergence would again follow as in Theorem 2.1.

### 3. THE MINIMIZING-MOVEMENTS SCHEME FOR THE DAMPED HARMONIC OSCILLATOR

In this section, we analyze the simplest GENERIC system fulfilling conditions (2.10)-(2.13). We consider the case of the damped harmonic oscillator, namely

$$mq'' + \nu q' + \kappa q + \lambda \theta = 0, \quad (3.1)$$

$$c\theta' = \nu(q')^2 + \lambda q' \theta. \quad (3.2)$$

Here,  $q \in \mathbb{R}$  represents the position of the harmonic oscillator and  $\theta > 0$  is its absolute temperature. Note that the case of  $\theta$  being the given, constant temperature of a surrounding heat bath is considered in [37] instead. The positive constants  $m$ ,  $\nu$ ,  $\kappa$ ,  $\lambda$  and  $c$  are the mass of the oscillator, the viscosity of the medium, the elastic modulus, the thermal-exchange coefficient, and the heat capacity, respectively.

By letting  $p \in \mathbb{R}$  be the momentum of the harmonic oscillator, we rewrite (3.1)-(3.2) as the first-order system

$$q' = \frac{p}{m}, \quad (3.3)$$

$$p' = -\frac{\nu p}{m} - \kappa q - \lambda \theta, \quad (3.4)$$

$$\theta' = \frac{\nu p^2}{m^2 c} + \frac{\lambda p \theta}{mc}. \quad (3.5)$$

System (3.3)-(3.5) can be written in the GENERIC form (2.1) by letting  $y = (q, p, \theta) \in Y = \mathbb{R}^2 \times (0, \infty)$  represent the state of the harmonic oscillator and by defining the total energy and the total entropy as

$$E(q, p, \theta) = \frac{p^2}{2m} + c\theta + \frac{\kappa q^2}{2} \quad \text{and} \quad S(q, p, \theta) = -\lambda q + c + c \ln \theta.$$

These definitions comply with the classical Helmholtz relations under the choice for the free energy

$$\Phi(q, p, \theta) = \frac{\kappa q^2}{2} + \lambda q \theta - c \theta \ln \theta.$$

In particular,  $S = -\partial_\theta \Phi$  and  $E = p^2/(2m) + \Phi + \theta S$ . Note that both  $E$  and  $S$  are smooth,  $-S$  is convex, and  $S(y) > -\infty$  iff  $\theta > 0$ .

The operators  $L, K : \mathbb{R}^3 \rightarrow \mathbb{R}^{3 \times 3}$  are given by

$$L(x) = \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & -\lambda \theta / c \\ 0 & \lambda \theta / c & 0 \end{pmatrix} \quad \text{and} \quad K(x) = \nu \theta \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & -p/(mc) \\ 0 & -p/(mc) & p^2/(mc)^2 \end{pmatrix}.$$



One readily checks that  $L$  is antisymmetric and a tedious computation shows that the Jacobi identity holds. On the other hand,  $K$  is symmetric and positive semi-definite, while not being invertible. Note that  $K(y)$  has rank two for all  $p \neq 0$ , so that the construction in [37] does not apply here.

By computing the gradients

$$DE(y) = (\kappa q, p/m, c), \quad DS(y) = (-\lambda, 0, c/\theta),$$

one can easily check that the noninteraction condition (2.4) holds.

Letting  $\xi = (\xi_q, \xi_p, \xi_\theta) \in \mathbb{R}^3$  and  $\eta = (\eta_q, \eta_p, \eta_\theta) \in \mathbb{R}^3$ , the entropy-production potential and its dual read

$$\begin{aligned} \Psi(y, \xi) &= \frac{1}{2} \xi \cdot K(y) \xi = \frac{\nu \theta}{2} \left( \xi_p - \frac{p}{m c} \xi_\theta \right)^2, \\ \Psi^*(y, \eta) &= \begin{cases} \frac{1}{2} \frac{\eta_p^2}{\nu \theta} & \text{if } \eta_q = 0 \text{ and } p \eta_p + m c \eta_\theta = 0, \\ \infty & \text{otherwise.} \end{cases} \end{aligned}$$

With the definitions above, the incremental functional  $G : (0, \infty) \times \mathbb{R}^3 \times \mathbb{R}^3 \rightarrow (-\infty, \infty]$  for  $y = (q, p, \theta)$  and  $\eta = (\eta_q, \eta_p, \eta_\theta)$  takes the form

$$\begin{aligned} G(\tau, \eta; y) &= \lambda q - c \ln \theta + \frac{\tau}{2} \frac{1}{\nu \theta} \left( \frac{p - \eta_p}{\tau} + \kappa q + \lambda \theta \right)^2 + \frac{\tau}{2} \nu \theta \left( \frac{p}{m \theta} \right)^2 - \lambda \eta_q + c \ln \eta_\theta \\ &\text{if } \theta > 0, \quad \frac{q - \eta_q}{\tau} = \frac{p}{m}, \quad \text{and} \quad c \frac{\theta - \eta_\theta}{\tau} = -\frac{p}{m} \frac{p - \eta_p}{\tau} - \frac{\kappa p q}{m}, \end{aligned}$$

$$G(\tau, \eta; y) = \infty \quad \text{otherwise.}$$

Assume to be given the time partition  $0 = t_0 < \dots < t_N = T$  with  $\tau_i = t_i - t_{i-1}$  for  $i = 1, \dots, N$ . By letting  $\eta = (q_{i-1}, p_{i-1}, \theta_{i-1})$  and defining  $(q_0, p_0, \theta_0) = (q^0, p^0, \theta^0)$  for some given initial state  $(q^0, p^0, \theta^0) \in \mathbb{R}^2 \times (0, \infty)$ , the incremental minimization problem (2.9) becomes

$$\begin{aligned} \min_{(q_i, p_i, \theta_i)} \left\{ \lambda q_i - c \ln \theta_i + \frac{\tau_i}{2 \nu \theta_i} \left( \frac{p_i - p_{i-1}}{\tau_i} + \kappa q_i + \lambda \theta_i \right)^2 + \frac{\tau_i \nu p_i^2}{2 m \theta_i} \right. \\ \left. - \lambda q_{i-1} + c \ln \theta_{i-1} \right\} \end{aligned} \quad (3.6)$$

under the constraints

$$\theta_i > 0, \quad (3.7)$$

$$\frac{q_i - q_{i-1}}{\tau_i} = \frac{p_i}{m}, \quad (3.8)$$

$$c \frac{\theta_i - \theta_{i-1}}{\tau_i} = -\frac{\kappa q_i p_i}{m} - \frac{p_i}{m} \frac{p_i - p_{i-1}}{\tau_i}, \quad (3.9)$$

for  $i = 1, \dots, N$ .

We devote the remainder of this section to prove that the convergence result of Theorem 2.1 applies to the scheme (3.6)-(3.9), namely that conditions (2.10)-(2.13) are fulfilled. In particular, we check that the minimization problem admits a solution  $\{y_i\}_{i=0}^N = \{(q_i, p_i, \theta_i)\}_{i=0}^N$  (Subsection 3.1), that such solution is unique if time steps are small enough (Subsection 3.2), that each such solutions fulfills

condition (2.10) in the stronger sense  $G(\tau_i, y_{i-1}; y_i) \leq 0$  (Subsection 3.3), and that the bounds (2.11)-(2.13) can be established, independently of the partition (Subsection 3.4). Eventually, numerical simulations are presented in Subsection 3.5.

**3.1. Existence of minimizers.** Assume to be given  $\eta = (q_{i-1}, p_{i-1}, \theta_{i-1}) \in \mathbb{R}^2 \times (0, \infty)$ . As  $y \mapsto G(\tau_i, \eta; y)$  is smooth on its domain, in order to prove that the scheme (3.6)-(3.9) admits a solution, we just need to check coercivity, namely that sublevels of  $y \mapsto G(\tau_i, \eta; y)$  are bounded.

By using the constraints (3.8)-(3.9), we can express  $q_i$  and  $\theta_i$  in terms of  $p_i$  and the values  $(q_{i-1}, p_{i-1}, \theta_{i-1})$  in the form

$$q_i = \tau_i \frac{p_i}{m} + q_{i-1} \quad \text{and} \quad \theta_i = f(p_i), \quad (3.10)$$

where the function  $f$  is defined by

$$f(p_i) = \theta_{i-1} - \frac{\tau_i^2 \kappa}{m^2 c} p_i^2 - \frac{\tau_i \kappa}{m c} p_i q_{i-1} - \frac{1}{m c} p_i^2 + \frac{1}{m c} p_i p_{i-1}.$$

By substituting the two expressions in the minimum problem (3.6)-(3.9), we can reduce it to a minimization in the variable  $p_i$  only. Indeed, problem (3.6)-(3.9) is equivalent to

$$\min_{p_i} F(p_i) \quad \text{under the constraint} \quad f(p_i) > 0, \quad (3.11)$$

where we have defined

$$\begin{aligned} F(p_i) := & \frac{\tau_i \lambda p_i}{m} - c \ln f(p_i) + \frac{\tau_i}{2\nu f(p_i)} \left( \frac{p_i - p_{i-1}}{\tau_i} + \frac{\tau_i \kappa}{m} p_i + \kappa q_{i-1} + \lambda f(p_i) \right)^2 \\ & + \frac{\tau_i \nu p_i^2}{2m f(p_i)} + c \ln \theta_{i-1}. \end{aligned}$$

Again, as  $F$  is smooth on its domain, in order to solve the minimization problem (3.11) we just need to prove that  $F$  has bounded sublevels. This follows from the fact that  $F(p_i) \rightarrow \infty$  if  $f(p_i) \rightarrow 0^+$  and that  $f(p_i) > 0$  iff  $p_i$  belongs to a bounded interval, depending on  $(q_{i-1}, p_{i-1}, \theta_{i-1})$ . In fact, as  $f$  is quadratic and  $\theta_{i-1} > 0$ , it follows that  $f(p_i) > 0$  if and only if  $|p_i - \hat{p}| < \hat{r}$  with

$$\hat{p} = \frac{p_{i-1} - \tau_i \kappa q_{i-1}}{2 + 2\tau_i^2 \kappa / m} \quad \text{and} \quad \hat{r} = \frac{((p_{i-1} - \tau_i \kappa q_{i-1})^2 + 4(1 + \tau_i^2 \kappa / m) m c \theta_{i-1})^{1/2}}{2 + 2\tau_i^2 \kappa / m}.$$

Hence, for all given  $(q_{i-1}, p_{i-1}, \theta_{i-1}) \in \mathbb{R}^2 \times (0, \infty)$ , we can find a solution  $p_i$  of (3.11). We conclude from (3.10) that  $(q_i, p_i, \theta_i)$  solves (3.6)-(3.9). In particular, we have that  $\theta_i > 0$ .

**3.2. Uniqueness of minimizers.** Let us remark that the minimizers  $(q_i, p_i, \theta_i)$  of (3.6)-(3.9) need not be unique in general. Still, the function  $F$  is strictly convex for  $\tau_i$  small enough, depending on the values  $(q_{i-1}, p_{i-1}, \theta_{i-1})$  and on material parameters.

Let us start by rewriting  $F$  as

$$\begin{aligned} F(p_i) &= -c \ln f(p_i) + \frac{(1 + \tau_i^2 \kappa/m)^2 (p_i + h_i)^2}{2\tau_i \nu f(p_i)} \\ &\quad + \frac{\tau_i \nu p_i^2}{2m f(p_i)} + \frac{\tau_i \lambda^2 f(p_i)}{2\nu} + r_i(p_i), \end{aligned} \quad (3.12)$$

where  $h_i = (-p_{i-1} + \tau_i \kappa q_{i-1})/(1 + \tau_i^2 \kappa/m)$  and  $r_i$  is the affine function

$$r_i(p_i) = \frac{\tau_i \lambda p_i}{m} + \frac{\lambda(1 + \tau_i^2 \kappa/m)}{\nu} (p_i + h_i) + c \ln \theta_{i-1}.$$

Using the following facts

$$\begin{aligned} \left( \frac{(p_i + h_i)^2}{f(p_i)} \right)'' &= \frac{((2f^3(p_i))^{1/2} - (2f(p_i))^{1/2} (p_i + h_i) f'(p_i))^2}{f^4(p_i)} \\ &\quad - \frac{(p_i + h_i)^2 f^2(p_i) f''(p_i)}{f^4(p_i)} > 0, \\ \left( \frac{p_i^2}{f(p_i)} \right)'' &= \frac{((2f^3(p_i))^{1/2} - (2f(p_i))^{1/2} p_i f'(p_i))^2}{f^4(p_i)} \\ &\quad - \frac{p_i^2 f^2(p_i) f''(p_i)}{f^4(p_i)} > 0, \end{aligned}$$

and differentiating  $F$  in (3.12) twice, we obtain

$$\begin{aligned} F''(p_i) &> -c(\ln f(p_i))'' + \frac{\tau_i \lambda^2}{2\nu} f''(p_i) = \frac{c(f'(p_i))^2}{f^2(p_i)} + f''(p_i) \left( -\frac{c}{f(p_i)} + \frac{\tau_i \lambda^2}{2\nu} \right) \\ &\geq f''(p_i) \left( -\frac{c}{f(p_i)} + \frac{\tau_i \lambda^2}{2\nu} \right) =: g(p_i). \end{aligned}$$

As  $f''(p_i) = -2(1 + \tau_i^2 \kappa/m)/(mc) < 0$ , the function  $g$  is nonnegative as long as  $f(p_i) \leq 2\nu c/(\tau_i \lambda^2)$ . It hence suffices to choose  $\tau_i$  so small that

$$\max f \leq \frac{2\nu c}{\tau_i \lambda^2}. \quad (3.13)$$

This condition depends on the size of the time-step and on material parameters and implies that  $F$  is strictly convex, thus admitting a unique minimizer  $p_i$ . Consequently, problem (3.6)-(3.9) has a unique minimizer  $(q_i, p_i, \theta_i)$ .

We can make the time-step dependent bound (3.13) on  $\tau_i$  more explicit by directly computing the maximum of  $f$ :

$$\begin{aligned} \max f &= \frac{\left( \frac{-\tau_i \kappa q_{i-1}}{mc} + \frac{p_{i-1}}{mc} \right)^2 + 4 \left( \frac{\tau_i^2 \kappa}{m^2 c} + \frac{1}{mc} \right) \theta_{i-1}}{4 \left( \frac{\tau_i^2 \kappa}{m^2 c} + \frac{1}{mc} \right)} \\ &= \frac{(-\tau_i \kappa q_{i-1} + p_{i-1})^2}{4(\tau_i^2 \kappa c + mc)} + \theta_{i-1} \leq \frac{p_{i-1}^2}{2mc} + \frac{\tau_i^2 \kappa^2 q_{i-1}^2}{2mc} + \theta_{i-1}. \end{aligned}$$

In Subsection 3.4 we will check that above right-hand side can be bounded by a positive constant  $c_0$  in terms of the initial data and of material parameters only,

uniformly with respect to  $i$ , see (3.22). Condition (3.13) can hence be strengthened by asking the partition to be such that

$$c_0 \leq \min_i \frac{2\nu c}{\tau_i \lambda^2}. \quad (3.14)$$

This stronger condition ensures the unique solvability of the minimization problem (3.6)-(3.9) for all  $i = 1, \dots, N$ .

**3.3. Minimizers fulfill condition (2.10).** Following the discussion of Section 2, the convexity of  $-S$  ensures that condition (2.10) holds as soon as one can prove that the implicit Euler scheme (2.18) is solvable. In the current setting, the implicit Euler scheme reads

$$\frac{q_i - q_{i-1}}{\tau_i} = \frac{p_i}{m}, \quad (3.15)$$

$$\frac{p_i - p_{i-1}}{\tau_i} = -\frac{\nu p_i}{m} - \kappa q_i - \lambda \theta_i, \quad (3.16)$$

$$\frac{\theta_i - \theta_{i-1}}{\tau_i} = \frac{\nu p_i^2}{m^2 c} + \frac{\lambda p_i \theta_i}{m c} \quad (3.17)$$

for  $i = 1, \dots, N$ .

Let the initial data  $(q^0, p^0, \theta^0) \in \mathbb{R}^2 \times (0, \infty)$  be given and assume to have solved (3.15)-(3.17) up to level  $i - 1$ . In particular, let  $(q_{i-1}, p_{i-1}, \theta_{i-1}) \in \mathbb{R}^2 \times (0, \infty)$ . Relation (3.15) can be written as

$$q_i = \frac{\tau_i p_i}{m} + q_{i-1}$$

and we can substitute it into (3.16) obtaining

$$p_i = \frac{-\tau_i m \lambda}{m + \tau_i \nu + \tau_i^2 \kappa} \theta_i + \frac{-\tau_i m \kappa q_{i-1} + m p_{i-1}}{m + \tau_i \nu + \tau_i^2 \kappa} =: \alpha_i \theta_i + \beta_i \quad (3.18)$$

where  $\alpha_i$  and  $\beta_i$  depend just on material parameters,  $\tau_i$ , and on  $q_{i-1}$  and  $p_{i-1}$ .

We may hence rewrite equation (3.17) as

$$\begin{aligned} \theta_i &= \theta_{i-1} + \frac{\tau_i \nu p_i^2}{m^2 c} + \frac{\tau_i \lambda p_i \theta_i}{m c} \\ &= \theta_{i-1} + \frac{\tau_i \nu}{m^2 c} \left( \alpha_i^2 \theta_i^2 + 2\alpha_i \beta_i \theta_i + \beta_i^2 \right) + \frac{\tau_i \lambda}{m c} \left( \alpha_i \theta_i^2 + \beta_i \theta_i \right), \end{aligned}$$

which gives the expression

$$\begin{aligned} 0 &= \left( \frac{\tau_i \nu \alpha_i^2}{m^2 c} + \frac{\tau_i \lambda \alpha_i}{m c} \right) \theta_i^2 + \left( \frac{2\tau_i \nu \alpha_i \beta_i}{m^2 c} + \frac{\tau_i \lambda \beta_i}{m c} - 1 \right) \theta_i + \left( \theta_{i-1} + \frac{\tau_i \nu \beta_i^2}{m^2 c} \right) \\ &=: \gamma_i \theta_i^2 + \delta_i \theta_i + \epsilon_i, \end{aligned} \quad (3.19)$$

with  $\gamma_i$ ,  $\delta_i$ , and  $\epsilon_i$  depending on  $\alpha_i$ ,  $\beta_i$ ,  $\theta_{i-1}$ , and data. Note that

$$\gamma_i = \frac{\tau_i}{m c} \left( \frac{\nu \alpha_i}{m} + \lambda \right) \alpha_i = -\frac{\tau_i^2 \lambda^2}{c} \frac{m + \tau_i^2 \kappa}{(m + \tau_i \nu + \tau_i^2 \kappa)^2} < 0.$$

We infer from  $\theta_{i-1} > 0$  that  $\epsilon_i > 0$ . Hence, the second-order equation (3.19) has a unique solution  $\theta_i > 0$ . This uniquely defines  $p_i$  and  $q_i$  via (3.18) and (3.15), respectively. In particular, the implicit Euler scheme (3.15)-(3.17) has a unique solution  $y_i^e = (q_i, p_i, \theta_i)$ .

Let now  $y_i$  be a solution to the incremental minimization problem (3.6). Owing to (2.19), we conclude that

$$G(\tau_i, y_{i-1}; y_i) = \min_y G(\tau_i, y_{i-1}; y) \leq G(\tau_i, y_{i-1}; y_i^e) \stackrel{(2.19)}{=} 0,$$

where we have also used the fact that  $y_i^e$  fulfills the constraints (3.7)-(3.9). In particular, condition (2.10) holds in the even stronger form

$$G(\tau_i, y_{i-1}; y_i) \leq 0 \quad \text{for } i = 1, \dots, N. \quad (3.20)$$

**3.4. A priori bounds.** We now prove that condition (2.11)-(2.13) of Theorem 2.1 hold. This will follow by checking a priori bounds on minimizers  $\{y_i\}_{i=0}^N$  of (3.6), independently from the time partition.

Let us start by remarking that the energy is nonincreasing. By (3.8), one can equivalently rewrite the constraint (3.9) as

$$c\theta_i - c\theta_{i-1} = -\frac{\kappa q_i^2}{2} + \frac{\kappa q_{i-1}^2}{2} - \frac{\kappa}{2}|q_i - q_{i-1}|^2 - \frac{p_i^2}{2m} + \frac{p_{i-1}^2}{2m} - \frac{1}{2m}|p_i - p_{i-1}|^2.$$

A rearrangement of the terms leads to

$$E(y_i) + \frac{\kappa}{2}|q_i - q_{i-1}|^2 + \frac{1}{2m}|p_i - p_{i-1}|^2 = E(y_{i-1}).$$

Hence,  $E(y_i)$  is nonincreasing and  $E(y_i) = E(y_{i-1})$  iff  $q_i = q_{i-1}$  and  $p_i = p_{i-1}$ . By taking the sum for  $i = 1, \dots, j$  for  $j \leq N$ , we find that

$$E(y_j) + D_j^q + D_j^p = E(y^0) \quad \forall j = 1, \dots, N, \quad (3.21)$$

where we have set

$$D_j^q = \sum_{i=1}^j \frac{\kappa}{2}|q_i - q_{i-1}|^2 \quad \text{and} \quad D_j^p = \sum_{i=1}^j \frac{1}{2m}|p_i - p_{i-1}|^2.$$

The nonnegative terms  $D_j^q$  and  $D_j^p$  exactly quantify the energy dissipativity of the scheme. Owing to (3.21), we obtain the uniform bound

$$|p_i| + |\theta_i| + |q_i| \leq C \quad \forall i = 1, \dots, N, \quad (3.22)$$

where, here and in the following, the symbol  $C$  denotes a generic positive constant depending on the initial data  $y^0$  and on material parameters, but not on the time partition. Bound (3.22) and constraint (3.8) imply that

$$\left| \frac{q_i - q_{i-1}}{\tau_i} \right| \leq C \quad \forall i = 1, \dots, N. \quad (3.23)$$

Moreover, we readily check that

$$|L(y_i) DE(y_i)| = |(p_i/m, -kq_i - \lambda\theta_i, \lambda p_i \theta_i/m)| \stackrel{(3.22)}{\leq} C \quad \forall i = 1, \dots, N. \quad (3.24)$$

Let us take the sum for  $i = 1, \dots, j$  for  $j \leq N$  in (3.20) getting

$$-S(y_j) + \psi_j^* + \psi_j \leq -S(y^0),$$

where

$$\psi_j^* = \sum_{i=1}^j \frac{\tau_i}{2\nu\theta_i} \left( \frac{p_i - p_{i-1}}{\tau_i} + \kappa q_i + \lambda\theta_i \right)^2 \quad \text{and} \quad \psi_j = \sum_{i=1}^j \frac{\tau_i \nu p_i^2}{2m\theta_i}.$$

As  $\psi_j^*$  and  $\psi_j$  are nonnegative, we infer that  $S(y_i)$  is nondecreasing. In particular,

$$-c \ln \theta_j \leq -S(y^0) - \lambda q_j + c \stackrel{(3.22)}{\leq} C \quad \forall j = 1, \dots, N.$$

and consequently,

$$\theta_i \geq \theta_{\min} > 0 \quad \forall i = 1, \dots, N, \quad (3.25)$$

for some given  $\theta_{\min}$  depending just on  $y^0$  and the material parameters. This ensures that

$$|DS(y_i)| = |(-\lambda, 0, c/\theta_i)| \leq C \quad \forall i = 1, \dots, N. \quad (3.26)$$

It follows from the bound (3.22) on  $q_i$  and  $\theta_i$  that  $S(y_i) = -\lambda q_i + c + c \ln \theta_i \leq C$  for all  $i = 1, \dots, N$ . We conclude that  $\psi_N \leq -S(y^0) + S(y_N) \leq C$  and

$$\begin{aligned} & \sum_{i=1}^N \frac{\tau_i}{2\nu\theta_i} \left| \frac{p_i - p_{i-1}}{\tau_i} \right|^2 \\ &= \psi_N^* - \sum_{i=1}^N \frac{\tau_i}{2\nu\theta_i} (\kappa q_i + \lambda \theta_i)^2 - 2 \sum_{i=1}^N \frac{\tau_i}{2\nu\theta_i} \left( \frac{p_i - p_{i-1}}{\tau_i} \right) (\kappa q_i + \lambda \theta_i) \\ &\leq \psi_N^* + \sum_{i=1}^N \frac{\tau_i}{2\nu\theta_i} (\kappa q_i + \lambda \theta_i)^2 + \sum_{i=1}^N \frac{\tau_i}{4\nu\theta_i} \left| \frac{p_i - p_{i-1}}{\tau_i} \right|^2 \\ &\stackrel{(3.22)}{\leq} C + \sum_{i=1}^N \frac{\tau_i}{4\nu\theta_i} \left| \frac{p_i - p_{i-1}}{\tau_i} \right|^2. \end{aligned} \quad (3.27)$$

Using the fact that  $\theta_i$  is uniformly bounded by (3.22), we can hence bound

$$\sum_{i=1}^N \tau_i \left| \frac{p_i - p_{i-1}}{\tau_i} \right|^2 \leq 4\nu (\max_i \theta_i) \sum_{i=1}^N \frac{\tau_i}{4\nu\theta_i} \left| \frac{p_i - p_{i-1}}{\tau_i} \right|^2 \stackrel{(3.22)+(3.27)}{\leq} C. \quad (3.28)$$

Eventually, we infer from constraint (3.9) that

$$\sum_{i=1}^N \tau_i \left| \frac{\theta_i - \theta_{i-1}}{\tau_i} \right|^2 \leq 2 \sum_{i=1}^N \tau_i \left( \frac{\kappa^2 q_i^2 p_i^2}{m^2 c^2} + \frac{p_i^2}{m^2 c^2} \left| \frac{p_i - p_{i-1}}{\tau_i} \right|^2 \right) \leq C. \quad (3.29)$$

Bounds (3.22)-(3.23) and (3.28)-(3.29) imply condition (2.11) from Theorem 2.1. On the other hand, bounds (3.24) and (3.26) imply conditions (2.12) and (2.13), respectively. Hence, the convergence statement of Theorem 2.1 holds. In particular, given initial values  $(q^0, p^0, \theta^0) \in \mathbb{R}^3$  with  $\theta^0 > 0$ , a sequence of partitions  $0 = t_0^n < \dots < t_{N^n}^n = T$  with  $\tau^n = \max_i (t_i^n - t_{i-1}^n) \rightarrow 0$ , and corresponding minimizers  $\{(q_i^n, p_i^n, \theta_i^n)\}_{i=0}^{N^n}$  of (3.6), it follows that

$$(\widehat{q}^n, \widehat{p}^n, \widehat{\theta}^n) \rightharpoonup (q, p, \theta) \quad \text{in } H^1(0, T; \mathbb{R}^3), \quad (3.30)$$

where  $(q, p, \theta)$  solves the damped harmonic oscillator system (3.3)-(3.5) with initial value  $(q(0), p(0), \theta(0)) = (q^0, p^0, \theta^0)$ . Note that solutions of (3.3)-(3.5) are unique. Hence, convergence (3.30) holds for the whole sequence of discrete solutions, not just for a subsequence.

As  $E$  and  $S$  are smooth,  $\overline{y}^n$  is bounded, and  $\theta_i \geq \theta_{\min} > 0$ , we find that  $E(\overline{y}(\cdot)) \rightarrow E(y(\cdot)) \equiv E(y^0)$  and  $S(\overline{y}(\cdot)) \rightarrow S(y(\cdot))$  uniformly, together with their

time derivatives of all orders. More precisely, upon remarking that

$$D_N^q \leq \tau^n \sum_{i=1}^N \frac{\tau_i \kappa}{2} \left| \frac{q_i - q_{i-1}}{\tau_i} \right|^2 \stackrel{(3.23)}{\leq} C\tau^n,$$

$$D_N^p \leq \tau^n \sum_{i=1}^N \frac{\tau_i}{2m} \left| \frac{p_i - p_{i-1}}{\tau_i} \right|^2 \stackrel{(3.28)}{\leq} C\tau^n,$$

we may use (3.21) and check that

$$-C\tau^n + E(\bar{y}^n(t)) \leq E^0 \quad \forall t \in [0, T].$$

In particular, we control the energy dissipation as follows:

$$\max_{t \in [0, T]} \|E(\bar{y}^n(t)) - E(y^0)\|_{L^\infty(0, T)} \leq C\tau^n. \quad (3.31)$$

**3.5. Numerical tests.** We record here some numerical evidence on the performance of the minimizing-movements scheme (3.6)-(3.9). All computations are performed in Matlab. In the following, we choose

$$m = \nu = \kappa = \lambda = c = 1, \quad y^0 = (q^0, p^0, \theta^0) = (1, 1, 1), \quad T = 15. \quad (3.32)$$

Given a uniform time-partition with time step  $\tau = T/N$ , we find a solution  $\{y_i\}_{i=0}^N$  of the minimizing-movements scheme (3.6)-(3.9) via Newton's method. Note that, by choosing (3.32) and recalling the bound (3.21), one has that the constant  $c_0$  in (3.14) can be bounded from above by  $E(y^0) = 2$ , for all  $\tau \leq 1$ . In particular, condition (3.14) is fulfilled by all  $\tau \leq 1$  and the solution of the minimizing-movements scheme is unique.

The unique solution  $\{y_i^e\}_{i=0}^N$  of the implicit Euler scheme (3.15)-(3.17) is obtained directly from (3.18)-(3.19). Eventually, the numerical reference solution  $t \mapsto y(t)$  is calculated by means of the Matlab solver ode45 choosing  $10^{-4}$  for the maximal time step and  $10^{-8}$  for the absolute tolerance.

Figures 1-3 illustrate the numerical reference solution and the time-discrete solutions for  $\tau = 1/4$ . Both the minimizing-movements and the Euler scheme dissipate energy, see Figure 3 left. This is of course an undesired effect, which is however attenuated as  $\tau \rightarrow 0$ , see (3.31). Energy dissipation seems to be more pronounced for the Euler scheme. On the other hand, entropy is nondecreasing for the minimizing-movements scheme whereas the Euler scheme shows a nonmonotone entropy, which is nonphysical, see Figure 3 right.

A second set of experiments is illustrated in Figure 4. For the same choices in (3.32) and different time steps

$$\tau^n = 2^{-n} \quad \text{for } n = -1, 0, \dots, 11, \quad (3.33)$$

we compute the uniform errors of the temperature component of the solution and of the energy with respect to the numerical reference solution.

As  $\tau$  converges to 0, our computations confirm that both the minimizing-movements and the Euler scheme are of order  $\tau$ , see also (3.31). Let us mention that a proof of first-order convergence for the minimizing-movements scheme in the nondissipative regime ( $L = 0$ ,  $K$  independent of the state) is given in [25, Prop. 4.3]. Both schemes do not conserve energy. Still, the minimizing-movements scheme is more accurate than the Euler scheme when the time steps are large.




FIGURE 1. Position  $q$  (left) and momentum  $p$  (right) with respect to time for the numerical reference solution (red, dotted), the minimizing-movements scheme (solid), and the Euler scheme (blue, dash-dotted).




FIGURE 2. Temperature with respect to time for the numerical reference solution (red, dotted), the minimizing-movements scheme (solid), and the Euler scheme (blue, dash-dotted).




FIGURE 3. Energy (left) and entropy (right) as function of time for the numerical reference solution (red dotted), the minimizing-movements scheme (solid), and the Euler scheme (blue, dash-dotted).

#### ACKNOWLEDGEMENTS

This research is supported by Austrian Science Fund (FWF) project F 65 and W 1245. AJ is supported by the FWF projects P 30000 and P 33010. US is supported by the FWF projects I4354 and P 32788 and by the Vienna Science and Technology Fund (WWTF) through Project MA14-009.






FIGURE 4. Errors with respect to  $1/\tau^n$  from (3.33), logarithmic scales for the variable  $\theta$  (on the left) and the energy (on the right). The (green) dotted lines indicate the order of convergence 1. Left:  $\max_{t \in [0, T]} |\theta(t) - \hat{\theta}(t)|$  (solid),  $\max_{t \in [0, T]} |\theta(t) - \hat{\theta}^e(t)|$  (blue, dash-dotted); right:  $\max_{t \in [0, T]} |E(\hat{y}(t)) - E(y^0)|$  (solid), and  $\max_{t \in [0, T]} |E(\hat{y}^e(t)) - E(y^0)|$  (blue, dash-dotted).

## REFERENCES

- [1] L. Ambrosio. Minimizing movements. *Rend. Accad. Naz. Sci. XL Mem. Mat. Appl.* (5), 19 (1995), 191–246.
- [2] L. Ambrosio, N. Gigli, G. Savaré. *Gradient Flows in Metric Spaces and in the Space of Probability Measures*, second ed., Birkhäuser Verlag, Basel, 2008.
- [3] F. Auricchio, E. Boatti, A. Reali, U. Stefanelli. Gradient structures for the thermomechanics of shape-memory materials. *Comput. Methods Appl. Mech. Engrg.* 299 (2016), 440–469
- [4] A. Bacho, E. Emmrich, A. Mielke. An existence result and evolutionary  $\Gamma$ -convergence for perturbed gradient systems. *J. Evol. Equ.* 19 (2019), 479–522.
- [5] P. Betsch, M. Schiebl. GENERIC-based formulation and discretization of initial boundary value problems for finite strain thermoelasticity. *Comput. Mech.* 65 (2020), 503–531.
- [6] P. Betsch, M. Schiebl. Energymomentumentropy consistent numerical methods for large strain thermoelasticity relying on the GENERIC formalism. *Internat. J. Numer. Methods Engrg.* 119 (2019), 1216–1244.
- [7] H. Brézis. *Opérateurs maximaux monotones et semi-groupes de contractions dans les espaces de Hilbert*. Number 5 in North Holland Math. Studies. North-Holland, Amsterdam, 1973.
- [8] E. Celledoni, V. Grimm, R. McLachlan, D. McLaren, D. O’Neale, B. Owren, G. Quispel. Preserving energy resp. dissipation in numerical PDEs using the “Average Vector Field” method. *J. Comput. Phys.* 231 (2012), 6770–6789.
- [9] S. Conde Martín, P. Betsch, J. C. García Orden. A temperature based thermodynamically consistent integration scheme for discrete thermo-elastodynamics. *Commun. Nonlinear Sci. Numer. Simul.* 32 (2016), 63–80.
- [10] S. Conde Martín, J. C. García Orden. On energyentropy momentum integration methods for discrete thermo-viscoelastodynamics. *Comput. Struct.* 181 (2017), 3–20.
- [11] E. De Giorgi, A. Marino, M. Tosques. Problems of evolution in metric spaces and maximal decreasing curve, *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur.* (8), 68 (1980), 180–187.
- [12] P. Dondl, T. Frenzel, A. Mielke. A gradient system with a wiggly energy and relaxed EDP-convergence. *ESAIM Control Optim. Calc. Var.* 25 (2019), Art. 68, 45 pp.
- [13] M. H. Duong. Formulation of the relativistic heat equation and the relativistic kinetic Fokker-Planck equations using GENERIC. *Phys. A*, 439 (2015), 3443.
- [14] M. H. Duong, M. A. Peletier, J. Zimmer. GENERIC formalism of a Vlasov–Fokker–Planck equation and connection to large-deviation principles. *Nonlinearity*, 26 (2013), 2951–2971.
- [15] J. C. García Orden, I. Romero. Energy-Entropy-Momentum integration of discrete thermo-visco-elastic dynamics. *Europ. J. Mech. Solids*, 32 (2012), 76–87.
- [16] O. Gonzalez. Time integration and discrete Hamiltonian systems. *J. Nonlin. Sci.* 6 (1996), 449–467.
- [17] O. Gonzalez. Exact energy-momentum conserving algorithms for general models in nonlinear elasticity. *Comp. Methods Appl. Mech. Engrg.* 190 (2000), 1763–1783.

- [18] M. Grmela, H. C. Öttinger. Dynamics and thermodynamics of complex fluids. I. Development of a general formalism. *Phys. Rev. E*, 56 (1997), 6620–6632.
- [19] E. Hairer, C. Lubich, G. Wanner. *Geometric Numerical Integration*, second edition, Springer, Berlin, 2006.
- [20] M. Hoyuelos. GENERIC framework for the Fokker-Planck equation. *Phys. A*, 442 (2016), 350–358.
- [21] M. Hütter, T. A. Tervoort. Finite anisotropic elasticity and material frame indifference from a nonequilibrium thermodynamics perspective. *J. Non Newton. Fluid. Mech.* 152 (2008), 45–52.
- [22] M. Hütter, B. Svendsen. On the formulation of continuum thermodynamic models for solids as general equations for non-equilibrium reversible/irreversible coupling. *J. Elast.* 104 (2011), 357–368.
- [23] M. Hütter, B. Svendsen. Thermodynamic model formulation for viscoplastic solids as general equations for nonequilibrium reversible/irreversible coupling. *Contin. Mech. Thermodyn.* 24 (2012), 211–227.
- [24] A. Jelić, M. Hütter, H. C. Öttinger. Dissipative electromagnetism from a nonequilibrium thermodynamics perspective. *Phys. Review E*, 74 (2006), 041126, 8 pp.
- [25] A. Jüngel, U. Stefanelli, L. Trussardi. Two structure-preserving time discretizations for gradient flows. *Appl. Math. Optim.* 80 (2020), 733–764.
- [26] R. C. Kraaij, A. Lazarescu, C. Maes, M. Peletier. Fluctuation symmetry leads to GENERIC equations with non-quadratic dissipation. *Stochastic Process. Appl.* 130 (2020), 139–170.
- [27] R. Kraaij, A. Lazarescu, C. Maes, M. Peletier. Deriving GENERIC from a generalized fluctuation symmetry. *J. Stat. Phys.* 170 (2018), 492–508.
- [28] M. Krüger, M. Groß, P. Betsch. An energyentropy-consistent time stepping scheme for non-linear thermo-viscoelastic continua. *ZAMM Z. Angew. Math. Mech.* 96 (2016), 141–178.
- [29] M. Liero, A. Mielke, M. A. Peletier, D. R. M. Renger. On microscopic origins of generalized gradient structures. *Discrete Contin. Dyn. Syst. Ser. S*, 10 (2017), 1–35.
- [30] R. McLachlan, G. Quispel, N. Robidoux. Geometric integration using discrete gradients. *Phil. Trans. R. Soc. Lond. A*, 357 (1999), 1021–1045.
- [31] A. Mielke. Formulation of thermoelastic dissipative material behavior using GENERIC. *Contin. Mech. Thermodyn.* 23 (2011), 233–256.
- [32] A. Mielke. Dissipative quantum mechanics using GENERIC, Proc. of the conference on *Recent Trends in Dynamical Systems*, vol. 35 of Proceedings in Mathematics & Statistics, Springer, 2013, pp. 555–585.
- [33] A. Mielke, A. Montefusco, M. Peletier. Exploring families of energy-dissipation landscapes via tilting – three types of EDP convergence. arXiv:2001.01455, 2020.
- [34] A. Mielke, R. Rossi, G. Savaré. BV solutions and viscosity approximations of rate-independent systems. *ESAIM Control Optim. Calc. Var.* 18 (2012), 36–80.
- [35] A. Montefusco, F. Consonni, G. Beretta. Essential equivalence of the general equation for the nonequilibrium reversible-irreversible coupling (GENERIC) and steepest-entropy-ascent models of dissipation for nonequilibrium thermodynamics. *Phys. Rev. E (3)*, 91 (2015), 042138, 19 pp.
- [36] H. C. Öttinger. *Beyond Equilibrium Thermodynamics*. John Wiley, New Jersey, 2005.
- [37] H. C. Öttinger. Generic integrators: structure preserving time integration for thermodynamic systems. *J. Non Equil. Thermodyn.* 43 (2018), 89–100.
- [38] D. Portillo, J. C. García Orden, I. Romero. Energyentropy momentum integration schemes for general discrete non-smooth dissipative problems in thermomechanics. *Internat. J. Numer. Methods Engrg.* 112 (2017), 776–802.
- [39] L. Portinale, U. Stefanelli. Penalization via global functionals of optimal-control problems for dissipative evolution. *Adv. Math. Sci. Appl.* 28 (2019), 425–447.
- [40] G. Quispel, D. McLaren. A new class of energy-preserving numerical integration methods. *J. Phys. A: Math. Theor.* 41 (2008), 045206, 7 pages.
- [41] T. Roche, R. Rossi, U. Stefanelli. Stability results for doubly nonlinear differential inclusions by variational convergence. *SIAM J. Control Optim.* 52 (2014), 1071–1107.
- [42] I. Romero. Thermodynamically consistent time-stepping algorithms for non-linear thermo-mechanical systems. *Internat. J. Numer. Methods Engrg.* 79 (2009), 706–732.
- [43] I. Romero. Algorithms for coupled problems that preserve symmetries and the laws of thermodynamics. Part I: monolithic integrators and their application to finite strain thermoelasticity. *Comput. Methods Appl. Mech. Engrg.* 199 (2010), 1841–1858.

- [44] I. Romero. Algorithms for coupled problems that preserve symmetries and the laws of thermodynamics. Part II: fractional step methods. *Comput. Methods Appl. Mech. Engrg.* 199 (2010), 2235–2248.
- [45] R. Rossi, G. Savaré. Gradient flows of non convex functionals in Hilbert spaces and applications. *ESAIM Control Optim. Calc. Var.* 12 (2006), 564–614.
- [46] R. Rossi, A. Mielke, G. Savaré. A metric approach to a class of doubly nonlinear evolution equations and applications. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 7 (2008), 97–169.
- [47] S. Sato, T. Matsuo, H. Suzuki, D. Furihata. A Lyapunov-type theorem for dissipative numerical integrators with adaptive time-stepping. *SIAM J. Numer. Anal.* 53 (2015), 2505–2518.
- [48] J. Simon. Compact sets in the space  $L^p(0, T; B)$ . *Ann. Mat. Pura Appl. (4)*, 146 (1987), 65–96.
- [49] Y. Suzuki, M. Ohnawa. GENERIC formalism and discrete variational derivative method for the two-dimensional vorticity equation. *J. Comput. Appl. Math.* 296 (2016), 690–708.

(Ansgar Jüngel) INSTITUTE FOR ANALYSIS AND SCIENTIFIC COMPUTING, VIENNA UNIVERSITY OF TECHNOLOGY, WIEDNER HAUPTSTRASSE 8-10, 1040 WIEN, AUSTRIA

*E-mail address:* [juengel@tuwien.ac.at](mailto:juengel@tuwien.ac.at)

*URL:* <http://www.asc.tuwien.ac.at/~juengel>

(Ulisse Stefanelli) FACULTY OF MATHEMATICS, UNIVERSITY OF VIENNA, OSKAR-MORGENSTERN-PLATZ 1, A-1090 VIENNA, AUSTRIA, VIENNA RESEARCH PLATFORM ON ACCELERATING PHOTOREACTION DISCOVERY, UNIVERSITY OF VIENNA, WÄHRINGERSTRASSE 17, 1090 WIEN, AUSTRIA, & ISTITUTO DI MATEMATICA APPLICATA E TECNOLOGIE INFORMATICHE *E. Magenes*, VIA FERRATA 1, I-27100 PAVIA, ITALY

*E-mail address:* [ulisse.stefanelli@univie.ac.at](mailto:ulisse.stefanelli@univie.ac.at)

*URL:* <http://www.mat.univie.ac.at/~stefanelli>

(Lara Trussardi) FACULTY OF MATHEMATICS, UNIVERSITY OF VIENNA, OSKAR-MORGENSTERN-PLATZ 1, 1090 WIEN, AUSTRIA.

*E-mail address:* [lara.trussardi@univie.ac.at](mailto:lara.trussardi@univie.ac.at)

*URL:* <http://www.mat.univie.ac.at/~trussardi>