

LARGE-TIME ASYMPTOTICS OF A FRACTIONAL DRIFT-DIFFUSION-POISSON SYSTEM VIA THE ENTROPY METHOD

FRANZ ACHLEITNER, ANSGAR JÜNGEL, AND MASAKAZU YAMAMOTO

ABSTRACT. The self-similar asymptotics for solutions to the drift-diffusion equation with fractional dissipation, coupled to the Poisson equation, is analyzed in the whole space. It is shown that in the subcritical and supercritical cases, the solutions converge to the fractional heat kernel with algebraic rate. The proof is based on the entropy method and leads to a decay rate in the $L^1(\mathbb{R}^d)$ norm. The technique is applied to other semilinear equations with fractional dissipation.

1. INTRODUCTION

In this paper, we investigate the large-time behavior of solutions to a drift-diffusion equation with fractional diffusion, coupled self-consistently to the Poisson equation. Such models describe the evolution of particles in a fluid under the influence of an acceleration field. The particle density $\rho(x, t)$ and potential $\psi(x, t)$ satisfy the equations

$$(1) \quad \partial_t \rho + (-\Delta)^{\theta/2} \rho = \operatorname{div}(\rho \nabla \psi), \quad -\Delta \psi = \rho \quad \text{in } \mathbb{R}^d, \quad t > 0,$$

with initial condition

$$(2) \quad \rho(\cdot, 0) = \rho_0 \quad \text{in } \mathbb{R}^d.$$

The fractional Laplacian $(-\Delta)^{\theta/2}$ is defined by $(-\Delta)^{\theta/2} \rho = \mathcal{F}^{-1}[|\xi|^\theta \mathcal{F}[\rho]]$, where \mathcal{F} is the Fourier transform, \mathcal{F}^{-1} its inverse, and $\theta > 0$. When $\theta = 2$, we recover the standard drift-diffusion-Poisson system arising in semiconductor theory and plasma physics [21]. Drift-diffusion-type equations with $\theta < 2$ were proposed to describe chemotaxis of biological cells whose behavior is not governed by Brownian motion [16]. For given acceleration field $\nabla \psi$ and $1 < \theta < 2$, the first equation in (1) was derived from the Boltzmann equation by Aceves-Sanchez and Mellet [1], based on the moment method of Mellet [28]. For the convenience of the reader, we present a formal derivation of the coupled system in Appendix B. Note, however, that we consider equations (1) in the range $0 < \theta \leq 2$.

Date: February 26, 2018.

2000 Mathematics Subject Classification. 35R11, 35B40, 35K45.

Key words and phrases. Drift-diffusion-Poisson system, fractional dissipation, self-similar asymptotics, large-time behavior.

The first and second author acknowledge support from the Austrian Science Fund (FWF), grant F65. The second author is supported by the Austrian Science Fund (FWF), grants P27352, P30000, and W1245. The last author is partially supported by the Japan Society for the Promotion of Science (JSPS), Grant-in-aid for Early-Career Scientists (B) 15K17566.

The aim of this paper is to compute the decay rate of the solution to (1) to self-similarity in the $L^1(\mathbb{R}^d)$ norm using the entropy method. In previous works [7, 26, 42], the self-similar asymptotics of various model variations were shown in $L^p(\mathbb{R}^d)$ norms but the decay rate is zero when $p = 1$. If additionally the first moment exists, i.e. if $|x|\rho_0 \in L^1(\mathbb{R}^d)$, the decay rate of the self-similar asymptotics in the $L^1(\mathbb{R}^d)$ norm is $1/\theta$, which is optimal [31, Lemma 5.1]. The entropy method provides an alternative way to analyze the self-similar asymptotics in the $L^1(\mathbb{R}^d)$ norm. Moreover, its strength is its robustness, i.e., the method can be easily applied to other semilinear equations with fractional dissipation. We give two examples in Section 4.

Before stating our main result and the key ideas of the technique, we review the state of the art for drift-diffusion equations. The *global existence of solutions* to the drift-diffusion-Poisson system was shown for $\theta = 2$ in [25], for the subcritical case $1 < \theta \leq 2$ in [26, 31], and for the supercritical case $0 < \theta < 1$ in [26, 35]. For suitable initial data, the solution to (1) satisfies

$$(3) \quad \rho \in C^0([0, \infty); L^1(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d)), \quad \rho(t) \geq 0 \quad \text{in } \mathbb{R}^d, \quad t > 0,$$

$$(4) \quad \|\rho(t)\|_{L^1(\mathbb{R}^d)} = \|\rho_0\|_{L^1(\mathbb{R}^d)}, \quad \|\rho(t)\|_{L^p(\mathbb{R}^d)} \leq C(1+t)^{-\frac{d}{\theta}(1-\frac{1}{p})}, \quad 1 \leq p \leq \infty;$$

see [31, Theorem 1.1] for $1 < \theta \leq 2$, [41, Theorem 1] for $\theta = 1$ and $d \geq 3$, and [26, Theorem 1.7] for $0 < \theta < 1$ and $d = 2$. The existence result of [31] for the bipolar drift-diffusion system was extended by Granero-Belinchón [18] by allowing for different fractional exponents. The existence of solutions in Besov spaces was proved for $1 < \theta < 2d$ and $d \geq 2$ in [44], and for $0 < \theta \leq 1$ and $d \geq 3$ in [35].

The self-similar asymptotics of the *fractional heat equation* was studied by Vazquez [37, Theorem 3.2], showing that if $0 < \theta < 2$ and the initial datum satisfies $(1+|x|)\rho_0 \in L^1(\mathbb{R}^d)$, we have $\|\rho(t) - MG_\theta(t)\|_{L^1(\mathbb{R}^d)} \leq Ct^{-1/\theta}$ with optimal rate, where $M = \int_{\mathbb{R}^d} \rho_0 dx$ is the initial mass and G_θ is the fundamental solution to the fractional heat equation (see Section 2.1). Exploiting the self-similar structure, this proves the exponential decay with rate $1/\theta$ to the fractional Fokker-Planck equation with quadratic potential. The exponential decay in L^1 spaces with weight $1+|x|^k$ and $k < \theta$ was proved by Tristani [36]. The fractional Laplacian can be replaced by more general Lévy operators, and the large-time asymptotics of so-called Lévy-Fokker-Planck equations were investigated by Biler and Karch [6] as well as Gentil and Imbert [17].

The large-time behavior of solutions to *drift-diffusion-Poisson systems* with $d = 2$ and $\theta = 2$ was studied by Nagai [30], showing the decay of the solutions to zero. A similar result for $0 < \theta < 2$ was proven by Li et al. [26]. The self-similar asymptotics in $L^p(\mathbb{R}^d)$ with $1 \leq p \leq \infty$ was shown in [23] for $\theta = 2$ and in [31] for $1 < \theta < 2$. In the latter reference, also the decay of the first-order asymptotic expansion of the solutions was computed. Higher-order expansions were studied for $1 < \theta \leq 2$ in [40], for $0 < \theta \leq 1$ in [42], and for the critical case $\theta = 1$ in [43]. However, in most of these references, the decay rate for $p = 1$ is zero.

The exponential decay in the relative entropy for solutions to Lévy-Fokker-Planck equations was proved in [6, 17]. Via the Csiszár-Kullback inequality (see, e.g., [3]), this implies

decay in the $L^1(\mathbb{R}^d)$ norm. In fact, we are using the techniques of [17], combined with tools from harmonic analysis and semigroup theory, to achieve self-similar decay of solutions to (1).

Our main result is as follows.

Theorem 1. *Let $d \geq 2$ and $0 < \theta \leq 2$ with $\theta < d/2$. Let $\rho_0 \in L^1(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d)$ be nonnegative such that $|x|^d \rho_0 \in L^q(\mathbb{R}^d)$ for some $q > d/\theta$. Furthermore, let ρ be a solution to (1) satisfying (3)-(4). Then, for all $t > 0$,*

$$(5) \quad \|\rho(t) - MG_\theta(t)\|_{L^1(\mathbb{R}^d)} \leq C(1 + \theta t)^{-1/2},$$

where $C > 0$ depends on ρ_0 and θ , $M = \int_{\mathbb{R}^d} \rho_0 dx$, and $G_\theta(x, t) = \mathcal{F}^{-1}[e^{-|\xi|^\theta t}](x)$ is the fundamental solution to $\partial_t u + (-\Delta)^{\theta/2} u = 0$ in \mathbb{R}^d .

Let us comment on the theorem. We need the finiteness of the moment $|x|^d \rho_0$ in $L^q(\mathbb{R}^d)$ to guarantee the well-posedness of the entropy functional; see Lemma 10 and Step 4 in the proof of Theorem 1. The lower bound of q is rather natural since $|x|^d G_\theta \in L^q(\mathbb{R}^d)$ if and only if $q > d/\theta$. If ρ_0 grows like $(1+r)^{-\alpha}$ for large values of the radius $r = |x|$, the condition $|x|^d \rho_0 \in L^q(\mathbb{R}^d)$ for large values of q is only slightly stronger than $\rho_0 \in L^1(\mathbb{R}^d)$. Indeed, in the latter case, we need $\alpha > d$, while $\alpha > (1+1/q)d$ is required in the former case. The condition $\theta < d/2$ is needed to estimate the nonlinear drift term; see the proof of Lemma 9. The decay rate is not optimal. This may be due to the fact the first moment of the fundamental solution G_θ is finite for all $1 < \theta < 2$ but not for $0 < \theta < 1$. The derivation of (1) leads to a drift term $\operatorname{div}(\rho D \nabla \psi)$ involving the drift matrix $D \in \mathbb{R}^{d \times d}$. We explain in Appendix C that we are able to treat only the case when D equals the identity matrix (times a factor and up to adding a skew-symmetric matrix).

As already mentioned, the idea of the proof is to employ the entropy method, originally developed for stochastic processes by Bakry and Emery [5] and later extended to linear and nonlinear diffusion equations (see, e.g., [3, 8]). First, we reformulate (1) in terms of the rescaled function

$$u(x, t) = \frac{e^{dt}}{M} \rho \left(e^t x, \frac{e^{\theta t} - 1}{\theta} \right), \quad x \in \mathbb{R}^d, \quad t > 0.$$

This function solves a drift-diffusion-Poisson system with the confinement potential $V(x) = \frac{1}{2}|x|^2$ and with the nonnegative steady state $u_\infty = G_\theta(1/\theta)$. Next, we show that the relative entropy

$$E_p \left[\frac{u}{u_\infty} \right] = \int_{\mathbb{R}^d} \left(\frac{u}{u_\infty} \right)^p u_\infty dx - \left(\int_{\mathbb{R}^d} u dx \right)^p$$

satisfies the inequality

$$\frac{dE_p}{dt} \leq -(\theta + \sigma(t))E_p + \sigma(t)$$

for some function $\sigma(t)$ which comes from the drift term involving ψ and which decays to zero exponentially fast. For this result, we need some results for Lévy operators due to [17] and a modified logarithmic Sobolev inequality due to Wu [39] and Chafaï [9]. By Gronwall's lemma, we conclude the exponential convergence of $t \mapsto E_p[u(t)/u_\infty]$. Then

the Csiszár-Kullback inequality implies that $u(t) - u_\infty$ converges exponentially fast in the $L^1(\mathbb{R}^d)$ norm. Finally, scaling back to the original variable, we deduce the algebraic decay for $\rho(t) - MG_\theta(t)$ in the $L^1(\mathbb{R}^d)$ norm.

The strength of the entropy method is that it is quite robust. It can be applied to other equations with fractional dissipation, at least if the regularity and decay properties (3)-(4) hold. As examples, we consider the two-dimensional quasi-geostrophic equation and a generalized fractional Burgers equation in one space dimension; see Section 4.

The paper is organized as follows. We summarize some results on the fractional heat equation and Lévy operators in Section 2. The proof of Theorem 1 is given in Section 3. In Section 4, the entropy method is applied to other equations. In Appendix A, a weighted $L^q(\mathbb{R}^d)$ estimate for ρ is shown. Appendix B is concerned with the formal derivation of (1), summarizing the ideas of [1]. Finally, we explain in Appendix C that we can only treat drift matrices of the form $D = aI + B$, where $a > 0$, I is the unit matrix, and B is a skew-symmetric matrix.

2. PRELIMINARIES

We need two ingredients for our analysis. The first one are properties of the fractional heat kernel, the second one is a modified logarithmic Sobolev inequality which relates the entropy E_p and the entropy production $-dE_p/dt$. For the convenience of the reader, we collect the needed results.

2.1. Fractional heat equation. We consider the fractional heat equation

$$(6) \quad \partial_t u + (-\Delta)^{\theta/2} u = 0 \quad \text{in } \mathbb{R}^d, \quad t > 0,$$

for $0 < \theta \leq 2$. The operator $(-\Delta)^{\theta/2}$ is defined for $\theta > 0$ and functions u in the Schwartz space of rapidly decaying functions on \mathbb{R}^d via Fourier transformation by $(-\Delta)^{\theta/2} u = \mathcal{F}^{-1}[|\xi|^\theta \mathcal{F}[u]]$, where $\mathcal{F}[u](x) = (2\pi)^{-d/2} \int_{\mathbb{R}^d} u(\xi) e^{-ix \cdot \xi} d\xi$ is the Fourier transform of u , and \mathcal{F}^{-1} is its inverse. In the limit $\theta \rightarrow 2$, the Laplace operator $-\Delta$ is recovered, but for $\theta \neq 2$, $(-\Delta)^{\theta/2}$ is a nonlocal operator. The fractional Laplacian can be expressed as the singular integral operator [15]

$$(-\Delta)^{\theta/2} u = -c_{d,\theta} \text{p.v.} \int_{\mathbb{R}^d} \frac{u(x+y) - u(x)}{|y|^{d+\theta}} dy, \quad c_{d,\theta} = \frac{\theta}{2\pi^{d/2+\theta}} \frac{\Gamma(\frac{1}{2}(d+\theta))}{\Gamma(\frac{1}{2}(2-\theta))},$$

where u is a suitable function, p.v. denotes the Cauchy principal value, and Γ is the Gamma function. The principal value can be avoided for $0 < \theta < 2$, and the integral becomes a standard one when $0 < \theta < 1$ [15, Theorem 1].

The fundamental solution G_θ of the fractional heat equation (6) is given by

$$(7) \quad G_\theta(x, t) = \mathcal{F}^{-1}[e^{-t|\xi|^\theta}](x), \quad x \in \mathbb{R}^d, \quad t > 0.$$

We recall some qualitative properties of G_θ :

- Normalization: $\int_{\mathbb{R}^d} G_\theta(x, t) dx = 1$ for $t > 0$.
- Self-similar form: A direct computation shows that

$$(8) \quad \lambda^d G_\theta(\lambda x, \lambda^\theta t) = G_\theta(x, t) \quad \text{for all } x \in \mathbb{R}^d, \quad t > 0, \quad \lambda > 0.$$

- Pointwise estimates [24, Theorem 7.3.1]: For any $K > 0$, there exists a constant $C > 1$, which can be chosen uniformly for θ from any compact interval in $(0, 2)$, such that

$$(9) \quad \begin{aligned} \frac{1}{Ct^{d/\theta}} &\leq G_\theta(x, t) \leq \frac{C}{t^{d/\theta}} && \text{if } |x| \leq Kt^{1/\theta}, t > 0, \\ \frac{t}{C|x|^{d+\theta}} &\leq G_\theta(x, t) \leq \frac{Ct}{|x|^{d+\theta}} && \text{if } |x| \geq Kt^{1/\theta}, t > 0. \end{aligned}$$

- Gradient estimate [24, Theorem 7.3.2]: There exists $C > 0$ such that

$$(10) \quad |\nabla G_\theta(x, t)| \leq C \min \{t^{-1/\theta}, |x|^{-1}\} G_\theta(x, t), \quad x \in \mathbb{R}^d, t > 0,$$

holds uniformly for θ from any compact interval in $(0, 2)$.

Here and in the following, $C > 0$ denotes a generic constant independent of x and t .

2.2. Lévy operators. We recall some results for Lévy operators. A Lévy operator \mathcal{I} is the infinitesimal generator associated with a Lévy process. According to the Lévy-Khinchine formula [33], it can be written in the form

$$\mathcal{I}[u] = \operatorname{div}(\sigma \nabla u) - b \cdot \nabla u + \int_{\mathbb{R}^d} (u(x+y) - u(x) - \nabla u(x) \cdot yh(y)) \nu(dy),$$

defined for suitable functions u , where σ is a symmetric positive semidefinite $d \times d$ matrix, $b \in \mathbb{R}^d$, $h(y) = (1 + |y|^2)^{-1}$ is a truncation function, and ν denotes a nonnegative singular measure on \mathbb{R}^d satisfying

$$\nu(\{0\}) = 0 \quad \text{and} \quad \lim_{\varepsilon \rightarrow 0} \int_{\{|y| > \varepsilon\}} \min\{1, |y|^2\} \nu(dy) < \infty;$$

see, e.g., [17, Section 1]. Fractional Laplacians $-(-\Delta)^{\theta/2}$ for $0 < \theta < 2$ are Lévy operators with $\sigma = 0$, $b = 0$, and Lévy measure $\nu(dy) = dy/|y|^{d+\theta}$. For more details on Lévy operators, we refer to [2, 33].

Let $\Phi : \mathbb{R}_+ \rightarrow \mathbb{R}$ be a smooth convex function and u_∞ a positive function satisfying $\int_{\mathbb{R}^d} u_\infty dx = 1$. The Φ -entropy is defined by

$$\operatorname{Ent}_{u_\infty}^\Phi(f) = \int_{\mathbb{R}^d} \Phi(f) u_\infty dx - \Phi\left(\int_{\mathbb{R}^d} f u_\infty dx\right),$$

for non-negative functions f . Observe that Jensen's inequality implies that $\operatorname{Ent}_{u_\infty}^\Phi(f) \geq 0$.

We need two results for the Φ -entropy. In the following we only consider the family of smooth convex functions $\Phi_p : \mathbb{R}_+ \rightarrow \mathbb{R}$, $s \mapsto s^p$, with $p \in (1, 2]$. However, the results hold for any smooth convex function Φ such that the mappings $(a, b) \mapsto D_\Phi(a+b, b)$ and $(a, y) \mapsto \Phi''(a)y \cdot (\sigma y)$ are convex on $\{a+b \geq 0, b \geq 0\}$ and $\mathbb{R}_+ \times \mathbb{R}^{2d}$, respectively. Here, $D_\Phi(a, b) := \Phi(a) - \Phi(b) - \Phi'(b)(a-b)$ denotes the Bregman distance.

The first result is a modified logarithmic Sobolev inequality.

Proposition 2. *Let $p \in (1, 2]$ and consider $\Phi : \mathbb{R}_+ \rightarrow \mathbb{R}$, $s \mapsto s^p$. If u_∞ is the density of an infinitely divisible probability measure then for all smooth positive functions f*

$$(11) \quad \text{Ent}_{u_\infty}^\Phi(f) \leq \int_{\mathbb{R}^d} \Phi''(f) \nabla f \cdot (\sigma_\infty \nabla f) u_\infty \, dx + \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} D_\Phi(f(x), f(x+y)) \nu_\infty(dy) u_\infty \, dx,$$

where ν_∞ and σ_∞ are the Lévy measure and the diffusion matrix associated with u_∞ , respectively.

This result was first proved for particular cases by Wu [39] and generalized by Chafaï [9], see also [17, Theorem 2].

The second result is a formula for the time derivative of the Φ -entropy along solutions to the Lévy-Fokker-Planck equation

$$(12) \quad \partial_t u = \mathcal{I}[u] + \text{div}(xu) \quad \text{for } x \in \mathbb{R}^d, \quad t > 0,$$

see [17, Proposition 1]. In fact, we need this result only for the special case of fractional Laplacians $\mathcal{I} = -(-\Delta)^{\theta/2}$ with $0 < \theta < 2$.

Proposition 3 ([17, Theorem 1]). *Let $p \in (1, 2]$ and consider $\Phi : \mathbb{R}_+ \rightarrow \mathbb{R}$, $s \mapsto s^p$. Consider the Lévy-Fokker-Planck equation (12) for fractional Laplacians $\mathcal{I} = -(-\Delta)^{\theta/2}$ with $0 < \theta < 2$ and stationary solution $u_\infty(x) = G_\theta(x, 1/\theta)$. If u_0 is a nonnegative function with $\text{Ent}_{u_\infty}^\Phi(u_0/u_\infty) < \infty$, then the solution u of (12) with initial datum u_0 satisfies for all $t \geq 0$,*

$$\text{Ent}_{u_\infty}^\Phi(u(t)/u_\infty) \leq e^{-\theta t} \text{Ent}_{u_\infty}^\Phi(u_0/u_\infty).$$

Proof. We claim that $u_\infty(x) = G_\theta(x, 1/\theta)$ is the stationary solution of (12), i.e. $\mathcal{I}[u_\infty] + \text{div}(xu_\infty) = 0$. Indeed, the Fourier transform of this equation is $|\xi|^\theta \widehat{u}_\infty + \xi \cdot \nabla \widehat{u}_\infty = 0$, whose solution is given by $\widehat{u}_\infty(\xi) = e^{-|\xi|^\theta/\theta}$, and the claim follows from the definition of the fractional heat kernel. Moreover, $u_\infty(x) = G_\theta(x, 1/\theta)$ is the density of an infinitely divisible probability measure with $\sigma_\infty = 0$ and Lévy measure $\nu_\infty(dy) = dy/\theta|y|^{d+\theta}$. Then, for $v = u/u_\infty$,

$$(13) \quad \begin{aligned} \frac{d}{dt} \text{Ent}_{u_\infty}^\Phi(v) &= - \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} D_\Phi(v(x), v(x-y)) \nu(dy) u_\infty(x) \, dx \\ &= - \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} D_\Phi(v(x), v(x+y)) \nu(dy) u_\infty(x) \, dx \\ &\leq -\theta \text{Ent}_{u_\infty}^\Phi(v). \end{aligned}$$

The first identity follows from [17, Proposition 1], in the second one we substituted $y \mapsto -y$ (and used $\nu(-dy) = \nu(dy)$), and the inequality is a consequence of the logarithmic Sobolev inequality (11). The factor θ comes from the relation between the Lévy measures ν and ν_∞ , see the comment after Theorem 1 in [17]. The final statement follows from Gronwall's inequality. \square

Lemma 4. *Let $0 < \theta \leq 2$ and G_θ be the fundamental solution to the fractional heat equation (6). Then for all $s > 0$,*

$$\|G_\theta(s+1/\theta) - G_\theta(s)\|_{L^1(\mathbb{R}^d)} \leq C(1+\theta s)^{-1/2}.$$

Proof. A direct computation shows that the function $U(x, t) = e^{dt}G_\theta(e^t x, (e^{\theta t} - 1)/\theta)$ solves the fractional Fokker-Planck equation

$$\partial_t U + (-\Delta)^{\theta/2} U = \operatorname{div}(xU) \quad \text{in } \mathbb{R}^d,$$

and $U_\infty = G_\theta(1/\theta)$ is the stationary solution. Setting $s = (e^{\theta t} - 1)/\theta$ and using the substitution $y = e^{-t}x$ and the self-similar form (8) with $\lambda = e^t$, we find that

$$\begin{aligned} \|G_\theta(s + 1/\theta) - G_\theta(s)\|_{L^1(\mathbb{R}^d)} &= \int_{\mathbb{R}^d} |G_\theta(x, e^{\theta t}/\theta) - G_\theta(x, (e^{\theta t} - 1)/\theta)| dx \\ &= \int_{\mathbb{R}^d} |G_\theta(e^t y, e^{\theta t}/\theta) - G_\theta(e^t y, (e^{\theta t} - 1)/\theta)| e^{dt} dy \\ &= \int_{\mathbb{R}^d} |G_\theta(y, 1/\theta) - e^{dt}G_\theta(e^t y, (e^{\theta t} - 1)/\theta)| dy \\ &= \|U_\infty - U(t)\|_{L^1(\mathbb{R}^d)}. \end{aligned}$$

Using Proposition 3 and the Csiszár-Kullback inequality (see [3] or [22, Theorem A.3])

$$\|U_\infty - U(t)\|_{L^1(\mathbb{R}^d)} \leq C e^{-\theta(t-t_0)/2} = C e^{-\theta t_0/2} (1 + \theta s)^{-1/2},$$

and the constant $C > 0$ depends on $\operatorname{Ent}_{U_\infty}^\Phi [U(t_0)/U_\infty]$ for some small $t_0 > 0$. This finishes the proof. \square

Remark 5. A direct estimate allows us to prove

$$(14) \quad \|G_\theta(s + 1/\theta) - G_\theta(s)\|_{L^1(\mathbb{R}^d)} \leq C(1 + s)^{-1}.$$

However, the constant C is of order $1/\theta^2$, hence, it is not uniformly bounded for $\theta \in (0, 2)$. In order to show (14), we observe that the integrand of

$$G_\theta(s + 1/\theta) - G_\theta(s) = \frac{1}{\theta} \int_0^1 \partial_t G_\theta(s + \lambda/\theta) d\lambda$$

can be written for $t = s + \lambda/\theta$ as

$$\begin{aligned} \partial_t G_\theta(x, t) &= \partial_t (t^{-d/\theta} G_\theta(t^{-1/\theta} x, 1)) \\ &= -\frac{1}{\theta t} \left(d t^{-d/\theta} G_\theta(t^{-1/\theta} x, 1) - t^{-d/\theta} (t^{-1/\theta} x) \cdot \nabla G_\theta(t^{-1/\theta} x, 1) \right), \end{aligned}$$

using the self-similar property (8). By the pointwise bounds (9) and (10), we find that

$$\|G_\theta(s + 1/\theta) - G_\theta(s)\|_{L^1(\mathbb{R}^d)} \leq C \int_0^1 (s + \lambda/\theta)^{-1} d\lambda \leq C \int_0^1 s^{-1} d\lambda = C s^{-1}.$$

Taking into account the bound

$$\|G_\theta(s + 1/\theta) - G_\theta(s)\|_{L^1(\mathbb{R}^d)} \leq \|G_\theta(s + 1/\theta)\|_{L^1(\mathbb{R}^d)} + \|G_\theta(s)\|_{L^1(\mathbb{R}^d)} \leq 2,$$

the claim (14) follows.

3. PROOF OF THE MAIN RESULT

We split the proof of Theorem 1 into several steps.

Step 1: Time-dependent rescaling of the equation. Let $M = \int_{\mathbb{R}^d} \rho_0 \, dx > 0$. We introduce the rescaled function

$$(15) \quad u(x, t) = \frac{e^{dt}}{M} \rho \left(e^t x, \frac{e^{\theta t} - 1}{\theta} \right), \quad x \in \mathbb{R}^d, \quad t > 0.$$

Lemma 6. *The function u solves the confined drift-diffusion-Poisson system*

$$(16) \quad \begin{aligned} \partial_t u + (-\Delta)^{\theta/2} u &= \operatorname{div}(u \nabla V) + M e^{-(d-\theta)t} \operatorname{div}(u \nabla \phi) \quad \text{in } \mathbb{R}^d, \quad t > 0, \\ -\Delta \phi &= u, \quad u(\cdot, 0) = u_0 \quad \text{in } \mathbb{R}^d, \end{aligned}$$

where $V(x) = \frac{1}{2}|x|^2$ and $u_0 = \rho_0/M$.

Proof. First, we observe that u fulfills

$$(17) \quad \partial_t u - \operatorname{div}(u \nabla V) = \frac{1}{M} e^{(d+\theta)t} \partial_t \rho \left(e^t x, \frac{e^{\theta t} - 1}{\theta} \right)$$

and, substituting $y = e^t x$,

$$\begin{aligned} \widehat{u}(\xi, t) &= \frac{1}{M} (2\pi)^{-d/2} \int_{\mathbb{R}^d} e^{dt} \rho \left(e^t x, \frac{e^{\theta t} - 1}{\theta} \right) e^{-ix \cdot \xi} \, dx \\ &= \frac{1}{M} (2\pi)^{-d/2} \int_{\mathbb{R}^d} \rho \left(y, \frac{e^{\theta t} - 1}{\theta} \right) e^{-iy \cdot (\exp(-t)\xi)} \, dy \\ &= \frac{1}{M} \widehat{\rho} \left(e^{-t}\xi, \frac{e^{\theta t} - 1}{\theta} \right). \end{aligned}$$

With this expression and the substitution $\eta = e^{-t}\xi$, we find that

$$(18) \quad \begin{aligned} (-\Delta)^{\theta/2} u &= (2\pi)^{-d/2} \int_{\mathbb{R}^d} |\xi|^\theta \widehat{u}(\xi, t) e^{ix \cdot \xi} \, d\xi \\ &= \frac{1}{M} (2\pi)^{-d/2} \int_{\mathbb{R}^d} |\xi|^\theta \widehat{\rho} \left(e^{-t}\xi, \frac{e^{\theta t} - 1}{\theta} \right) e^{ix \cdot \xi} \, d\xi \\ &= \frac{1}{M} (2\pi)^{-d/2} e^{(d+\theta)t} \int_{\mathbb{R}^d} |\eta|^\theta \widehat{\rho} \left(\eta, \frac{e^{\theta t} - 1}{\theta} \right) e^{i(\exp(t)x) \cdot \eta} \, d\eta \\ &= \frac{1}{M} e^{(d+\theta)t} (-\Delta)^{\theta/2} \rho \left(e^t x, \frac{e^{\theta t} - 1}{\theta} \right). \end{aligned}$$

Adding (17)-(18) and inserting (1) leads to

$$(19) \quad \begin{aligned} \partial_t u + (-\Delta)^{\theta/2} u - \operatorname{div}(u \nabla V) &= \frac{1}{M} e^{(d+\theta)t} \operatorname{div}(\rho \nabla \psi) \\ &= \frac{1}{M} e^{(d+\theta)t} (\nabla \rho \cdot \nabla \psi) \left(e^t x, \frac{e^{\theta t} - 1}{\theta} \right) - \frac{1}{M} e^{(d+\theta)t} \rho^2 \left(e^t x, \frac{e^{\theta t} - 1}{\theta} \right). \end{aligned}$$

It remains to express the right-hand side in terms of u .

By the representation formula for solutions of the Poisson equation $-\Delta\phi = u$ and the substitution $z = e^t y$, it follows that

$$(20) \quad \begin{aligned} \nabla\phi(x) &= c_d \int_{\mathbb{R}^d} \frac{x-y}{|x-y|^d} u(y,t) dy = \frac{c_d}{M} e^{dt} \int_{\mathbb{R}^d} \frac{x-y}{|x-y|^d} \rho\left(e^t y, \frac{e^{\theta t} - 1}{\theta}\right) dy \\ &= \frac{c_d}{M} e^{(d-1)t} \int_{\mathbb{R}^d} \frac{e^t x - z}{|e^t x - z|^d} \rho\left(z, \frac{e^{\theta t} - 1}{\theta}\right) dz = \frac{1}{M} e^{(d-1)t} \nabla\psi\left(e^t x, \frac{e^{\theta t} - 1}{\theta}\right), \end{aligned}$$

where $c_d = \Gamma(d/2)/(2\pi^{d/2})$. Hence, since

$$\nabla u(x,t) = \frac{1}{M} e^{(d+1)t} \nabla\rho\left(e^t x, \frac{e^{\theta t} - 1}{\theta}\right),$$

the first term on the right-hand side of (19) becomes

$$\frac{1}{M} e^{(d+\theta)t} (\nabla\rho \cdot \nabla\psi)\left(e^t x, \frac{e^{\theta t} - 1}{\theta}\right) = M e^{-(d-\theta)t} (\nabla u \cdot \nabla\phi)(x,t).$$

Taking the square of the definition of u , the second term on the right-hand side of (19) can be written as

$$\frac{1}{M} e^{(d+\theta)t} \rho^2\left(e^t x, \frac{e^{\theta t} - 1}{\theta}\right) = M e^{-(d-\theta)t} u^2(x,t).$$

Inserting the previous two expressions in (19) and using $\nabla u \cdot \nabla\phi - u^2 = \operatorname{div}(u\nabla\phi)$, we finish the proof. \square

Lemma 7. *The solution u to (16) satisfies $u \in L^\infty(0, \infty; L^1(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d))$.*

Proof. We use the estimate in (4) and the definition of u to find that

$$\begin{aligned} \|u(t)\|_{L^p(\mathbb{R}^d)}^p &= \frac{e^{dpt}}{M^p} \int_{\mathbb{R}^d} \rho\left(e^t x, \frac{e^{\theta t} - 1}{\theta}\right)^p dx = \frac{e^{dt(p-1)}}{M^p} \int_{\mathbb{R}^d} \rho\left(y, \frac{e^{\theta t} - 1}{\theta}\right)^p dy \\ &\leq \frac{C}{M^p} e^{dt(p-1)} \left(1 + \frac{e^{\theta t} - 1}{\theta}\right)^{-d(p-1)/\theta} \leq \frac{C}{M^p} \left(\frac{\theta e^{\theta t}}{\theta + e^{\theta t} - 1}\right)^{d(p-1)/\theta} \leq \frac{C}{M^p}. \end{aligned}$$

Consequently, $u(t)$ is bounded in any L^p norm uniformly in time and we conclude by passing to the limit $p \rightarrow \infty$. \square

Step 2: Time decay of $u(t)$. We show that u converges exponentially fast to the fundamental solution $G_\theta(1/\theta)$ (see Section 2.1). First, we relate the difference of ρ to the fundamental solution G_θ and the difference of u to G_θ .

Lemma 8. *Let $s = (e^{\theta t} - 1)/\theta$. Then it holds that*

$$\|\rho(s) - M G_\theta(s + 1/\theta)\|_{L^1(\mathbb{R}^d)} = M \|u(t) - G_\theta(1/\theta)\|_{L^1(\mathbb{R}^d)}.$$

Proof. By the definition of $u(x,t)$ and the substitution $y = e^t x$, we compute

$$M \|u(t) - G_\theta(1/\theta)\|_{L^1(\mathbb{R}^d)} = \int_{\mathbb{R}^d} \left| e^{dt} \rho\left(e^t x, \frac{e^{\theta t} - 1}{\theta}\right) - M G_\theta(x, 1/\theta) \right| dx$$

$$= \int_{\mathbb{R}^d} \left| \rho \left(y, \frac{e^{\theta t} - 1}{\theta} \right) - M e^{-dt} G_\theta(e^{-t} y, 1/\theta) \right| dy.$$

We set $s = (e^{\theta t} - 1)/\theta$ and use the self-similar form (8) with $\lambda = e^{-t} = (1 + s\theta)^{-1/\theta}$:

$$\begin{aligned} M \|u(t) - G_\theta(1/\theta)\|_{L^1(\mathbb{R}^d)} &= \int_{\mathbb{R}^d} \left| \rho(x, s) - M(1 + s\theta)^{-d/\theta} G_\theta((1 + s\theta)^{-1/\theta} x, 1/\theta) \right| dx \\ &= \int_{\mathbb{R}^d} \left| \rho(x, s) - M G_\theta(x, (1 + s\theta)/\theta) \right| dx \\ &= \|\rho(s) - M G_\theta(s + 1/\theta)\|_{L^1(\mathbb{R}^d)}, \end{aligned}$$

which concludes the proof. \square

The previous lemma shows that it is sufficient to estimate $u(t) - G_\theta(1/\theta)$.

Lemma 9. *Let the assumptions of Theorem 1 hold. Then the solution u to (16) satisfies*

$$\|u(t) - G_\theta(1/\theta)\|_{L^1(\mathbb{R}^d)} \leq C e^{-\theta t/2}, \quad t > 0,$$

where $C > 0$ is some constant.

Proof. Let $1 < p < 2$ and set $v(t) = u(t)/G_\theta(1/\theta)$ and

$$(21) \quad E_p[v(t)] = \int_{\mathbb{R}^d} v(x, t)^p G_\theta(x, 1/\theta) dx - \left(\int_{\mathbb{R}^d} v(x, t) G_\theta(x, 1/\theta) dx \right)^p.$$

We prove in Step 4 below that this functional is well defined. Note that $E_p[v] = \text{Ent}_{u_\infty}^\Phi(v)$, where $\Phi(s) = s^p$ and $\text{Ent}_{u_\infty}^\Phi$ is the entropy defined in Section 2.2. We differentiate $E_p[v]$ with respect to time. The derivative of the second integral in $E_p[v]$ vanishes since $\int_{\mathbb{R}^d} v G_\theta(1/\theta) dx = \int_{\mathbb{R}^d} u dx = 1$. Therefore, by (16),

$$\begin{aligned} \frac{dE_p}{dt}[v(t)] &= p \int_{\mathbb{R}^d} v(x, t)^{p-1} \partial_t u(x, t) dx \\ &= p \int_{\mathbb{R}^d} v^{p-1} \left(-(-\Delta)^{\theta/2} u + \text{div}(u \nabla V) \right) dx + p M e^{-(d-\theta)t} \int_{\mathbb{R}^d} v^{p-1} \text{div}(u \nabla \phi) dx. \end{aligned}$$

Using the Lévy operator $\mathcal{I}[u] = \text{div}(x \nabla V) - (-\Delta)^{\theta/2}$ with $V(x) = \frac{1}{2}|x|^2$, the calculations in the proof of [17, Proposition 1] show that

$$\begin{aligned} &p \int_{\mathbb{R}^d} v^{p-1} \left(-(-\Delta)^{\theta/2} u + \text{div}(u \nabla V) \right) dx \\ &= - \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} D_p(v(x, t), v(x + y, t)) \nu(dy) G_\theta(x, 1/\theta) dx, \end{aligned}$$

where $D_p(a, b) = a^p + b^p - p b^{p-1}(a - b)$ is the Bregman distance (see Section 2.2) and $\nu(dy) = dy/|y|^{d+\theta}$. Moreover, the modified logarithmic Sobolev inequality (11) gives

$$\int_{\mathbb{R}^d} \int_{\mathbb{R}^d} D_p(v(x, t), v(x + y, t)) \nu(dy) G_\theta(x, 1/\theta) dx \geq \theta E_p[v],$$

since $u_\infty(x) = G_\theta(x, 1/\theta)$ is the density of an infinitely divisible probability measure with $\sigma_\infty = 0$ and Lévy measure $\nu_\infty(dy) = dy/\theta|y|^{d+\theta}$. Putting these estimates together leads to

$$(22) \quad \frac{dE_p}{dt}[v] \leq -\theta E_p[v] + pM e^{-(d-\theta)t} \int_{\mathbb{R}^d} v^{p-1} \operatorname{div}(v G_\theta(1/\theta) \nabla \phi) dx,$$

and it remains to estimate the last integral. To this end, we differentiate and integrate by parts:

$$\begin{aligned} p \int_{\mathbb{R}^d} v^{p-1} \operatorname{div}(v G_\theta(1/\theta) \nabla \phi) dx &= \int_{\mathbb{R}^d} (p v^p \operatorname{div}(G_\theta(1/\theta) \nabla \phi) + \nabla(v^p) \cdot (G_\theta(1/\theta) \nabla \phi)) dx \\ &= (p-1) \int_{\mathbb{R}^d} v^p \operatorname{div}(G_\theta(1/\theta) \nabla \phi) dx. \end{aligned}$$

Differentiating $G_\theta(1/\theta) \nabla \phi$ and using $-\Delta \phi = u$, we obtain

$$\begin{aligned} &\left| \int_{\mathbb{R}^d} v^p \operatorname{div}(G_\theta(1/\theta) \nabla \phi) dx \right| \\ &\leq \int_{\mathbb{R}^d} |v^p \nabla \log G_\theta(1/\theta) \cdot \nabla \phi| G_\theta(1/\theta) dx + \int_{\mathbb{R}^d} v^p u G_\theta(1/\theta) dx \\ &\leq (\|\nabla \log G_\theta(1/\theta)\|_{L^\infty(\mathbb{R}^d)} \|\nabla \phi\|_{L^\infty(\mathbb{R}^d)} + \|u\|_{L^\infty(\mathbb{R}^d)}) \int_{\mathbb{R}^d} v^p G_\theta(1/\theta) dx. \end{aligned}$$

We claim that the L^∞ norms are bounded uniformly in time. Indeed, by Lemma 7, the L^∞ norm of $u(t)$ is uniformly bounded. By (10), we find that

$$|\nabla \log G_\theta(1/\theta)| = \frac{|\nabla G_\theta(1/\theta)|}{G_\theta(1/\theta)} \leq C \min\{\theta^{1/\theta}, |x|^{-1}\} \leq C,$$

which shows the bound for $\|\nabla \log G_\theta(1/\theta)\|_{L^\infty(\mathbb{R}^d)}$. Finally, we deduce from Poisson's representation formula that

$$\begin{aligned} |\nabla \phi(x, t)| &\leq C \int_{\mathbb{R}^d} \frac{|u(y, t)|}{|x-y|^{d-1}} dy \\ &= C \int_{\{|x-y|<1\}} \frac{|u(y, t)|}{|x-y|^{d-1}} dy + C \int_{\{|x-y|\geq 1\}} \frac{|u(y, t)|}{|x-y|^{d-1}} dy \\ &\leq C \|u(t)\|_{L^\infty(\mathbb{R}^d)} \int_{\{|x-y|<1\}} \frac{dy}{|x-y|^{d-1}} + C \int_{\mathbb{R}^d} |u(y, t)| dy \\ &\leq C (\|u(t)\|_{L^\infty(\mathbb{R}^d)} + \|u(t)\|_{L^1(\mathbb{R}^d)}), \end{aligned}$$

and we have already seen that the right-hand side is bounded, due to Lemma 7. Hence, (22) becomes

$$\frac{dE_p}{dt}[v] \leq -\theta E_p[v] + (p-1)C M e^{-(d-\theta)t} \int_{\mathbb{R}^d} v^p G_\theta(1/\theta) dx,$$

where $C := \sup_{0<t<\infty} (\|\nabla \log G_\theta(1/\theta)\|_{L^\infty(\mathbb{R}^d)} \|\nabla \phi\|_{L^\infty(\mathbb{R}^d)} + \|u\|_{L^\infty(\mathbb{R}^d)})$. By definition of the entropy $E_p[v]$ and the mass conservation $\int_{\mathbb{R}^d} u dx = 1$, the integral on the right-hand side

equals

$$\int_{\mathbb{R}^d} v^p G_\theta(1/\theta) \, dx = E_p[v] + \left(\int_{\mathbb{R}^d} v G_\theta \, dx \right)^p = E_p[v] + 1,$$

and we end up with

$$\frac{dE_p}{dt}[v] \leq (-\theta + \sigma(t))E_p[v] + \sigma(t), \quad \text{where } \sigma(t) = (p-1)CM e^{-(d-\theta)t}.$$

We apply the Gronwall inequality to infer that

$$E_p[v(t)] \leq e^{-\theta t + S(t)} \left(E_p[v(0)] + \int_0^t e^{\theta s - S(s)} \sigma(s) \, ds \right),$$

where $S(t) = \int_0^t \sigma(s) \, ds$ is bounded uniformly in $t \in (0, \infty)$ since $d - \theta > 0$. It follows from $e^{\theta s} \sigma(s) = (p-1)CM e^{(2\theta-d)s}$ and $2\theta - d < 0$ that, for some constants $C > 0$,

$$E_p[v(t)] \leq C e^{-\theta t} \left(E_p[v(0)] + \int_0^t e^{(2\theta-d)s} \, ds \right) \leq C e^{-\theta t}.$$

Finally, we use the Csiszár-Kullback inequality (see [3] or [22, Theorem A.3]), applied to the function $z \mapsto z^p - 1$ for $1 < p < 2$,

$$\begin{aligned} \|u(t) - G_\theta(1/\theta)\|_{L^1(\mathbb{R}^d)}^2 &\leq \frac{2}{p(p-1)} \int_{\mathbb{R}^d} \left\{ \left(\frac{u}{G_\theta(1/\theta)} \right)^p - 1 \right\} G_\theta(1/\theta) \, dx \\ &= \frac{2}{p(p-1)} \int_{\mathbb{R}^d} (v^p - 1) G_\theta(1/\theta) \, dx \\ &= \frac{2}{p(p-1)} \left(\int_{\mathbb{R}^d} v^p G_\theta(1/\theta) \, dx - 1 \right) \\ &= \frac{2}{p(p-1)} E_p[v(t)] \leq C e^{-\theta t}, \end{aligned}$$

where we used the fact that $G_\theta(1/\theta)$ is normalized. □

Step 3: Time decay of $\rho(t)$. Lemmas 8 and 9 show that

$$\begin{aligned} \|\rho(s) - MG_\theta(s + 1/\theta)\|_{L^1(\mathbb{R}^d)} &= M \|u(t) - G_\theta(1/\theta)\|_{L^1(\mathbb{R}^d)} \\ &\leq C M e^{-\theta t/2} = C M (1 + \theta s)^{-1/2}, \end{aligned}$$

since $s = (e^{\theta t} - 1)/\theta$ is equivalent to $e^{\theta t} = 1 + \theta s$. We infer from Lemma 4 that

$$\begin{aligned} \|\rho(s) - MG_\theta(s)\|_{L^1(\mathbb{R}^d)} &\leq \|\rho(s) - MG_\theta(s + 1/\theta)\|_{L^1(\mathbb{R}^d)} \\ &\quad + M \|G_\theta(s + 1/\theta) - G_\theta(s)\|_{L^1(\mathbb{R}^d)} \\ &\leq C M (1 + \theta s)^{-1/2}. \end{aligned}$$

Step 4: Well-posedness of the entropy (21). It remains to verify that (21) is well defined. We reformulate $E_p[v]$ in terms of ρ . With the substitutions $y = e^t x$ and $s = (e^{\theta t} - 1)/\theta$,

we find from definition (15) that

$$\begin{aligned} E_p[v] &= \frac{e^{dpt}}{M^p} \int_{\mathbb{R}^d} \rho \left(e^t x, \frac{e^{\theta t} - 1}{\theta} \right)^p G_\theta(x, 1/\theta)^{1-p} dx - \frac{e^{dpt}}{M^p} \left(\int_{\mathbb{R}^d} \rho \left(e^t x, \frac{e^{\theta t} - 1}{\theta} \right) dx \right)^p \\ &= \frac{e^{d(p-1)t}}{M^p} \int_{\mathbb{R}^d} \rho(y, s)^p G_\theta(e^{-t}y, 1/\theta)^{1-p} dy - \frac{1}{M^p} \left(\int_{\mathbb{R}^d} \rho(y, s) dy \right)^p. \end{aligned}$$

Using the self-similar form (8) with $\lambda = e^{-t}$ yields

$$E_p[v] = \frac{1}{M^p} \int_{\mathbb{R}^d} \left(\frac{\rho(y, s)}{G_\theta(y, s + 1/\theta)} \right)^p G_\theta(y, s + 1/\theta) dy - \frac{1}{M^p} \left(\int_{\mathbb{R}^d} \rho(y, s) dy \right)^p.$$

Therefore, it is sufficient to show that the integral

$$(23) \quad I(t) := \int_{\mathbb{R}^d} \left(\frac{\rho(x, t)}{G_\theta(x, t + 1/\theta)} \right)^p G_\theta(x, t + 1/\theta) dx$$

converges for any fixed $t > 0$ and some $1 < p < 2$. The trivial estimate (taking into account (9))

$$I(t) \leq \|\rho(t)\|_{L^\infty(\mathbb{R}^d)}^p \int_{\mathbb{R}^d} G(x, t + 1/\theta)^{1-p} dx \leq Ct^{-d/\theta} \int_{\mathbb{R}^d} (1 + |x|)^{(d+\theta)(p-1)} dx$$

cannot be used since $p - 1 > 0$, so we have to derive a finer estimate for $\rho(t)$ in some $L^q(\mathbb{R}^d)$. This is done in the following lemma.

Lemma 10. *Let $d \geq 2$ and either $1 < \theta \leq 2$, $q > d/(d - \theta)$ or $0 < \theta \leq 1$, $q > d/\theta$. Furthermore, let $|x|^d \rho_0 \in L^q(\mathbb{R}^d)$, and the solution to (1) satisfies (3)-(4). Then*

$$\||x|^d \rho(t)\|_{L^q(\mathbb{R}^d)} \leq C(1+t)^{d/(\theta q)}, \quad t > 0.$$

We postpone the (lengthy) proof to Appendix A.

We claim that Lemma 10 and estimate (9) imply that (23) converges. To see this, we first observe that (23) converges for $p = 1$. For $p > 1$, we fix $t > 0$ and split (23) into two parts, $I(t) = I_1 + I_2$, where

$$\begin{aligned} I_1 &= \int_{B_R(0)} \rho(x, t)^p G_\theta(x, t + 1/\theta)^{1-p} dx, \\ I_2 &= \int_{\mathbb{R}^d \setminus B_R(0)} \rho(x, t)^p G_\theta(x, t + 1/\theta)^{1-p} dx, \end{aligned}$$

and $B_R(0)$ is the ball of radius $R > 0$ centered at the origin. Let $K = R(t + 1/\theta)^{-1/\theta}$. Consider $|x| < R = K(t + 1/\theta)^{1/\theta}$. Then we can apply estimates (9) and (4):

$$I_1 \leq C(t + 1/\theta)^{\frac{d}{\theta}(p-1)} \int_{\mathbb{R}^d} \rho(x, t)^p dx \leq C(t + 1/\theta)^{\frac{d}{\theta}(p-1)} t^{-\frac{d}{\theta}(p-1)} \leq C.$$

To estimate I_2 , we wish to apply Lemma 10 for large $q > d/\theta$. If $0 < \theta < 1$, the inequality $q > d/\theta$ shows that the assumption of Lemma 10 is satisfied. If $1 < \theta \leq 2$, it follows that our assumption $\theta < d/2$ (in Theorem 1) is equivalent to $d/\theta > d/(d - \theta)$ such that

$q > d/(d - \theta)$, and Lemma 10 is applicable also in this case. Consider $1 < p < 2$ and $\min\{p, d/\theta\} < q$. We employ Hölder's inequality for $r = q/(q - p)$ to find that

$$(24) \quad \begin{aligned} I_2 &= \int_{\mathbb{R}^d \setminus B_R(0)} |x|^{dp} \rho(x, t)^p G_\theta(x, t + 1/\theta)^{1-p} |x|^{-dp} dx \\ &\leq \left\| |x|^d \rho(x, t) \right\|_{L^q(\mathbb{R}^d)}^p \left(\int_{\mathbb{R}^d \setminus B_R(0)} G_\theta(x, t + 1/\theta)^{(1-p)r} |x|^{-dpr} dx \right)^{1/r}. \end{aligned}$$

Using the pointwise estimates (9), the second factor can be estimated as

$$\begin{aligned} \int_{\mathbb{R}^d \setminus B_R(0)} G_\theta(x, t + 1/\theta)^{(1-p)r} |x|^{-dpr} dx &\leq \int_{\mathbb{R}^d \setminus B_R(0)} (C(t + 1/\theta) |x|^{-d-\theta})^{(1-p)r} |x|^{-dpr} dx \\ &\leq C(t + 1/\theta)^{(1-p)r} \int_{\mathbb{R}^d \setminus B_R(0)} |x|^{(d+\theta)(p-1)r - dpr} dx \\ &\leq C(t + 1/\theta)^{(1-p)r} \int_R^\infty \eta^{(d+\theta)(p-1)r - dpr + d - 1} d\eta. \end{aligned}$$

The last integral exists if and only if $(d + \theta)(p - 1)r - dpr + d - 1 < -1$ or equivalently $q < dp/((p - 1)\theta)$. Since $dp/((p - 1)\theta) \rightarrow \infty$ as $p \rightarrow 1$, for any large $q > d/\theta$, we can choose a sufficiently small $p > 1$ such that $d/\theta < q < dp/((p - 1)\theta)$. Collecting the estimates starting from (24) with $R = K(t + 1/\theta)^{1/\theta}$ and using Lemma 10, we deduce that

$$\begin{aligned} I_2 &\leq C(1 + t)^{dp/(\theta q)} (t + 1/\theta)^{1-p} |R|^{(d+\theta)(p-1) - dp + d/r} \\ &\leq C(1 + t)^{dp/(\theta q)} (t + 1/\theta)^{1-p} (t + 1/\theta)^{((d+\theta)(p-1) - dp + d/r)/\theta} \leq C \end{aligned}$$

is uniformly bounded in time. Thus, the integral $I(t)$ in (23) and consequently $E_p[v(t)]$ are well-defined for all $t \geq 0$. This finishes the proof of Theorem 1.

4. APPLICATION TO OTHER EQUATIONS

The entropy method can be applied to other semilinear equations, and in this section, we give two illustrative examples.

4.1. Quasi-geostrophic equations. The two-dimensional quasi-geostrophic equations (for sufficiently smooth solutions) read as

$$(25) \quad \partial_t \rho + (-\Delta)^{\theta/2} \rho = \operatorname{div}(\rho \nabla^\perp \psi), \quad (-\Delta)^{1/2} \psi = -\rho \quad \text{in } \mathbb{R}^2, \quad t > 0,$$

with the initial condition $\rho(0) = \rho_0 \geq 0$ in \mathbb{R}^2 , $0 < \theta \leq 2$, and $\nabla^\perp = (-\partial_2, \partial_1)^T$. This model approximates the atmospheric and oceanic fluid flow in a certain physical regime. The variables ρ and ψ refer to the temperature and the stream function, respectively. For the geophysical background, we refer to [11]. The existence of global solutions in a Sobolev space setting was shown in [20], for instance. The time decay in the $L^\infty(\mathbb{R}^2)$ norm was investigated in [13]. Estimate (4) can be obtained from standard estimates since the nonlinear part vanishes after multiplication by ρ^{p-1} ($p > 1$) and integration over \mathbb{R}^2 :

$$\int_{\mathbb{R}^2} \rho^{p-1} \operatorname{div}(\rho \nabla^\perp \psi) dx = (p - 1) \int_{\mathbb{R}^2} \rho^{p-1} \nabla \rho \cdot \nabla^\perp \psi dx = 0.$$

For the analysis of the asymptotic profile, we use the transformation (15) and $\phi(x, t) := (e^t/M)\psi(e^t x, (e^{\theta t} - 1)/\theta)$, yielding

$$\begin{aligned} \partial_t u + (-\Delta)^{\theta/2} u &= \operatorname{div}(u \nabla V) + M e^{-(3-\theta)t} \operatorname{div}(u \nabla^\perp \phi), \\ (-\Delta)^{1/2} \phi &= -u \quad \text{in } \mathbb{R}^2, \quad t > 0, \quad u(0) = \rho_0/M, \end{aligned}$$

where $V(x) = \frac{1}{2}|x|^2$. Defining the entropy as in (21), a computation similar to the one in the proof of Lemma 9 gives for $v = u/G_\theta(1/\theta)$:

$$\frac{dE_p}{dt}[v] \leq -\theta E_p[v] + p M e^{-(3-\theta)t} \int_{\mathbb{R}^2} v^{p-1} G_\theta(1/\theta)^{1-1/p} u \nabla(G_\theta(1/\theta)^{1-1/p}) \cdot \nabla^\perp \phi \, dx.$$

Estimate (10) shows that $\nabla(G_\theta(1/\theta)^{1-1/p}) \in L^\infty(\mathbb{R}^2)$. Moreover,

$$\sup_{t>0} \|\nabla^\perp \phi\|_{L^p(\mathbb{R}^2)} = \sup_{t>0} \|\nabla^\perp (-\Delta)^{-1/2} u\|_{L^p(\mathbb{R}^2)} \leq C \sup_{t>0} \|u\|_{L^p(\mathbb{R}^d)}$$

is finite. The regularity of ρ implies that $u \in L^\infty(0, \infty; L^\infty(\mathbb{R}^2))$. Therefore, using Hölder's and Young's inequalities, we find that

$$\begin{aligned} & \left| \int_{\mathbb{R}^2} v^{p-1} G_\theta(1/\theta)^{1-1/p} u \nabla(G_\theta(1/\theta)^{1-1/p}) \cdot \nabla^\perp \phi \, dx \right| \\ & \leq \|\nabla(G_\theta(1/\theta)^{1-1/p})\|_{L^\infty(\mathbb{R}^2)} \|u\|_{L^\infty(\mathbb{R}^2)} \|\nabla^\perp \phi\|_{L^p(\mathbb{R}^2)} \left(\int_{\mathbb{R}^2} v^p G_\theta(1/\theta) \, dx \right)^{1-1/p} \\ & \leq C(E_p[v] + 1). \end{aligned}$$

Thus, the same procedure as in Section 3 shows that the solution to (25) satisfies (5).

4.2. Generalized Burgers equation. The generalized Burgers equation in one space dimension

$$(26) \quad \partial_t \rho + (-\partial_x^2)^{\theta/2} \rho + \rho \partial_x \rho = 0 \quad \text{in } \mathbb{R}, \quad t > 0, \quad \rho(0) = \rho_0,$$

with $\theta = 2$ is a basic model for one-dimensional fluid flows with interacting nonlinear and dissipative phenomena, and it describes for $0 < \theta < 2$ the overdriven detonation in gases [10] or the anomalous diffusion in semiconductor growth [38]. The value $\theta = 1$ is a threshold for the occurrence of singularities [14], and the existence of global solutions holds for $1 < \theta < 2$. Global solutions in Besov spaces were also shown to exist [19], and in the same paper, the self-similar asymptotics have been analyzed. The function u , defined by (15), solves

$$\partial_t u + (-\partial_x^2)^{\theta/2} u = \partial_x(u \partial_x V) - M e^{-(2-\theta)t} u \partial_x u,$$

and the entropy method yields, similar as in the previous subsection,

$$\begin{aligned} \frac{dE_p}{dt}[v] &\leq -\theta E_p[v] + p M e^{-(2-\theta)t} \int_{\mathbb{R}} \partial_x u v^p G_\theta(1/\theta) \, dx \\ &\leq -\theta E_p[v] + p M e^{-(2-\theta)t} \|\partial_x u\|_{L^\infty(\mathbb{R})} (E_p[v] + 1). \end{aligned}$$

Assuming that the solution to (26) satisfies the bound $\|\partial_x \rho(t)\|_{L^\infty(\mathbb{R})} \leq Ct^{-2/\theta}$, it follows that $\partial_x u \in L^\infty(0, \infty; L^\infty(\mathbb{R}))$, and we obtain the decay estimate (5), where a restriction for the range of θ is expected from estimating the nonlinear terms.

Observe that the entropy method does not work in the case $\theta = 2$ (see the discussion at the end of the introduction in [19]). Moreover, when $0 < \theta < 1$, there are initial data such that the solution develops gradient blowup in finite time [14].

APPENDIX A. PROOF OF LEMMA 10

For later reference, we note that, by the Poisson representation formula and property (4) with $p = 1$ and $p = \infty$,

$$\begin{aligned}
|\nabla \psi(x, t)| &= |\nabla(-\Delta)^{-1} \rho(x, t)| = c_d \left| \int_{\mathbb{R}^d} \frac{x-y}{|x-y|^d} \rho(y, t) dy \right| \\
&\leq c_d \int_{\{|x-y| \leq (1+t)^{1/\theta}\}} \frac{|\rho(y, t)|}{|x-y|^{d-1}} dy + c_d \int_{\{|x-y| > (1+t)^{1/\theta}\}} \frac{|\rho(y, t)|}{|x-y|^{d-1}} dy \\
&\leq C(1+t)^{1/\theta} \|\rho(t)\|_{L^\infty(\mathbb{R}^d)} + C(1+t)^{-(d-1)/\theta} \|\rho(t)\|_{L^1(\mathbb{R}^d)} \\
(27) \quad &\leq C(1+t)^{-d/\theta+1/\theta}.
\end{aligned}$$

Step 1: $1 < \theta \leq 2$. The solution to (1) can be formulated as the mild solution

$$\rho(t) = G_\theta(t) * \rho_0 + \int_0^t \nabla G_\theta(t-s) * (\rho \nabla(-\Delta)^{-1} \rho)(s) ds.$$

We multiply this equation by $|x|^d$,

$$\begin{aligned}
\||x|^d \rho(t)\| &\leq \||x|^d G_\theta(t) * \rho_0\| \\
&\quad + C \int_0^t \int_{\mathbb{R}^d} (|x-y|^d + |y|^d) |\nabla G_\theta(t-s, x-y)| |(\rho \nabla(-\Delta)^{-1} \rho)(s, y)| dy ds,
\end{aligned}$$

and take the $L^q(\mathbb{R}^d)$ norm:

$$\begin{aligned}
\||x|^d \rho(t)\|_{L^q(\mathbb{R}^d)} &\leq \||x|^d G_\theta(t)\|_{L^q(\mathbb{R}^d)} \|\rho_0\|_{L^1(\mathbb{R}^d)} + \|G_\theta(t)\|_{L^1(\mathbb{R}^d)} \||x|^d \rho_0\|_{L^q(\mathbb{R}^d)} \\
&\quad + C \int_0^t \||x|^d \nabla G_\theta(t-s)\|_{L^q(\mathbb{R}^d)} \|\rho(s)\|_{L^1(\mathbb{R}^d)} \|(\nabla(-\Delta)^{-1} \rho)(s)\|_{L^\infty(\mathbb{R}^d)} ds \\
&\quad + C \int_0^t \|\nabla G_\theta(t-s)\|_{L^1(\mathbb{R}^d)} \||x|^d \rho(s)\|_{L^q(\mathbb{R}^d)} \|(\nabla(-\Delta)^{-1} \rho)(s)\|_{L^\infty(\mathbb{R}^d)} ds \\
(28) \quad &=: I_1 + I_2 + I_3.
\end{aligned}$$

We estimate term by term.

Estimates (9) show that $\||x|^d G_\theta(t)\|_{L^\infty(\mathbb{R}^d)} \leq C$ and $\||x|^d G_\theta(t)\|_{L^q(\mathbb{R}^d)} \leq C(1+t)^{d/(q\theta)}$ for $q > d/\theta$, and $\|G_\theta(t)\|_{L^1(\mathbb{R}^d)} = 1$. We deduce that

$$I_1 \leq C(1+t)^{d/(q\theta)}.$$

For the estimate of I_2 , we first infer from the pointwise estimates (10) that

$$\left\| |x|^d \nabla G_\theta(t) \right\|_{L^q(\mathbb{R}^d)} \leq C \left\| \min\{t^{-1/\theta}, |x|^{-1}\} \right\|_{L^\infty(\mathbb{R}^d)} \left\| |x|^d G_\theta(t) \right\|_{L^q(\mathbb{R}^d)} \leq C t^{-\frac{1}{\theta} + \frac{d}{q\theta}}.$$

Therefore, taking into account (27) and the inequality $\theta \leq d$,

$$I_2 \leq C \int_0^t (t-s)^{-1/\theta + d/(q\theta)} (1+s)^{-d/\theta + 1/\theta} ds \leq C(1+t)^{d/(q\theta) - d/\theta + 1} \leq C(1+t)^{d/(q\theta)}.$$

Note that $\theta > 1$ implies that $-1/\theta + d/(q\theta) > -1$ and so, $(t-s)^{-1/\theta + d/(q\theta)}$ is integrable in $(0, t)$.

For the integral I_3 , we first observe that, again by (9) and (10), $\|\nabla G_\theta(t-s)\|_{L^1(\mathbb{R}^d)} \leq C(t-s)^{-1/\theta}$. Hence, with (27),

$$\begin{aligned} I_3 &\leq C \int_0^t (t-s)^{-1/\theta} \left\| |x|^d \rho(s) \right\|_{L^q(\mathbb{R}^d)} (1+s)^{-d/\theta + 1/\theta} ds \\ &\leq C \sup_{0 < s < t} \left((1+s)^{\frac{d}{\theta}(1-\frac{1}{q}) - \frac{d}{\theta}} \left\| |x|^d \rho(s) \right\|_{L^q(\mathbb{R}^d)} \right) \int_0^t (t-s)^{-1/\theta} (1+s)^{-\frac{d}{\theta}(1-\frac{1}{q}) + \frac{1}{\theta}} ds \\ &\leq C \sup_{0 < s < t} \left((1+s)^{\frac{d}{\theta}(1-\frac{1}{q}) - \frac{d}{\theta}} \left\| |x|^d \rho(s) \right\|_{L^q(\mathbb{R}^d)} \right) \tau(t), \end{aligned}$$

where $\tau(t) = (1+t)^{-\frac{d}{\theta}(1-\frac{1}{q})+1}$. Note that again, $\theta > 1$ is needed to ensure that $(t-s)^{-1/\theta}$ is integrable in $(0, t)$. Since $q > d/(d-\theta)$, it follows that $\tau(t) \rightarrow 0$ as $t \rightarrow \infty$.

We conclude from (28) that

$$\left\| |x|^d \rho(t) \right\|_{L^q(\mathbb{R}^d)} \leq C(1+t)^{d/(q\theta)} + C\tau(t) \sup_{0 < s < t} \left((1+s)^{-d/(q\theta)} \left\| |x|^d \rho(s) \right\|_{L^q(\mathbb{R}^d)} \right).$$

This can be written as

$$\begin{aligned} &\sup_{0 < s < t} \left((1+s)^{-d/(q\theta)} \left\| |x|^d \rho(s) \right\|_{L^q(\mathbb{R}^d)} \right) \\ &\leq C + (1+t)^{-d/(q\theta)} \tau(t) \sup_{0 < s < t} \left((1+s)^{-d/(q\theta)} \left\| |x|^d \rho(s) \right\|_{L^q(\mathbb{R}^d)} \right). \end{aligned}$$

Choosing $t > 0$ sufficiently large, we infer that

$$\sup_{0 < s < t} \left((1+s)^{-d/(q\theta)} \left\| |x|^d \rho(s) \right\|_{L^q(\mathbb{R}^d)} \right) \leq C,$$

proving the claim.

Step 2: $0 < \theta \leq 1$. We first study the even dimensional case, $d = 2m$ for some $m \in \mathbb{N}$. The function $P(x, t) = |x|^d \rho(x, t)$ solves

$$(29) \quad \partial_t P + (-\Delta)^{\theta/2} P - \operatorname{div}(P \nabla \psi) = [(-\Delta)^{\theta/2}, |x|^d] \rho + [|x|^d, \operatorname{div}](\rho \nabla \psi),$$

where $[A, B] = AB - BA$ is the commutator. The first term on the right-hand side becomes

$$[(-\Delta)^{\theta/2}, |x|^d] \rho = \mathcal{F}^{-1} [[|\xi|^\theta, (-\Delta)^m] \hat{\rho}].$$

The highest order derivative of $\hat{\rho}$ in the commutator cancels, since

$$[|\xi|^\theta, (-\Delta)^m] \hat{\rho} = |\xi|^\theta (-\Delta)^m \hat{\rho} - (-\Delta)^m (|\xi|^\theta \hat{\rho}),$$

and the chain rule gives the sum of derivatives up to order $2m - 1 = d - 1$. Hence, we infer from the Fourier convolution formula that

$$\begin{aligned} [(-\Delta)^{\theta/2}, |x|^d]\rho &= \sum_{|\alpha|+|\beta|=d, |\beta|\leq d-1} a_{\alpha,\beta} \mathcal{F}^{-1}[(i\nabla)^\alpha(|\xi|^\theta)(i\nabla)^\beta \widehat{\rho}] \\ &= \sum_{|\alpha|+|\beta|=d, |\beta|\leq d-1} a_{\alpha,\beta} \mathcal{F}^{-1}[(i\nabla)^\alpha(|\xi|^\theta)] * (x^\beta \rho), \end{aligned}$$

where $\alpha, \beta \in \mathbb{N}_0^d$ are multi-indices, $a_{\alpha,\beta} \in \mathbb{R}$ are some constants, and $(i\nabla)^\beta$ and x^β are to be understood as products of the components of the corresponding vectors. Evaluating the commutator, the second term on the right-hand side of (29) equals

$$[|x|^d, \operatorname{div}](\rho \nabla \psi) = |x|^d \operatorname{div}(\rho \nabla \psi) - \operatorname{div}(|x|^d \rho \nabla \psi) = -d|x|^{d-2} \rho x \cdot \nabla \psi.$$

Hence, multiplying (29) by P^{q-1} with $q > d/\theta > 1$, we have

$$\begin{aligned} (30) \quad & \frac{1}{q} \frac{d}{dt} \int_{\mathbb{R}^d} P^q dx + \int_{\mathbb{R}^d} P^{q-1} (-\Delta)^{\theta/2} P dx + \frac{q-1}{q} \int_{\mathbb{R}^d} \nabla \psi \cdot \nabla P^q dx \\ &= \sum_{|\alpha|+|\beta|=d, |\beta|\leq d-1} a_{\alpha,\beta} \int_{\mathbb{R}^d} P^{q-1} \mathcal{F}^{-1}[(i\nabla)^\alpha(|\xi|^\theta)] * (x^\beta \rho) dx \\ & \quad - d \int_{\mathbb{R}^d} P^{q-1} |x|^{d-2} \rho x \cdot \nabla \psi dx. \end{aligned}$$

The Stroock-Varopoulos inequality (see, e.g., [31, Prop. 2.1], [12, Lemma 1] or [27, Theorem 2.1]) gives for the second term on the left-hand side:

$$\int_{\mathbb{R}^d} P^{q-1} (-\Delta)^{\theta/2} P dx \geq \frac{2}{q} \int_{\mathbb{R}^d} |(-\Delta)^{\theta/4} (P^{q/2})|^2 dx.$$

Furthermore, integrating by parts and using the Poisson equation, the third term on the left-hand side of (30) becomes

$$\frac{q-1}{q} \int_{\mathbb{R}^d} \nabla \psi \cdot \nabla P^q dx = \frac{q-1}{q} \int_{\mathbb{R}^d} P^q \rho dx \geq 0.$$

Thus, multiplying (30) by $q(1+t)^{-\gamma q}$ for some $0 < \gamma < d/(\theta q)$ and integrating over time,

$$\begin{aligned} & (1+t)^{-\gamma q} \|P(t)\|_{L^q(\mathbb{R}^d)}^q + \gamma q \int_0^t (1+s)^{-\gamma q-1} \|P(s)\|_{L^q(\mathbb{R}^d)}^q ds \\ & \quad + 2 \int_0^t (1+s)^{-\gamma q} \|(-\Delta)^{\theta/4} (P(s)^{q/2})\|_{L^2(\mathbb{R}^d)}^2 ds \\ & \quad + (q-1) \int_0^t (1+s)^{-\gamma q} \int_{\mathbb{R}^d} P(s)^q \rho(s) dx ds \\ & \leq q \sum_{|\alpha|+|\beta|=d, |\beta|\leq d-1} a_{\alpha,\beta} \int_0^t (1+s)^{-\gamma q} \int_{\mathbb{R}^d} P(s)^{q-1} \mathcal{F}^{-1}[(i\nabla)^\alpha(|\xi|^\theta)] * (x^\beta \rho(s)) dx ds \end{aligned}$$

$$(31) \quad -d \int_0^t (1+s)^{-\gamma q} \int_{\mathbb{R}^d} P(s)^{q-1} |x|^{d-2} \rho(s) x \cdot \nabla \psi(s) dx ds + \| |x|^d \rho_0 \|_{L^q(\mathbb{R}^d)}^q.$$

The third and fourth terms on the left-hand side will be neglected, and we need to estimate only the terms on the right-hand side.

Let us consider the first term on the right-hand side. The Hölder inequality with $\eta = q/(q-1) > 1$ and $\eta' = q$ implies that

$$\left| \int_{\mathbb{R}^d} P^{q-1} \mathcal{F}^{-1}[(i\nabla)^\alpha(|\xi|^\theta)] * (x^\beta \rho) dx ds \right| \leq \|P\|_{L^q(\mathbb{R}^d)}^{q-1} \|\mathcal{F}^{-1}[(i\nabla)^\alpha(|\xi|^\theta)] * (x^\beta \rho)\|_{L^q(\mathbb{R}^d)}.$$

Since $(i\nabla)^\alpha |\xi|^\theta = |\xi|^{\theta-2|\alpha|} P_{|\alpha|}(\xi)$ for some homogeneous polynomial $P_{|\alpha|}$ of ξ with order $|\alpha|$, we see that $\mathcal{F}^{-1}[(i\nabla)^\alpha |\xi|^\theta] * (x^\beta \rho) = |\nabla|^{\theta-2|\alpha|} P_{|\alpha|}(i\nabla)(x^\beta \rho)$. The operator $|\nabla|^{-|\alpha|} P_{|\alpha|}(i\nabla)$ is bounded in $L^q(\mathbb{R}^d)$ since it is a polynomial of Riesz transforms. Hence, we can apply the Hardy-Littlewood-Sobolev inequality [34, Section V.1.1, Theorem 1] with $1/r = 1/q + (|\alpha| - \theta)/d$ to obtain

$$\| |\nabla|^{\theta-2|\alpha|} P_{|\alpha|}(i\nabla)(x^\beta \rho) \|_{L^q(\mathbb{R}^d)} \leq C \| |\nabla|^{\theta-|\alpha|} (x^\beta \rho) \|_{L^q(\mathbb{R}^d)} \leq C \| x^\beta \rho \|_{L^r(\mathbb{R}^d)}.$$

We infer that

$$\left| \int_{\mathbb{R}^d} P^{q-1} \mathcal{F}^{-1}[(i\nabla)^\alpha(|\xi|^\theta)] * (x^\beta \rho) dx ds \right| \leq C \|P\|_{L^q(\mathbb{R}^d)}^{q-1} \|x^\beta \rho\|_{L^r(\mathbb{R}^d)}.$$

We proceed by applying the Hölder inequality with $\eta = dq/(|\beta|r) > 1$ and $\eta' = dq/(dq - |\beta|r)$ to the last norm:

$$\| |x|^{|\beta|} \rho \|_{L^r(\mathbb{R}^d)}^r = \int_{\mathbb{R}^d} |x|^{|\beta|r} \rho^{|\beta|r/d} \cdot \rho^{r(1-|\beta|/d)} dx \leq \| |x|^d \rho \|_{L^q(\mathbb{R}^d)}^{|\beta|r/d} \| \rho \|_{L^\nu(\mathbb{R}^d)}^{r(1-|\beta|/d)},$$

where $\nu = (d-|\beta|)qr/(dq-|\beta|r)$. Note that $\nu > 1$. Indeed, inserting $1/r = 1/q + (|\alpha| - \theta)/d$ and $|\alpha| + |\beta| = d$ yields

$$\nu = \frac{d-|\beta|}{d/r - |\beta|/q} = \frac{d-|\beta|}{d/q + |\alpha| - \theta - |\beta|/q} = \frac{|\alpha|}{|\alpha|/q + |\alpha| - \theta},$$

and this is larger than one if and only if $|\alpha|/q < \theta$. This is true since $|\alpha|/q \leq d/q < \theta$ by the choice of q . Then, splitting the integrand,

$$\begin{aligned} & \left| \int_0^t (1+s)^{-\gamma q} \int_{\mathbb{R}^d} P(s)^{q-1} \mathcal{F}^{-1}[(i\nabla)^\alpha(|\mu|^\theta)] * (x^\beta \rho(s)) dx ds \right| \\ & \leq C \int_0^t (1+s)^{-\gamma q} \|P\|_{L^q(\mathbb{R}^d)}^{|\beta|/d+q-1} \|\rho\|_{L^\nu(\mathbb{R}^d)}^{1-|\beta|/d} ds \\ & \leq C \int_0^t ((1+s)^{-\gamma q + \mu - 1})^{1/\mu} \|\rho\|_{L^\nu(\mathbb{R}^d)}^{1-|\beta|/d} ((1+s)^{-\gamma q - 1})^{1-1/\mu} \|P\|_{L^q(\mathbb{R}^d)}^{|\beta|/d+q-1} ds \\ & = C \int_0^t \left((1+s)^{-\gamma q + \mu - 1} \|\rho\|_{L^\nu(\mathbb{R}^d)}^q \right)^{1/\mu} \left((1+s)^{-\gamma q - 1} \|P\|_{L^q(\mathbb{R}^d)}^q \right)^{(\mu-1)/\mu} ds. \end{aligned}$$

Young's inequality with $\eta = \mu > 1$ and $\eta' = \mu/(\mu - 1)$ and any $\varepsilon > 0$ yields

$$(32) \quad \left| \int_0^t (1+s)^{-\gamma q} \int_{\mathbb{R}^d} P(s)^{q-1} \mathcal{F}^{-1}[(i\nabla)^\alpha(|\xi|^\theta)] * (x^\beta \rho(s)) \, dx \, ds \right| \\ \leq C_\varepsilon \int_0^t (1+s)^{-\gamma q + \mu - 1} \|\rho\|_{L^\nu(\mathbb{R}^d)}^q \, ds + \varepsilon \int_0^t (1+s)^{-\gamma q - 1} \|P\|_{L^q(\mathbb{R}^d)}^q \, ds.$$

For sufficiently small $\varepsilon > 0$, the last term is absorbed by the second term on the left-hand side of (31). In view of (4), the first term on the right-hand side of (32) is estimated according to

$$\int_0^t (1+s)^{-\gamma q + \mu - 1} \|\rho\|_{L^\nu(\mathbb{R}^d)}^q \, ds \leq C(1+t)^{-\gamma q + \mu - q \frac{d}{\theta} (1 - \frac{1}{\nu})}.$$

Using the definitions of μ , ν , and $1/r = 1/q + (|\alpha| - \theta)/d$ as well the property $|\alpha| + |\beta| = d$, it follows that

$$\begin{aligned} \mu - q \frac{d}{\theta} \left(1 - \frac{1}{\nu}\right) &= \frac{dq}{d - |\beta|} - \frac{dq}{\theta} \left(1 - \frac{dq - |\beta|r}{(d - |\beta|)qr}\right) \\ &= \frac{dq}{d - |\beta|} \left(1 - \frac{d - |\beta|}{\theta} + \frac{d}{\theta r} - \frac{|\beta|}{q\theta}\right) \\ &= \frac{dq}{d - |\beta|} \left(1 - \frac{d}{\theta} + \frac{|\beta|}{\theta} + \frac{d}{q\theta} + \frac{|\alpha| - \theta}{\theta} - \frac{|\beta|}{q\theta}\right) \\ &= \frac{dq}{d - |\beta|} \left(\frac{d}{q\theta} - \frac{|\beta|}{q\theta}\right) = \frac{d}{\theta}. \end{aligned}$$

Therefore,

$$\int_0^t (1+s)^{-\gamma q + \mu - 1} \|\rho\|_{L^\nu(\mathbb{R}^d)}^q \, ds \leq C(1+t)^{-\gamma q + d/\theta}.$$

It remains to bound the second term on the right-hand side of (31). We apply Hölder's inequality with $\eta = q/(q - 1) > 1$ and $\eta' = q$ to obtain

$$(33) \quad \left| \int_{\mathbb{R}^d} P^{q-1} |x|^{d-1} \rho |\nabla \psi| \, dx \right| \leq \|P\|_{L^q(\mathbb{R}^d)}^{q-1} \| |x|^{d-1} \rho \|_{L^q(\mathbb{R}^d)} \|\nabla \psi\|_{L^\infty(\mathbb{R}^d)}.$$

The second factor is estimated using Hölder's inequality again with $\eta = d/(d - 1) > 1$, $\eta' = 1/(1 - 1/\eta) = d$ and using (4):

$$\begin{aligned} \| |x|^{d-1} \rho \|_{L^q(\mathbb{R}^d)}^q &= \int_{\mathbb{R}^d} |x|^{(d-1)q} \rho^{q/\eta} \cdot \rho^{q(1-1/\eta)} \, dx \leq \left(\int_{\mathbb{R}^d} |x|^{dq} \rho^q \, dx \right)^{1/\eta} \left(\int_{\mathbb{R}^d} \rho^q \, dx \right)^{1-1/\eta} \\ &= \| |x|^d \rho \|_{L^q(\mathbb{R}^d)}^{q(1-\frac{1}{d})} \|\rho\|_{L^q(\mathbb{R}^d)}^{q/d} \leq C \|P\|_{L^q(\mathbb{R}^d)}^{q(1-\frac{1}{d})} (1+s)^{-q/\theta + 1/\theta}. \end{aligned}$$

We conclude from (27) and (33) that

$$\left| \int_{\mathbb{R}^d} P^{q-1} |x|^{d-1} \rho |\nabla \psi| \, dx \right| \leq C(1+s)^{-d/\theta + 1/(q\theta)} \|P\|_{L^q(\mathbb{R}^d)}^{q-1/d}.$$

Thus, using the Young inequality with $\eta = dq$, $\eta' = dq/(dq - 1)$ (we only need that $\eta > 0$), for any $\varepsilon > 0$, the second term on the right-hand side of (31) becomes

$$\begin{aligned}
& \left| \int_0^t (1+s)^{-\gamma q} \int_{\mathbb{R}^d} P(s)^{q-1} |x|^{d-2} \rho x \cdot \nabla \psi(s) \, dx \, ds \right| \\
& \leq C \int_0^t (1+s)^{-\gamma q - d/\theta + 1/(q\theta)} \|P\|_{L^q(\mathbb{R}^d)}^{q-1/d} \, ds \\
& = C \int_0^t (1+s)^{1+1/(q\theta) - 1/(dq) - d/\theta - \gamma/d} \left((1+s)^{-\gamma q - 1} \|P\|_{L^q(\mathbb{R}^d)}^q \right)^{1-1/(dq)} \, ds \\
& \leq C_\varepsilon \int_0^t (1+s)^{dq+d/\theta - 1 - d^2q/\theta - \gamma q} \, ds + \varepsilon \int_0^t (1+s)^{-\gamma q - 1} \|P\|_{L^q(\mathbb{R}^d)}^q \, ds. \\
& \leq C(1+t)^{-\gamma q + d/\theta} + \varepsilon \int_0^t (1+s)^{-\gamma q - 1} \|P\|_{L^q(\mathbb{R}^d)}^q \, ds,
\end{aligned}$$

where we used $dq - d^2q/\theta \leq 0$ which is equivalent to $\theta \leq d$, and this property has been assumed. The last expression can be absorbed by the second term on the left-hand side of (31) for sufficiently small $\varepsilon > 0$.

Summarizing the estimates, we conclude from (31) (for sufficiently small $\varepsilon > 0$) that

$$(1+t)^{-\gamma q} \|P(t)\|_{L^q(\mathbb{R}^d)}^q \leq C(1+t)^{-\gamma q + d/\theta},$$

which gives the desired result.

When the dimension is odd, $d = 2m + 1$ for some $n \in \mathbb{N}_0$, we choose $P(x, t) = |x|^{d-1} x \rho(x, t)$ and proceed as above. This concludes the proof.

APPENDIX B. FORMAL DERIVATION OF THE FRACTIONAL DRIFT-DIFFUSION-POISSON SYSTEM

Fractional diffusion may be derived from linear kinetic transport models by using the Fourier-Laplace transform [29] or by exploiting the harmonic extension definition of the fractional diffusion operator [4]. An alternative is Mellet's moment method [28], which was used to derive the first equation in (1) with fixed force field $E = \nabla \psi$ [1]. In this section, we sketch a formal derivation of the drift-diffusion-Poisson system following [1] to explain the origin of equations (1).

The starting point is the scaled Boltzmann equation

$$(34) \quad \varepsilon^{\theta-1} \partial_t f_\varepsilon + v \cdot \nabla_x f_\varepsilon + \varepsilon^{\theta-2} \nabla_x \psi_\varepsilon \cdot \nabla_v f_\varepsilon = \varepsilon^{-1} Q(f_\varepsilon),$$

for the distribution function $f_\varepsilon(x, v, t)$, where $x \in \mathbb{R}^d$ is the spatial variable, $v \in \mathbb{R}^d$ is the velocity, and $t > 0$ is the time. We prescribe the initial condition $f(x, v, 0) = f_I(x, v)$ for $(x, v) \in \mathbb{R}^d \times \mathbb{R}^d$. The electric potential $\psi_\varepsilon(x, t)$ is governed by the Poisson equation

$$(35) \quad -\Delta_x \psi_\varepsilon = \rho_\varepsilon, \quad \rho_\varepsilon := \int_{\mathbb{R}^d} f_\varepsilon \, dv,$$

where ρ_ε is the particle density, and Q is the linear Boltzmann operator

$$Q(f) = \int_{\mathbb{R}^d} (\sigma(v, v')M(v)f(v') - \sigma(v', v)M(v')f(v)) dv',$$

where $\sigma(v, v') > 0$ is the scattering rate. The parameter $\varepsilon > 0$ measures the collision frequency, and $\theta > 0$ fixes the relation between diffusion, due to scattering, and advection, due to the acceleration field.

System (34)-(35) models the evolution of particles in a dilute gas subject to a self-consistent acceleration field $E_\varepsilon = \nabla_x \psi_\varepsilon$. For instance, ρ_ε may describe the density of electrons in a semiconductor, and ψ_ε is the electric potential. The function $M(v)$ models the thermodynamic equilibrium. In semiconductor theory, it is typically given by the Maxwellian, and in this case $\theta = 2$ [21]. Here, we are interested in the case $1 < \theta < 2$. The main assumptions on the equilibrium M are that it is positive, normalized, even, and heavy-tailed, i.e.

$$M(v) \sim \frac{\gamma}{|x|^{d+\theta}} \quad \text{as } |x| \rightarrow \infty$$

for some $\gamma > 0$ and $1 < \theta < 2$.

Equations (1) are derived in the limit $\varepsilon \rightarrow 0$. To this end, we expand $f_\varepsilon = F_\varepsilon + \varepsilon^{\theta/2} r_\varepsilon$, where r_ε is a remainder term. In contrast to standard kinetic theory, F_ε is not given by the equilibrium distribution $M(v)$ (up to a factor) but it is the unique solution to

$$(36) \quad \varepsilon^{\theta-1} E_\varepsilon \cdot \nabla_v F_\varepsilon = Q(F_\varepsilon), \quad \int_{\mathbb{R}^d} F_\varepsilon dv = 1.$$

We can interpret this equation in two ways. First, because of $\theta > 1$, (36) converges formally to $Q(F_0) = 0$ as $\varepsilon \rightarrow 0$ with $F_0 = \lim_{\varepsilon \rightarrow 0} F_\varepsilon$, such that (36) is an approximation of the equilibrium condition $Q(F_0) = 0$. Second, we can write $F_\varepsilon(x, v, t) = F(v, \varepsilon^{\theta-1} E_\varepsilon(x, t))$, where F is the unique solution to $E_\varepsilon \cdot \nabla_v F = Q(F)$, $\int_{\mathbb{R}^d} F dv = 1$, and this transformation eliminates the factor in ε . This equation appears in the high-field limit, first studied in kinetic theory for semiconductors by Poupaud [32].

Inserting the expansion $f_\varepsilon = F_\varepsilon + \varepsilon^{\theta/2} r_\varepsilon$ into the Boltzmann equation (34), dividing the resulting equation by $\varepsilon^{\theta/2-1}$, and observing that F_ε solves (36), we find that

$$(37) \quad \varepsilon^{\theta/2} \partial_t f_\varepsilon + \varepsilon^{1-\theta/2} v \cdot \nabla_x (F_\varepsilon + \varepsilon^{\theta/2} r_\varepsilon) = -\varepsilon^{\theta-1} E_\varepsilon \cdot \nabla_x r_\varepsilon + Q(r_\varepsilon).$$

Formally, the left-hand side converges to zero as $\varepsilon \rightarrow 0$ (since $\theta < 2$). Therefore $R_\varepsilon := -\varepsilon^{\theta-1} E_\varepsilon \cdot \nabla_x r_\varepsilon + Q(r_\varepsilon) \rightarrow 0$. In fact, it can be even proven that $\varepsilon^{-\theta/2} R_\varepsilon \rightarrow 0$; this is contained in [1, Prop. 4.1]. We assume that $f_\varepsilon \rightarrow f$ as $\varepsilon \rightarrow 0$. Hence $\rho_\varepsilon \rightarrow \rho = \int_{\mathbb{R}^d} f dv$. In view of the Poisson equation, this implies that $E_\varepsilon \rightarrow E$, where $E = \nabla_x \psi$ and $-\Delta_x \psi = \rho$.

Next, following [1], we multiply (37) by some test function χ and integrate over $S = \mathbb{R}^d \times \mathbb{R}^d \times (0, T)$:

$$(38) \quad \iiint_S \partial_t f_\varepsilon \chi dx dv dt + \varepsilon^{1-\theta} \iiint_S v \cdot \nabla_x (F_\varepsilon + \varepsilon^{\theta/2} r_\varepsilon) \chi dx dv dt = \varepsilon^{-\theta/2} \iiint_S R_\varepsilon \chi dx dv dt.$$

The key idea is to choose $\chi = \chi_\varepsilon$ as the unique solution to

$$\varepsilon v \cdot \nabla_x \chi_\varepsilon = \nu(v)(\chi_\varepsilon - \phi) \quad \text{in } \mathbb{R}^d,$$

where $\phi(x, t)$ is some test function and $\nu(v) = \int_{\mathbb{R}^d} \sigma(v', v) M(v') dv'$. Formally, $\chi_\varepsilon \rightarrow \delta_v \phi$ as $\varepsilon \rightarrow 0$, where δ_v is the Dirac delta distribution in the velocity space. Therefore, for the first term in (38),

$$\iiint_S \partial_t f_\varepsilon \chi_\varepsilon dx dv dt \rightarrow \int_0^T \int_{\mathbb{R}^d} \partial_t \rho \phi dx dt.$$

The right-hand side of (38) converges to zero. It remains to treat the second term in (38). Integrating by parts and using the equation for χ_ε , we obtain

$$\begin{aligned} & \varepsilon^{1-\theta} \iiint_S v \cdot \nabla_x (F_\varepsilon + \varepsilon^{\theta/2} r_\varepsilon) \chi_\varepsilon dx dv dt \\ &= -\varepsilon^{1-\theta} \iiint_S v \cdot \nabla_x \chi_\varepsilon F_\varepsilon dx dv dt - \varepsilon^{1-\theta/2} \iiint_S v \cdot \nabla_x \chi_\varepsilon r_\varepsilon dx dv dt \\ &= -\varepsilon^{-\theta} \iiint_S \nu F_\varepsilon (\chi_\varepsilon - \phi) dx dv dt - \varepsilon^{1-\theta/2} \iiint_S v \cdot \nabla_x \chi_\varepsilon r_\varepsilon dx dv dt. \end{aligned}$$

Since $1-\theta/2 > 0$, the last term converges (formally) to zero. By [1, Prop. 4.2] and $E_\varepsilon \rightarrow E$, also the first term converges:

$$\begin{aligned} & -\varepsilon^{-\theta} \iiint_S \nu(v) F(v, \varepsilon^{\theta-1} E_\varepsilon(x, t)) (\chi_\varepsilon - \phi) dx dv dt \\ & \rightarrow \int_0^T \int_{\mathbb{R}^d} (\kappa(-\Delta)^{\theta/2} \phi + (DE) \cdot \nabla_x \phi) \rho dx dt, \end{aligned}$$

where $\kappa > 0$ depends only on d, γ, θ , and the behavior of $\nu(v)$ as $|v| \rightarrow \infty$, and

$$D = \int_{\mathbb{R}^d} \lambda(v) \otimes v dv, \quad \text{where } Q(\lambda) = \nabla_v M.$$

Therefore, integrating by parts, the limit $\varepsilon \rightarrow 0$ in (38) leads to

$$\int_0^T \int_{\mathbb{R}^d} \partial_t \rho \phi dx dt = \int_0^T \int_{\mathbb{R}^d} (-\kappa(-\Delta)^{\theta/2} \rho + \text{div}(DE\rho)) \phi dx dt,$$

which is the weak formulation of the first equation in (1) since $E = \nabla_x \psi$.

Finally, we remark that when the scattering rate is constant, $\sigma(v, v') = 1$, the collision operator simplifies to $Q(f) = M \int_{\mathbb{R}^d} f dv - f$, and the unique solution to $Q(\lambda) = \nabla_v M$, $\int_{\mathbb{R}^d} \lambda dv = 0$ equals $\lambda = -\nabla_v M$. Consequently, $D_{ij} = 0$ for $i \neq j$ and $D_{ii} = -\int_{\mathbb{R}^d} \partial_{v_i} M v_i dv = \int_{\mathbb{R}^d} M dv = 1$, so D equals the unit matrix.

APPENDIX C. GENERAL DRIFT TERMS

The drift-diffusion equation of [1] is of the form (see Appendix B)

$$(39) \quad \partial_t \rho + \kappa(-\Delta)^{\theta/2} \rho = \text{div}(DE\rho) \quad \text{in } \mathbb{R}^d,$$

where $\kappa > 0$ and $D \in \mathbb{R}^{d \times d}$ is a (constant) matrix. We suppose that $E = \nabla\psi$ and the potential ψ is a solution to the Poisson equation $-\Delta\psi = \rho$. In this section, we will illustrate that our method can be applied only when $D = aI + B$, where $a > 0$ and B is skew-symmetric.

This restriction appears when deriving $L^q(\mathbb{R}^d)$ estimates for $\rho(t)$. Indeed, multiplying (39) by ρ^{q-1} for some $q > 1$ and using the Stroock-Varopoulos inequality as in Appendix A, we obtain

$$(40) \quad \begin{aligned} \frac{1}{q} \frac{d}{dt} \|\rho(t)\|_{L^q(\mathbb{R}^d)}^q + \frac{2\kappa}{q} \|(-\Delta)^{\theta/4}(\rho^{q/2})\|_{L^2(\mathbb{R}^d)}^2 &\leq -\frac{q-1}{q} \int_{\mathbb{R}^d} \nabla(\rho^q) \cdot (D\nabla\psi) \, dx \\ &= \frac{q-1}{q} \int_{\mathbb{R}^d} \rho^q \operatorname{div}(D\nabla\psi) \, dx. \end{aligned}$$

If D equals the unit matrix, the last integral has a sign:

$$\int_{\mathbb{R}^d} \rho^q \operatorname{div}(D\nabla\psi) \, dx = \int_{\mathbb{R}^d} \rho^q \Delta\psi \, dx = - \int_{\mathbb{R}^d} \rho^{q+1} \, dx \leq 0.$$

For general matrices D , we argue as follows. Let $R = \nabla(-\Delta)^{-1/2}$ be the Riesz transform, which can be also characterized as a Fourier multiplier, $Rf = \mathcal{F}^{-1}[-i(\xi/|\xi|)\widehat{f}]$. It has the property $R \cdot R = -I$. Then, since D is a constant matrix and $\psi = (-\Delta)^{-1}\rho$,

$$\begin{aligned} \operatorname{div}(D\nabla\psi) &= \nabla \cdot (D\nabla(-\Delta)^{-1}\rho) = \nabla(-\Delta)^{-1/2} \cdot (D\nabla(-\Delta)^{-1/2}\rho) \\ &= R \cdot (DR\rho) = \frac{1}{d} \operatorname{tr}(D)(R \cdot R)\rho - R \cdot \left(\frac{1}{d} \operatorname{tr}(D) - D \right) R\rho. \end{aligned}$$

We claim that the last term can be written as

$$R \cdot \left(\frac{1}{d} \operatorname{tr}(D) - D \right) R\rho = c_d \int_{\mathbb{R}^d} \left(\frac{1}{d} \operatorname{tr}(D)|y|^2 - y \cdot (Dy) \right) \frac{\rho(x-y)}{|y|^{d+2}} \, dy,$$

for some constant $c_d > 0$ only depending on the dimension d . Set $P(\xi) = \operatorname{tr}(D)|\xi|^2/d + \xi \cdot (D\xi)$. Since R is a Fourier multiplier, we have

$$\begin{aligned} R \cdot \left(\frac{1}{d} \operatorname{tr}(D) - D \right) R\rho &= -\mathcal{F}^{-1} \left[\left(\frac{1}{d} \operatorname{tr}(D) + \frac{\xi}{|\xi|} \cdot D \frac{\xi}{|\xi|} \right) \widehat{\rho} \right] = -\mathcal{F}^{-1} \left[\frac{P(\xi)}{|\xi|^2} \widehat{\rho} \right] \\ &= -\mathcal{F}^{-1} \left[c_d \mathcal{F} \left[\frac{P(x)}{|x|^{d+2}} \right] \mathcal{F}[\rho] \right] = -c_d \frac{P(\xi)}{|\xi|^{d+2}} * \rho \\ &= -c_d \int_{\mathbb{R}^d} \frac{P(\xi)}{|\xi|^{d+2}} \rho(x-\xi) \, d\xi, \end{aligned}$$

where we used $c_d \mathcal{F}[P(x)/|x|^{d+2}] = P(\xi)/|\xi|^2$ (see [34, Theorem 5, Section III.3]) and the Fourier convolution formula. This shows that (40) can be written as

$$(41) \quad \begin{aligned} \frac{1}{q} \frac{d}{dt} \|\rho(t)\|_{L^q(\mathbb{R}^d)}^q + \frac{2\kappa}{q} \|(-\Delta)^{\theta/4}(\rho^{q/2})\|_{L^2(\mathbb{R}^d)}^2 &\leq -\frac{q-1}{q} \frac{\operatorname{tr}(D)}{d} \int_{\mathbb{R}^d} \rho^{q+1} \, dx \\ &\quad - \frac{q-1}{q} c_d \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \rho^q(x) \left(\frac{1}{d} \operatorname{tr}(D)|y|^2 - y \cdot (Dy) \right) \frac{\rho(x-y)}{|y|^{d+2}} \, dy \, dx. \end{aligned}$$

If $\text{tr}(D) \geq 0$, the first integral on the right-hand side is nonpositive. The second integral is nonpositive if $y \cdot Dy \leq \text{tr}(D)|y|^2/d$ for all $y \in \mathbb{R}^d$. However, Lemma 11 below shows that under this condition, D can be decomposed as $D = aI + B$, where $a > 0$ and B is skew-symmetric. Consequently, we are not able to treat general drift matrices D . Clearly, even if the second integral on the right-hand side of (41) is positive, it may happen that it is absorbed by the first integral such that the right-hand side is still nonpositive, but we are not able to prove this.

Lemma 11. *Let $D = (D_{ij}) \in \mathbb{R}^{d \times d}$ be a matrix such that $x \cdot Dx \leq \text{tr}(D)|x|^2/d$ for all $x \in \mathbb{R}^d$. Then there exists $a > 0$ and a skew-symmetric matrix such that $D = aI + B$.*

Proof. We can write $D = A + B$, where $A = (D + D^T)/2$ is symmetric and $B = (D - D^T)/2$ is skew-symmetric. We show that $x \cdot Ax = a|x|^2$ for all $x \in \mathbb{R}^d$ for some $a > 0$. Since A is symmetric and real, we may assume (by the spectral theorem), without loss of generality, that A is a diagonal matrix, $A = \text{diag}(a_1, \dots, a_d)$. Furthermore, by homogeneity, it is sufficient to show this result for all $x \in \mathbb{R}^d$ with $|x| = 1$. Then $x \cdot Dx \leq \text{tr}(D)/d$ is equivalent to

$$\sum_{i=1}^d a_i x_i^2 \leq \frac{1}{d} \sum_{j=1}^d a_j$$

or, since $\sum_{i=1}^d x_i^2 = 1$,

$$\sum_{i=1}^d \left(a_i - \frac{1}{d} \sum_{j=1}^d a_j \right) x_i^2 \leq 0.$$

Choosing $x_i = \delta_{ik}$, we see that $a_i - (1/d) \sum_{j=1}^d a_j \leq 0$, but this is only possible if $a := a_i = a_j$ for all $i \neq j$. This proves the lemma. \square

REFERENCES

- [1] Aceves-Sanchez, P., Mellet, M., Anomalous diffusion limit for a linear Boltzmann equation with external force field, *Math. Models Meth. Appl. Sci.* **27** (2017), 845–878.
- [2] Applebaum, D., *Lévy Processes and Stochastic Calculus*, Cambridge Studies Adv. Math. **93**, Cambridge University Press, Cambridge, 2004.
- [3] Arnold, A., Markowich, P., Toscani, G., Unterreiter, A., On convex Sobolev inequalities and the rate of convergence to equilibrium for Fokker-Planck type equations, *Commun. Part. Diff. Eqs.* **26** (2001), 43–100.
- [4] Bardos, C., Golse, F., Moyano, I., Linear Boltzmann equation and fractional diffusion, preprint arXiv:1708.09791, 2017.
- [5] Bakry, D., Emery, M., Diffusions hypercontractives, *Séminaire de probabilités XIX, 1983/84*, Lecture Notes Math. **1123**, pp. 177–206, Springer, Berlin, 1985.
- [6] Biler, P., Karch, G., Generalized Fokker-Planck equations and convergence to their equilibria, in: *Evolution Equations (Warsaw 2001)*, pp. 307–318, Banach Center Publ. **60**, Warsaw, 2003.
- [7] Biler, P., Karch, G., Woyczyński, W. A., Critical nonlinearity exponent and selfsimilar asymptotics for Lévy conservation laws, *Ann. H. Poincaré* **18** (2001), 613–637.
- [8] Carrillo, J.A., Jüngel, A., Markowich, P.A., Toscani, G., Unterreiter, A., Entropy dissipation methods for degenerate parabolic problems and generalized Sobolev inequalities, *Monatsh. Math.* **133** (2001), 1–82.

- [9] Chafaï, D., Entropies, convexity, and functional inequalities: On Φ -entropies and Φ -Sobolev inequalities, *J. Math. Kyoto Univ.* **44** (2004), 325–363.
- [10] P. Clavin, Instabilities and nonlinear patterns of overdriven detonations in gases, in: H. Berestycki and Y. Pomeau (eds.), *Nonlinear PDEs in Condensed Matter and Reactive Flows*, Kluwer, 2002, 49–97.
- [11] Constantin, P., Majda, A.J., Tabak, E., Formation of strong fronts in the 2-D quasigeostrophic thermal active scalar, *Nonlinearity* **7** (1994), 1495–1533.
- [12] Córdoba, A., Córdoba, D., A pointwise estimate for fractionary derivatives with applications to partial differential equations, *Proc. Natl. Acad. Sci. USA* **100** (2003), 15316–15317.
- [13] Córdoba, A., Córdoba, D., A maximum principle applied to quasi-geostrophic equations, *Commun. Math. Phys.* **249** (2004), 511–528.
- [14] Dong, H., Du, D., Li, D., Finite time singularities and global well-posedness for fractal Burgers equations, *Indiana Math. J.* **58** (2009), 807–821.
- [15] Droniou, J., Imbert, C., Fractal first order partial differential equations, *Arch. Ration. Mech. Anal.* **182** (2006), 299–331.
- [16] Escudero, C., The fractional Keller–Segel model, *Nonlinearity* **19** (2006), 2909–2918.
- [17] Gentil, I., Imbert, C., The Lévy-Fokker-Planck equation: Φ -entropies and convergence to equilibrium, *Asymptot. Anal.* **59** (2008), 125–138.
- [18] Granero-Belinchón, R., On a drift-diffusion system for semiconductor devices, *Ann. H. Poincaré* **17** (2016), 3473–3498.
- [19] Iwabuchi, T., Global solutions for the critical Burgers equation in the Besov spaces and the large time behavior, *Ann. H. Poincaré Anal. Non Linéaire* **32** (2015), 687–713.
- [20] Ju, N., Existence and uniqueness of the solution to the dissipative 2D quasi-geostrophic equations in the Sobolev space, *Commun. Math. Phys.* **251** (2004), 365–376.
- [21] Jüngel, A., *Transport Equations for Semiconductors*, Lecture Notes Phys. **773**, Springer, Berlin, 2009.
- [22] Jüngel, A., *Entropy Methods for Diffusive Partial Differential Equations*, BCAM Springer Briefs, Springer, 2016.
- [23] Kobayashi, R., Kawashima, S., Decay estimates and large time behavior of solutions to the drift-diffusion system, *Funkcialaj Ekvacioj* **51** (2008), 371–394.
- [24] Kolokoltsov, V., *Markov Processes, Semigroups and Generators*, De Gruyter, Berlin/New York, 2011.
- [25] Kurokiba, M., Ogawa, T., Well-posedness for the drift-diffusion system in L^p arising from the semiconductor device simulation, *J. Math. Anal. Appl.* **342** (2008), 1052–1067.
- [26] Li, D., Rodrigo, J. L., Zhang, X., Exploding solutions for a nonlocal quadratic evolution problem, *Rev. Mat. Iberoamer.* **26** (2010), 295–332.
- [27] Liskevich, V. A., Semenov, Yu. A., Some problems on Markov semigroups, in: *Schrödinger Operators, Markov Semigroups, Wavelet Analysis, Operator Algebras*, Math. Top., vol. **11**, Partial Differential Equations, Akademie Verlag, Berlin, 1996.
- [28] Mellet, A., Fractional diffusion limit for collisional kinetic equations: a moments method, *Indiana Univ. Math. J.* **59** (2010), 1333–1360.
- [29] Mellet, A., Mischler, S., Mouhot, C., Fractional diffusion limit for collisional kinetic equations, *Arch. Ration. Mech. Anal.* **199** (2011), 493–525.
- [30] Nagai, T., Convergence to self-similar solutions for a parabolic-elliptic system of drift-diffusion type in \mathbb{R}^2 , *Adv. Diff. Eqs.* **16** (2011), 839–866.
- [31] Ogawa, T., Yamamoto, M., Asymptotic behavior of solutions to drift-diffusion system with generalized dissipation, *Math. Models Meth. Appl. Sci.* **19** (2009), 939–967.
- [32] Poupaud, F., Runaway phenomena and fluid approximation under high fields in semiconductor kinetic theory, *Z. Angew. Math. Mech.* **72** (1992), 359–372.
- [33] Sato, K., *Lévy processes and infinitely divisible distributions*, Cambridge Studies in Advanced Mathematics **68**, 2nd edition, Cambridge University Press, Cambridge, 2013.

- [34] Stein, E.M., Singular Integrals and Differentiability Properties of Functions, Vol. **30**, Princeton University Press, Princeton, 1970.
- [35] Sugiyama, Y., Yamamoto, M., Kato, K., Local and global solvability and blow up for the drift-diffusion equation with the fractional dissipation with the critical space, *J. Diff. Eqs.* **258** (2015), 2983–3010.
- [36] Tristani, I., Fractional Fokker-Planck equation, *Commun. Math. Sci.* **13** (2015), 1243–1260.
- [37] Vázquez, J. L., Asymptotic behaviour for the fractional heat equation in the Euclidean space, *Complex Variables Ellipt. Eqs.*, 2017. Online first, <https://doi.org/10.1080/17476933.2017.1393807>.
- [38] W. Woyczyński, Lévy processes in the physical sciences, in: O. Barndorff-Nielsen, S. Resnick, and T. Mikosch (eds.), *Lévy Processes: Theory and Applications*, Birkhäuser, Boston, 2001, 241–266.
- [39] Wu, L., A new modified logarithmic Sobolev inequality for Poisson point processes and several applications, *Prob. Theory Related Fields* **118** (2000), 427–438.
- [40] Yamamoto, M., Large-time behavior of solutions to the drift-diffusion equation with fractional dissipation, *Diff. Int. Eqs.* **25** (2012), 731–758.
- [41] Yamamoto, M., Kato, K., Sugiyama, Y., Existence and analyticity of solutions to the drift-diffusion equation with critical dissipation, *Hiroshima Math. J.* **44** (2014), 275–313.
- [42] Yamamoto, M., Sugiyama, Y., Asymptotic expansion of solutions to the drift-diffusion equation with fractional diffusion, *Nonlin. Anal.* **141** (2016), 57–87.
- [43] Yamamoto, M., Sugiyama, Y., Asymptotic behavior of solutions to the drift-diffusion equation with critical dissipation, *Ann. H. Poincaré* **17** (2016), 1331–1352.
- [44] Zhao, J., Liu, Q., On the Cauchy problem for the fractional drift-diffusion system in critical Besov spaces, *Appl. Anal.* **93** (2014), 1431–1450.

FACULTY OF MATHEMATICS, UNIVERSITY OF VIENNA, OSKAR-MORGENSTERN-PLATZ 1, 1090 WIEN, AUSTRIA

E-mail address: `franz.achleitner@univie.ac.at`

INSTITUTE FOR ANALYSIS AND SCIENTIFIC COMPUTING, VIENNA UNIVERSITY OF TECHNOLOGY, WIEDNER HAUPTSTRASSE 8–10, 1040 WIEN, AUSTRIA

E-mail address: `juengel@tuwien.ac.at`

GRADUATE SCHOOL OF SCIENCE AND TECHNOLOGY, NIIGATA UNIVERSITY, NIIGATA, 950-2181, JAPAN

E-mail address: `masakazu@eng.niigata-u.ac.jp`