

CONVERGENCE OF A FINITE VOLUME SCHEME AND DISSIPATIVE MEASURE-VALUED-STRONG STABILITY FOR A HYPERBOLIC-PARABOLIC CROSS-DIFFUSION SYSTEM

KATHARINA HOPF AND ANSGAR JÜNGEL

ABSTRACT. This article is concerned with the approximation of hyperbolic-parabolic cross-diffusion systems modeling segregation phenomena for populations by a fully discrete finite-volume scheme. It is proved that the numerical scheme converges to a dissipative measure-valued solution of the PDE system and that, whenever the latter possesses a strong solution, the convergence holds in the strong sense. Furthermore, the “parabolic density part” of the limiting measure-valued solution is atomic and converges to its constant state for long times. The results are based on Young measure theory and a weak-strong stability estimate combining Shannon and Rao entropies. The convergence of the numerical scheme is achieved by means of discrete entropy dissipation inequalities and an artificial diffusion, which vanishes in the continuum limit.

1. INTRODUCTION

The segregation of multi-species populations can be modeled at a macroscopic level by cross-diffusion equations. Segregation typically requires the associated diffusion matrix to have a nontrivial kernel. In this situation, solutions may have spatial discontinuities; see, e.g., [2] for a two-species model. The segregation models have been derived, for an arbitrary number of species, from interacting particle systems in a mean-field-type limit [9]. The class considered here has recently been found to possess a symmetric hyperbolic-parabolic structure [17]. In this paper, we establish the global existence of dissipative measure-valued solutions as a limit of finite-volume approximations, the uniqueness of strong solutions among dissipative measure-valued solutions, and a result on the long-time asymptotic behavior.

Date: March 31, 2023.

2020 Mathematics Subject Classification. 35M33, 35R06, 65M12, 92D25.

Key words and phrases. Cross diffusion, segregating populations, parametrized measure, dissipative measure-valued solution, finite-volume method, entropy method, weak-strong uniqueness, long-time behavior.

The second author acknowledges partial support from the Austrian Science Fund (FWF), grants P33010 and F65. He thanks Antoine Zurek (Compiègne) and Flore Nabet (Palaiseau) for helpful discussions and pointing out the papers [40, 41]. This work has received funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme, ERC Advanced Grant no. 101018153.

1.1. **Equations.** The segregation cross-diffusion equations for the vector $u = (u_1, \dots, u_n)$ of the population densities u_i are systems of continuity equations

$$(1) \quad \partial_t u_i + \operatorname{div}(u_i \mathbf{v}_i) = 0, \quad \mathbf{v}_i = -\nabla p_i(u), \quad \text{in } \Omega, \quad t > 0, \quad i = 1, \dots, n,$$

where $p_i(u) = \sum_{j=1}^n a_{ij} u_j$ and $\Omega \subset \mathbb{R}^d$ ($d \geq 1$) is a bounded Lipschitz domain, supplemented with the no-flux boundary and initial conditions

$$(2) \quad u_i \nabla p_i(u) \cdot \nu = 0 \quad \text{on } \partial\Omega, \quad t > 0, \quad u_i(0) = u_i^{\text{in}} \quad \text{in } \Omega, \quad i = 1, \dots, n,$$

where ν denotes the exterior unit normal vector to $\partial\Omega$. The variables (u_i) represent, for instance, densities of animal populations [2], healthy and tumor cell densities [39], or heights of thin fluid layers [14, 37].

The parameters $a_{ij} \geq 0$ are assumed to satisfy the following two conditions: The matrix $A = (a_{ij}) \in \mathbb{R}^{n \times n}$ is semistable, i.e., the real parts of all its eigenvalues are nonnegative, and it satisfies the detailed-balance condition, i.e., there exist $\pi_1, \dots, \pi_n > 0$ such that

$$(3) \quad \pi_i a_{ij} = \pi_j a_{ji} \quad \text{for all } i, j = 1, \dots, n, \quad i \neq j.$$

These equations can be recognized as the detailed-balance condition for the Markov chain associated to A , and the vector (π_i) is an invariant measure. Under condition (3), the change of variables $u_j \mapsto \pi_j u_j := \tilde{u}_j$ brings the equation in the form $\partial_t \tilde{u}_i = \operatorname{div}(\tilde{u}_i \nabla (B \tilde{u})_i)$, where the matrix $B = (a_{ij} \pi_j^{-1})_{ij}$ is symmetric and positive semidefinite. Thus, from now on we consider, without loss of generality, the equations

$$(4) \quad \partial_t u_i = \operatorname{div}(u_i \nabla p_i(u)), \quad p_i(u) = \sum_{j=1}^n b_{ij} u_j \quad \text{in } \Omega, \quad t > 0, \quad i = 1, \dots, n,$$

where $B = (b_{ij}) \in \mathbb{R}^{n \times n}$ is symmetric positive semidefinite and $b_{ij} \geq 0$ for all $i, j = 1, \dots, n$. We note that if $b_{ii} = 0$ for some i , then $b_{ji} = b_{ij} = 0$ for $j = 1, \dots, n$ due to the positive semidefiniteness of B . Thus, in this case, the dynamics of u_i become trivial and the i th species can be removed from the system. We may therefore further assume that $b_{ii} > 0$ for all $i = 1, \dots, n$. If $\operatorname{rank} B = n$ and $u_i > 0$ for all $i = 1, \dots, n$, equation (4) is parabolic in the sense of Petrovskii, which at the linear level is a minimal condition for the generation of an analytic semigroup on $L^p(\Omega)$ [1]. The existence of global weak solutions in the case $\operatorname{rank} B = n$ was investigated in [32, Theorem 17]. If B has a nontrivial kernel, it is positive definite only on the subspace $(\ker B)^\perp$, and we lose the parabolic structure. This is the situation we are primarily concerned with in this paper.

1.2. **State of the art.** Equations (1) with nontrivial kernel have been studied in the literature in special cases. The first work is [2], where the global existence of segregated solutions for two species in one space dimension with $a_{11} = a_{12} = 1$ and $a_{21} = a_{22} = k > 0$ was shown. This result relies on a change to mass variables. The analysis was generalized in [3] to several space dimensions if $k = 1$. The idea is to introduce new variables $w_1 = u_1 + u_2$ and $w_2 = u_1 / (u_1 + u_2)$. It turns out that w_1 solves a porous-medium equation with quadratic nonlinearity and w_2 solves a transport equation, demonstrating the hyperbolic-parabolic nature of the system. The same idea was used in [27] for a related system, where $p_i(u) = (u_1 + u_2)^\gamma$ for $\gamma > 1$. Notice that the choice $k = 1$ means that the corresponding

velocity fields \mathbf{v}_i in (1) are independent of i , so that the motion of the two species is governed by a single velocity field.

The existence of an infinite family of minimizers of the entropy (or free energy) functional for different local and nonlocal variants was proved in [6], showing that both segregation and mixing of species is possible. If the pressure is the variational derivative of a certain functional, one may formulate (1) for $n = 2$ as a formal gradient flow. This property has been exploited in [6, 15] to prove the convergence of a minimizing scheme.

The one-velocity two-species case was generalized to an arbitrary number of species in [18], proving the global existence of classical and weak solutions by decomposing the system into one decoupled porous-medium equation and $n - 1$ transport equations. This approach was generalized in [17] to the case of multiple velocity fields with associated diffusion matrices of arbitrary rank $r \in \{1, \dots, n\}$ to show the local-in-time existence of classical solutions. Segregating solutions for one-velocity multi-species reactive systems were constructed in [30].

There exist related cross-diffusion models with rank-deficient diffusion matrices in the literature, for instance the Maxwell–Stefan equations for fluid mixtures [4], where the diffusion matrix has a one-dimensional kernel. In contrast to the present problem, the kernel can be removed by taking into account the volume-filling assumption $\sum_{i=1}^n u_i = 1$, which allows one to reduce the system for the densities u_1, \dots, u_n to a parabolic one for the variables u_1, \dots, u_{n-1} via $u_n = 1 - \sum_{i=1}^{n-1} u_i$ [33].

The analysis and the convergence of approximation schemes to equations (4) for general rank-deficient matrices A is challenging, since the decomposition of the parabolic and hyperbolic parts is involved. Moreover, in view of the results of [2], we cannot expect weak solutions in $H^1(\Omega)$, and the hyperbolic part makes it difficult to obtain (entropy) solutions in the distributional sense. In the present paper, we choose to enlarge the solution space by considering dissipative measure-valued solutions, which allow us to encode information about the oscillation properties of the approximate solutions.

DiPerna introduced the concept of (entropy) *measure-valued solutions* to conservation laws [16]. In this framework, solutions are no longer integrable functions but Young measures (parametrized probability measures), which are able to capture the limiting behavior of sequences of oscillating functions. This concept is based on an earlier work by Tartar [44], who characterized weak limits of sequences of bounded functions. Due to the lack of uniqueness results, the framework of measure-valued solutions does not allow one to identify the physically relevant solutions, and further structural conditions on the solutions are necessary.

One idea to resolve this issue is to require an integrated form of the entropy or energy inequality, which leads to the concept of *dissipative solutions*. It has been introduced by P.-L. Lions [38, Sec. 4.4] in the context of the incompressible Euler equations. In [5] it is shown that dissipative measure-valued solutions to the incompressible Euler equations enjoy the weak–strong uniqueness property, i.e., the dissipative measure-valued solution is atomic and coincides with the strong or classical solution of the same initial-value problem if the latter exists. This idea was further applied to models from polyconvex elastodynamics [11],

to the compressible Euler and Navier–Stokes equations [28, 21], to hyperbolic–parabolic systems in thermoviscoelasticity [10], and to various other, mainly fluid mechanical models.

In the present paper, we obtain dissipative measure-valued solutions to (4), (2) by passing to the limit from discrete *finite-volume solutions*. We further show that they enjoy the weak–strong uniqueness property (in the sense of measure-valued–strong uniqueness), which entails important consequences for the numerical approximation. Indeed, one may expect that reasonable structure-preserving approximation schemes generate a dissipative measure-valued solution. If this measure-valued solution turns out to be atomic, i.e. taking the form of a Dirac measure at each point in space-time, Young measure theory implies that the underlying approximate solutions converge in the strong sense. This idea has, for instance, been exploited in the proof of the convergence of finite-volume-type schemes for the compressible Navier–Stokes and Euler equations [22, 23]. For a further discussion on the use of measure-valued solutions in the numerical context, we refer to [24].

The novelty of this paper is the analysis of equations (4) with general rank-deficient matrices B by combining the measure-valued framework, entropy methods, and finite-volume schemes.

1.3. Key tools and main results. The analysis of (4) is based on the observation that the system possesses two Lyapunov functionals, respectively, the Shannon and Rao entropies

$$(5) \quad H_S(u) = \int_{\Omega} h_S(u) dx, \quad h_S(u) = \sum_{i=1}^n (u_i (\log u_i - 1) + 1),$$

$$(6) \quad H_R(u) = \int_{\Omega} h_R(u) dx, \quad h_R(u) = \frac{1}{2} \sum_{i,j=1}^n b_{ij} u_i u_j.$$

The Shannon (-Boltzmann) entropy is related to the thermodynamic entropy of the system, while the Rao entropy measures the functional diversity of the species [43].

The functionals have two important properties. First, a computation shows that, along smooth solutions to (4), (2),

$$(7) \quad \frac{dH_S}{dt}(u) + \sum_{i,j=1}^n \int_{\Omega} b_{ij} \nabla u_i \cdot \nabla u_j dx = 0,$$

$$(8) \quad \frac{dH_R}{dt}(u) + \sum_{i=1}^n \int_{\Omega} u_i |\nabla p_i(u)|^2 dx = 0.$$

Since the matrix B is positive semidefinite, the Shannon entropy dissipation term (the integral term in (7)) is nonnegative and consequently, $t \mapsto H_S(u(t))$ is nonincreasing. The expression $p_i(u)$ can be interpreted as the i th partial pressure and $-\nabla p_i(u)$ as the i th partial velocity (by Darcy’s law). Thus, we may interpret the Rao entropy dissipation integral as the total kinetic energy of the system, and $t \mapsto H_R(u(t))$ is also nonincreasing.

Second, the Shannon entropy density h_S is strictly convex on $(0, \infty)^n$, but has only logarithmically superlinear growth as $\sum_{i=1}^n u_i \rightarrow \infty$. This growth appears to be insufficient to derive a weak-strong stability estimate based on the associated Bregman distance $h_S(u|v) := h_S(u) - h_S(v) - h'_S(v) \cdot (u-v)$, since we do not generally have a priori bounds in $L^\infty(\Omega)$ for solutions to (4). However, the Rao entropy density h_R , which is convex but not necessarily strictly convex, has quadratic growth as $|u| \rightarrow \infty$, $u \in (0, \infty)^n$, as soon as $b_{ij} \geq 0$ and $b_{ii} > 0$ for all $i, j = 1, \dots, n$. By considering the sum of the Shannon and the Rao entropy densities, we obtain a strictly convex function whose associated Bregman distance turns out to grow fast enough as $\sum_i u_i \rightarrow \infty$ to deduce a closed weak-strong stability estimate via Gronwall's lemma.

Identities (7)–(8) provide estimates for u_i in $L^\infty(0, T; L^2(\Omega))$ and for $(Bu)_i$ in $L^2(0, T; H^1(\Omega))$, $T > 0$. Thus, if B is rank-deficient, these bounds do not provide gradient estimates for the whole vector u . Notice that the weak convergence for u_m and $\nabla p_i(u_m) = \nabla(Bu_m)_i$ in $L^2(\Omega \times (0, T))$, which may be expected for suitable approximating sequences u_m , does not allow us to identify the weak limit of $u_{m,i} \nabla(Bu_m)_i$. This issue is overcome by a suitable concept of dissipative measure-valued solutions. Let us mention that the estimates coming from (8) allow us to control $u_{m,i} \nabla(Bu_m)_i$ in $L^2(0, T; L^{4/3}(\Omega))$, thus ruling out potential concentrations in this term.

Before giving the definition of the measure-valued solutions, we introduce some notation. We rewrite equation (4) as

$$\partial_t u_i = \operatorname{div}(u_i \nabla(Bu)_i), \quad i = 1, \dots, n,$$

and set $L := \ker B \subsetneq \mathbb{R}^n$. Then $L^\perp = \operatorname{ran} B \supsetneq \{0\}$. Let P_{L^\perp} be the projection onto L^\perp and set $\hat{s} := P_{L^\perp} s$ for any $s \in \mathbb{R}^n$. Any vector-valued function u is written as $u = (u_1, \dots, u_n)$. We define $\mathbb{R}_{\geq} = [0, \infty)$ and let $\mathcal{P}(W)$ be the space of probability measures on

$$W := \mathbb{R}_{\geq}^n \times (L^\perp)^d.$$

The space $L_w^\infty(\Omega \times [0, \infty); \mathcal{P}(W))$ is the set of weakly* measurable, essentially bounded functions of $\Omega \times [0, \infty)$ taking values in $\mathcal{P}(W)$. We henceforth use the notation

$$\langle \nu, f(s, p) \rangle := \int_W f(s, p) \, d\nu(s, p) \quad \text{for } \nu \in \mathcal{P}(W), f \in C_0(W),$$

and let $\Omega_T = \Omega \times (0, T)$ for $T > 0$.

Definition 1 (Dissipative measure-valued solution). *Suppose that $u^{\text{in}} \in L^2(\Omega; \mathbb{R}_{\geq}^n)$. We call a parametrized measure*

$$\mu \in L_w^\infty(\Omega \times [0, \infty); \mathcal{P}(W))$$

with barycenters $u := \langle \mu, s \rangle$, $y := \langle \mu, p \rangle$ a dissipative measure-valued solution to (4), (2) if the following is satisfied for all $T > 0$:

- *Regularity: It holds $u_i \geq 0$ a.e. in Ω_T for $i = 1, \dots, n$ and*

$$u_i \in L^\infty(0, \infty; L^2(\Omega)), \quad \partial_t u \in L^2(0, \infty; W^{1,4}(\Omega)^*), \quad y \in L^2(\Omega_T; (L^\perp)^d), \quad y = \nabla \hat{u}.$$

- Evolution equation: *It holds for all $i = 1, \dots, n$ and $\phi \in C_c^1(\overline{\Omega} \times [0, T])$ that*

$$(9) \quad \int_0^T \int_{\Omega} u_i \partial_t \phi dx dt + \int_{\Omega} u_i^{\text{in}} \phi(0) dx = \int_0^T \int_{\Omega} \langle \mu_{x,t}, s_i(Bp)_i \rangle \cdot \nabla \phi dx dt.$$

- Shannon and Rao entropy inequalities: *It holds for a.e. $t > 0$ that*

$$(10) \quad H_S^{\text{mv}}(u(t)) + \int_0^t \int_{\Omega} |\nabla(B^{1/2}\widehat{u})|^2 dx d\tau \leq H_S(u^{\text{in}}),$$

$$(11) \quad H_R(u(t)) + \sum_{i=1}^n \int_0^t \int_{\Omega} \langle \mu_{x,t}, s_i |Bp_i|^2 \rangle dx d\tau \leq H_R(u^{\text{in}}),$$

where H_S and H_R are defined in (5)–(6) and $H_S^{\text{mv}}(u(t)) := \int_{\Omega} \langle \mu_{x,t}, h_S(s) \rangle dx$.

We show in Remark 5 that the definition of dissipative measure-valued solutions is consistent with the definition of weak solutions. Our main results can be sketched as follows; we refer to Section 2.5 for the precise statements.

- Existence of finite-volume approximations: There exists a sequence of approximate solutions (u_m) , where $m \in \mathbb{N}$ indicates the fineness of the mesh, to an implicit Euler finite-volume scheme. The numerical scheme preserves the structure of the equations, namely nonnegativity, conservation of mass, and entropy dissipation; see Theorem 3.
- Existence of global dissipative measure-valued solutions: Any Young measure μ generated by (u_m) is a dissipative measure-valued solution to (4), (2) in the sense of Definition 1, which further satisfies (26); see Theorem 4. For this result, we need to include some artificial diffusion in the scheme, which vanishes in the limit $m \rightarrow \infty$.
- Weak–strong uniqueness: If v is a positive classical solution to (4), (2) with initial datum $v(0) = u^{\text{in}}$ and μ is a dissipative measure-valued solution to (4), (2), then $\mu_{x,t} = \delta_{v(x,t)} \otimes \delta_{\nabla v(x,t)}$ for a.e. $(x, t) \in \Omega_T$; see Theorem 7.
- Long-time behavior: The density $\widehat{u}(t) := \langle \mu_{\cdot,t}, \widehat{s} \rangle$ converges strongly in the $L^2(\Omega)$ norm as $t \rightarrow \infty$ to a function $\widehat{u}^* \in L^2(\Omega; [0, \infty)^n)$ satisfying $\int_{\Omega} \widehat{u}^* dx = \int_{\Omega} u^{\text{in}} dx$ and $\nabla(B\widehat{u}^*) = 0$ in Ω ; see Theorem 9.

If equations (4), (2) admit a classical solution, the weak–strong uniqueness property implies that the sequence of finite-volume solutions converges, in the strong L^1 -sense, to this classical solution on the lifespan of the latter; see Corollary 8.

The paper is organized as follows. We introduce the numerical scheme and the precise statements of the theorems in Section 2. The four theorems are proved in Sections 3–6, and we conclude in Appendix A with some auxiliary lemmas.

2. NUMERICAL SCHEME AND MAIN RESULTS

First, we introduce the notation necessary to formulate our numerical method. Then we state the numerical scheme and the main results.

2.1. Spatial domain and mesh. Let $d \geq 2$ and let $\Omega \subset \mathbb{R}^d$ be a bounded polygonal domain (or polyhedral if $d \geq 3$). We associate to this domain an admissible mesh, given by (i) a family \mathcal{T} of open polygonal (or polyhedral) control volumes, which are also called cells, (ii) a family \mathcal{E} of edges (or faces if $d \geq 3$), and (iii) a family of points $(x_K)_{K \in \mathcal{T}}$ associated to the control volumes and satisfying [20, Definition 9.1]. This definition implies that the straight line $\overline{x_K x_L}$ between two centers of neighboring cells is orthogonal to the edge (or face) $\sigma = K|L$ between two cells. For instance, Voronoï meshes satisfy this condition [20, Example 9.2]. The size of the mesh is given by $\Delta x = \max_{K \in \mathcal{T}} \text{diam}(K)$. The family of edges \mathcal{E} is assumed to consist of interior edges \mathcal{E}_{int} satisfying $\sigma \subset \Omega$ and boundary edges $\sigma \in \mathcal{E}_{\text{ext}}$ satisfying $\sigma \subset \partial\Omega$. For a given $K \in \mathcal{T}$, \mathcal{E}_K denotes the set of edges of K , splitting into $\mathcal{E}_K = \mathcal{E}_{\text{int},K} \cup \mathcal{E}_{\text{ext},K}$. For any $\sigma \in \mathcal{E}$, there exists at least one cell $K \in \mathcal{T}$ such that $\sigma \in \mathcal{E}_K$.

We need a regularity assumption of the mesh. For given $\sigma \in \mathcal{E}$, we define the distance

$$d_\sigma = \begin{cases} d(x_K, x_L) & \text{if } \sigma = K|L \in \mathcal{E}_{\text{int},K}, \\ d(x_K, \sigma) & \text{if } \sigma \in \mathcal{E}_{\text{ext},K}, \end{cases}$$

where d is the Euclidean distance in \mathbb{R}^d , and the transmissibility coefficient

$$(12) \quad \tau_\sigma = \frac{\tilde{\mathbf{m}}(\sigma)}{d_\sigma},$$

where $\tilde{\mathbf{m}}(\sigma)$ denotes the $(d-1)$ -dimensional Hausdorff measure of σ . We suppose the following mesh regularity condition: There exists $\zeta > 0$ such that for all $K \in \mathcal{T}$ and $\sigma \in \mathcal{E}_K$,

$$(13) \quad d(x_K, \sigma) \geq \zeta d_\sigma.$$

This condition means that the mesh is locally quasi-uniform. We also use the geometric property

$$(14) \quad \sum_{\sigma \in \mathcal{E}_{\text{int},K}} \tilde{\mathbf{m}}(\sigma) d(x_K, \sigma) \leq d\mathbf{m}(K) \quad \text{for any } K \in \mathcal{T},$$

where \mathbf{m} denotes the d -dimensional Lebesgue measure. Inequalities (13) and (14) are needed, for instance, to derive a uniform bound for the discrete time derivative of the approximate solution; see Lemma 13.

2.2. Function spaces. Let $T > 0$, $N \in \mathbb{N}$ and introduce the time step size $\Delta t = T/N$ and the time steps $t_k = k\Delta t$ for $k = 0, \dots, N$. We denote by \mathcal{D} the admissible space-time discretization of $\Omega_T = \Omega \times (0, T)$ composed of an admissible mesh \mathcal{T} and the values $(\Delta t, N)$.

The space of piecewise constant functions is defined by

$$V_{\mathcal{T}} = \left\{ v : \Omega \rightarrow \mathbb{R} : \exists (v_K)_{K \in \mathcal{T}} \subset \mathbb{R}, v(x) = \sum_{K \in \mathcal{T}} v_K \mathbf{1}_K(x) \right\},$$

where $\mathbf{1}_K$ is the characteristic function on K . To define a norm on this space, we define for $v \in V_{\mathcal{T}}$, $K \in \mathcal{T}$, and $\sigma \in \mathcal{E}_K$,

$$v_{K,\sigma} = \begin{cases} v_L & \text{if } \sigma = K|L \in \mathcal{E}_{\text{int},K}, \\ v_K & \text{if } \sigma \in \mathcal{E}_{\text{ext},K}, \end{cases} \quad D_{K,\sigma}v := v_{K,\sigma} - v_K, \quad D_{\sigma}v := |D_{K,\sigma}v|.$$

Let $1 \leq q < \infty$ and $v \in V_{\mathcal{T}}$. The discrete $W^{1,q}(\Omega)$ norm on $V_{\mathcal{T}}$ is given by

$$\|v\|_{1,q,\mathcal{T}} = \left(\|v\|_{0,q,\mathcal{T}}^q + |v|_{1,q,\mathcal{T}}^q \right)^{1/q}, \quad \text{where}$$

$$\|v\|_{0,q,\mathcal{T}}^q = \sum_{K \in \mathcal{T}} \mathbf{m}(K) |v_K|^q, \quad |v|_{1,q,\mathcal{T}}^q = \sum_{\sigma \in \mathcal{E}} \tilde{\mathbf{m}}(\sigma) \left| \frac{D_{\sigma}v}{d_{\sigma}} \right|^q,$$

where $v \in V_{\mathcal{T}}$. If $v = (v_1, \dots, v_n) \in V_{\mathcal{T}}^n$ is a vector-valued function, we write for notational convenience

$$\|v\|_{1,q,\mathcal{T}} = \sum_{i=1}^n \|v_i\|_{1,q,\mathcal{T}}.$$

We associate to the discrete $W^{1,q}$ norm a dual norm with respect to the L^2 inner product:

$$\|v\|_{-1,q,\mathcal{T}} = \sup \left\{ \int_{\Omega} v w dx : w \in V_{\mathcal{T}}, \|w\|_{1,q,\mathcal{T}} = 1 \right\}.$$

Then the following property holds:

$$\left| \int_{\Omega} v w dx \right| \leq \|v\|_{-1,q,\mathcal{T}} \|w\|_{1,q,\mathcal{T}} \quad \text{for all } v, w \in V_{\mathcal{T}}, \quad 1 < p < \infty.$$

Finally, we introduce the space $V_{\mathcal{T},\Delta t}$ of piecewise constant functions with values in $V_{\mathcal{T}}$,

$$V_{\mathcal{T},\Delta t} = \left\{ v : \Omega \times [0, T] \rightarrow \mathbb{R} : \exists (v^k)_{k=1,\dots,N} \subset V_{\mathcal{T}}, \quad v(x, t) = \sum_{k=1}^N v^k(x) \mathbf{1}_{(t_{k-1}, t_k]}(t) \right\},$$

equipped with the discrete $L^2(0, T; H^1(\Omega))$ norm

$$\left(\sum_{k=1}^N \Delta t \|v^k\|_{1,2,\mathcal{T}}^2 \right)^{1/2} \quad \text{for all } v \in V_{\mathcal{T},\Delta t}.$$

2.3. Discrete gradient. The discrete gradient is defined on a dual mesh. For this, we define the cell $T_{K,\sigma}$ of the dual mesh for $K \in \mathcal{T}$ and $\sigma \in \mathcal{E}_K$:

- **Diamond:** Let $\sigma = K|L \in \mathcal{E}_{\text{int},K}$. Then $T_{K,\sigma}$ is that cell whose vertices are given by x_K , x_L , and the end points of the edge σ .
- **Triangle:** Let $\sigma \in \mathcal{E}_{\text{ext},K}$. Then $T_{K,\sigma}$ is that cell whose vertices are given by x_K and the end points of the edge σ .

The union of all diamonds and triangles $T_{K,\sigma}$ equals the domain Ω (up to a set of measure zero). The property that the straight line $\overline{x_K x_L}$ between two neighboring centers of cells is orthogonal to the edge $\sigma = K|L$ implies that

$$\tilde{\mathbf{m}}(\sigma) d(x_K, x_L) = d \mathbf{m}(T_{K,\sigma}) \quad \text{for all } \sigma = K|L \in \mathcal{E}_{\text{int}}.$$

The approximate gradient of $v \in V_{\mathcal{T}, \Delta t}$ is then defined by

$$\nabla^{\mathcal{D}} v(x, t) = \frac{\tilde{\mathbf{m}}(\sigma)}{\mathbf{m}(T_{K, \sigma})} D_{K, \sigma}(v^k) \nu_{K, \sigma} \quad \text{for } x \in T_{K, \sigma}, \quad t \in [t_{k-1}, t_k),$$

where $\nu_{K, \sigma}$ is the unit vector that is normal to σ and points outwards of K .

2.4. Numerical scheme. The initial functions are approximated by $u^0 \in V_{\mathcal{T}}^n$ defined via

$$(15) \quad u_{i, K}^0 = \frac{1}{\mathbf{m}(K)} \int_K u_i^{\text{in}}(x) dx \quad \text{for all } K \in \mathcal{T}, \quad i = 1, \dots, n.$$

Let $u^{k-1} = (u_1^{k-1}, \dots, u_n^{k-1}) \in V_{\mathcal{T}}^n$ be given. Then the values $u_{i, K}^k$ for all $K \in \mathcal{T}$ and $i = 1, \dots, n$ are determined by the implicit Euler finite-volume scheme

$$(16) \quad \mathbf{m}(K) \frac{u_{i, K}^k - u_{i, K}^{k-1}}{\Delta t} + \sum_{\sigma \in \mathcal{E}_K} \mathcal{F}_{i, K, \sigma}^k = 0,$$

$$(17) \quad \mathcal{F}_{i, K, \sigma}^k = -\tau_{\sigma} u_{i, \sigma}^k D_{K, \sigma} p_i(u^k) - \eta^{\alpha} \tau_{\sigma} D_{K, \sigma} u_i^k,$$

where $\eta = \max\{\Delta x, \Delta t\}$, $0 < \alpha < 2$, and τ_{σ} is given by (12). The mobility $u_{i, \sigma}^k$ is defined for $\sigma \in \mathcal{E}$ by the upwind scheme

$$(18) \quad u_{i, \sigma}^k = \begin{cases} u_{i, K, \sigma}^k & \text{if } D_{K, \sigma} p_i(u^k) \geq 0, \\ u_{i, K}^k & \text{if } D_{K, \sigma} p_i(u^k) < 0. \end{cases}$$

The upwind approximation allows us to derive the discrete Shannon entropy inequality; see Remark 1. We may also use a logarithmic mean function; see Remark 2.

We have added some artificial diffusion in the numerical flux $\mathcal{F}_{i, K, \sigma}^k$, which vanishes in the limit $\eta \rightarrow 0$. The term is needed to show the convergence of the scheme. In particular, it provides an η -dependent bound for the full gradient, compensating the incomplete gradient estimate. Note that the artificial diffusion is *not* needed to prove the existence of discrete solutions, and we may set $\eta = 0$ in this case. Artificial diffusion/viscosity is used in numerical approximations of the Euler equations to stabilize the scheme; see, e.g., [23, (3.8)].

The numerical fluxes $\mathcal{F}_{i, K, \sigma}^k$ are consistent approximations of the exact fluxes through the edges, since $\mathcal{F}_{i, K, \sigma} + \mathcal{F}_{i, L, \sigma} = 0$ for all edges $\sigma = K|L$ and $\mathcal{F}_{i, K, \sigma} = 0$ for all $\mathcal{E}_{\text{ext}, K}$. The following discrete integration-by-parts formula holds for $v = (v_K) \in V_{\mathcal{T}}$:

$$(19) \quad \sum_{K \in \mathcal{T}} \sum_{\sigma \in \mathcal{E}_K} \mathcal{F}_{i, K, \sigma} v_K = - \sum_{\sigma \in \mathcal{E}_{\text{int}}} \mathcal{F}_{i, K, \sigma} D_{K, \sigma} v.$$

Notice that the terms $\mathcal{F}_{i, K, \sigma} D_{K, \sigma} v$ on the right-hand side only depend on σ , but not on the specific control volume K satisfying $\sigma \in \mathcal{E}_K$. Hence, to evaluate the sum on the right, we may pick for each σ any K with $\sigma \in \mathcal{E}_K$ as long as we keep K fixed.

Remark 1 (Discrete gradient-flow property for upwind scheme). The upwind approximation implies a discrete gradient-flow property. Indeed, we first observe that the concavity of the logarithm gives

$$b(\log a - \log b) \leq a - b \leq a(\log a - \log b) \quad \text{for all } a, b > 0.$$

Combined with definition (18) of $u_{i,\sigma}^k$, this leads for $u_{i,K}^k > 0$ and $u_{i,L}^k > 0$ to

$$(20) \quad u_{i,\sigma}^k (p_i(u_L^k) - p_i(u_K^k)) (\log u_{i,L}^k - \log u_{i,K}^k) \geq (p_i(u_L^k) - p_i(u_K^k)) (u_{i,L}^k - u_{i,K}^k)$$

and therefore, by discrete integration by parts (19),

$$(21) \quad \begin{aligned} \sum_{i=1}^n \sum_{K \in \mathcal{T}} \sum_{\sigma \in \mathcal{E}_K} \tau_\sigma \mathcal{F}_{i,K,\sigma}^k \log u_{i,K}^k &= - \sum_{i=1}^n \sum_{\sigma \in \mathcal{E}_{\text{int}}} \tau_\sigma u_{i,\sigma}^k \mathbf{D}_{K,\sigma} p_i(u^k) \mathbf{D}_{K,\sigma} \log u_i^k \\ &\quad - \eta^\alpha \sum_{i=1}^n \sum_{\sigma \in \mathcal{E}_{\text{int}}} \tau_\sigma \mathbf{D}_{K,\sigma} u_i^k \mathbf{D}_{K,\sigma} \log u_i^k \leq - \sum_{i=1}^n \sum_{\sigma \in \mathcal{E}_{\text{int}}} \tau_\sigma b_{ij} \mathbf{D}_{K,\sigma} u_j^k \mathbf{D}_{K,\sigma} u_i^k, \end{aligned}$$

where we used the monotonicity of the logarithm implying that $\mathbf{D}_{K,\sigma} u_i^k \mathbf{D}_{K,\sigma} \log u_i^k \geq 0$. The right-hand side of (21) is nonpositive due to the positive semidefiniteness of $B = (b_{ij})$. We deduce from this inequality the discrete entropy inequality (24). \square

Remark 2 (Discrete gradient flow property for logarithmic mean). We may define $u_{i,\sigma}^k$ via the logarithmic mean

$$(22) \quad u_{i,\sigma}^k = \begin{cases} \frac{u_{i,L}^k - u_{i,K}^k}{\log u_{i,L}^k - \log u_{i,K}^k} & \text{if } u_{i,K}^k \neq u_{i,L}^k \text{ and } u_{i,K}^k > 0, u_{i,L}^k > 0, \\ u_{i,K}^k & \text{if } u_{i,K}^k = u_{i,L}^k > 0, \\ 0 & \text{else.} \end{cases}$$

We remark that the artificial diffusion in the numerical flux (17) allows us to show that $u_{i,K}^k$ is positive for all $K \in \mathcal{T}$ (see Section 3.5) such that $u_{i,\sigma}^k$ (for $\sigma = K|L$) is always defined by one of the first two cases. Definition (22) also leads to a discrete gradient-flow property. Indeed, observing that $u_{i,\sigma}^k \mathbf{D}_{K,\sigma} \log u_i^k = \mathbf{D}_{K,\sigma} u_i^k$ and multiplying (17) by $\log u_{i,K}^k$ and summing over all $i = 1, \dots, n, K \in \mathcal{T}$, and $\sigma \in \mathcal{E}_K$, we see that (21) holds too. Notice that (20) becomes an equality in this case. \square

Finally, we observe that the mobility satisfies in both cases the following properties:

$$(23) \quad u_{i,\sigma}^k \leq \max\{u_{i,K}^k, u_{i,L}^k\}, \quad |u_{i,\sigma}^k - u_{i,K}^k| \leq |u_{i,K}^k - u_{i,L}^k| \quad \text{for } \sigma = K|L.$$

2.5. Main results. We impose the following hypotheses.

- (H1) Data: $\Omega \subset \mathbb{R}^d$ is a bounded polygonal (or polyhedral if $d \geq 3$) domain, $T > 0$, and $u^{\text{in}} \in L^2(\Omega; \mathbb{R}_{\geq}^n)$ such that $\|u^{\text{in}}\|_{L^1(\Omega)} > 0$. We set $\Omega_T = \Omega \times (0, T)$.
- (H2) Discretization: \mathcal{D} is an admissible discretization of Ω_T satisfying (13).
- (H3) Diffusion coefficients: $B = (b_{ij}) \in \mathbb{R}_{\geq}^{n \times n}$ is symmetric positive semidefinite with $\text{rank } B \in \{1, \dots, n\}$ and $b_{ii} > 0$ for $i = 1, \dots, n$.

Note that since B is positive semidefinite, its square root $B^{1/2}$ exists and $z^T B z = |B^{1/2} z|^2$ for $z \in \mathbb{R}^n$. Moreover, with $\lambda > 0$ being the smallest positive eigenvalue of B , we have $|B^{1/2} z|^2 \geq \lambda |z|^2$.

Theorem 3. *Let Hypotheses (H1)–(H3) hold, $k \in \mathbb{N}$, $\eta \geq 0$, and let $u^{k-1} \in V_{\mathcal{T}}^n$ be given. Then there exists a solution $u^k = (u_1^k, \dots, u_n^k) \in V_{\mathcal{T}}^n$ to scheme (15)–(17) satisfying $u_{i,K}^k > 0$ for $i = 1, \dots, n$, $K \in \mathcal{T}$. Inductively, let $u^j \in V_{\mathcal{T}}^n$, $j = 1, \dots, k$, be the solution to scheme (15)–(17) with u^{k-1} replaced by u^{j-1} . Then $\{u^j\}$ obey the discrete entropy inequalities*

$$(24) \quad H_S(u^k) + \sum_{j=1}^k \Delta t |B^{1/2} u^j|_{1,2,\mathcal{T}}^2 + 4\eta^\alpha \sum_{j=1}^k \Delta t \sum_{i=1}^n |(u_i^j)^{1/2}|_{1,2,\mathcal{T}}^2 \leq H_S(u^0),$$

$$(25) \quad H_R(u^k) + \sum_{j=1}^k \Delta t \sum_{i=1}^n \sum_{\sigma \in \mathcal{E}} \tau_\sigma u_{i,\sigma}^j |D_\sigma(Bu^j)_i|^2 \leq H_R(u^0).$$

Moreover, $H_R(u^k) \leq H_R(u^{k-1})$.

The existence of finite-volume solutions to (15)–(17) was shown in [34] by using the Rao entropy only, but the proof needs matrices B with full rank. We can avoid this condition since we exploit the estimates coming from the Shannon entropy. Theorem 3 is proved by adding a discrete version of the regularizing term $\varepsilon(-\Delta w_i + w_i)$, where $w_i = \log u_i$ are the entropic variables [25, 31, 36], and a topological degree argument, similar as in [34]. Uniform estimates from the Shannon entropy inequality (24) allow us to perform the de-regularizing limit $\varepsilon \rightarrow 0$. Observe that the theorem is valid for $\eta = 0$, i.e., no artificial diffusion is needed here.

Theorem 3 and the subsequent results also hold for domains $\Omega \subset \mathbb{R}^d$ with curved (Lipschitz) boundary. Indeed, one may triangulate Ω in such a way that the control volumes have a curved boundary [40], or one may cover Ω by additional cells and estimate the integral error; we refer to Remark 14 for details.

For the convergence result, we introduce a family $(\mathcal{D}_m)_{m \in \mathbb{N}}$ of admissible space-time discretizations of Ω_T indexed by the size $\eta_m = \max\{\Delta x_m, \Delta t_m\}$ of the mesh, where $\Delta x_m = \max_{K \in \mathcal{T}_m} \text{diam}(K)$ and Δt_m is the time step size of the mesh \mathcal{D}_m , satisfying $\eta_m \rightarrow 0$ as $m \rightarrow \infty$. We denote by \mathcal{T}_m the corresponding meshes of Ω and set $\nabla^m := \nabla^{\mathcal{D}_m}$.

Theorem 4 (Convergence of the scheme). *Let Hypotheses (H1)–(H3) hold, and let (\mathcal{D}_m) be a family of admissible meshes satisfying (13) uniformly in $m \in \mathbb{N}$. Let (u_m) be a sequence of finite-volume solutions to (15)–(17) with $\eta = \eta_m > 0$, constructed in Theorem 3. Then, up to a subsequence, $(u_m, \nabla^m \hat{u}_m)$ generates a Young measure μ which is a dissipative measure-valued solution to (4), (2) in the sense of Definition 1. Moreover, $t \mapsto H_R(u(t))$ is nonincreasing and μ acts trivially on the \hat{s} -component,*

$$(26) \quad \langle \mu, f(s, p) \rangle = \langle \mu, f(\hat{u} + P_L s, p) \rangle$$

for all $f \in C_0(\mathbb{R}_{\geq}^n \times (L^\perp)^d)$, where C_0 is the space of continuous functions vanishing at infinity and $L^\perp = \text{ran } B$.

Property (26) implies, if $\text{rank } B = n$, that u fulfills (4), (2) in the usual weak sense, since then $P_L = 0$.

The strategy of the proof of Theorem 4 is as follows. The estimates from the discrete entropy inequalities and a uniform bound for the discrete time derivative of u_m allow us to apply the compactness result of [26] to conclude the strong convergence of (a subsequence of) \hat{u}_m in $L^2(\Omega_T)$ as $m \rightarrow \infty$. Moreover, (u_m) and $\nabla^m(B\hat{u}_m)$ are weakly converging in $L^2(\Omega_T)$. Clearly, these convergences are too weak to conclude the convergence of the non-linear flux (17). However, the sequence $(u_m, \nabla^m \hat{u}_m)$ generates a parametrized measure μ [42, Chap. 6] such that $\langle \mu, s_i(Bp)_i \rangle$ is the distributional limit of $u_{m,i,\sigma} \nabla^m(B\hat{u}_m)_i$. Moreover, because of the strong convergence of (\hat{u}_m) , we can separate this part, leading to (26).

Remark 5 (Consistency of the definition). The definition of dissipative measure-valued solutions is consistent with the definition of weak solutions. Indeed, any weak solution u to (4), (2) satisfying the regularity statements of Definition 1 and the Shannon and Rao entropy inequalities gives rise to a dissipative measure-valued solution μ via $\mu_{x,t} = \delta_{u(x,t)} \otimes \delta_{\nabla \hat{u}(x,t)}$. On the other hand, if a dissipative measure-valued solution μ is trivial in the sense that $\mu_{x,t} = \delta_{v(x,t)} \otimes \delta_{z(x,t)}$ for certain functions v and z , then $v = \langle \mu, s \rangle = u$ and $z = \langle \mu, p \rangle = y = \nabla \hat{u}$. We infer that

$$\langle \mu, s_i(Bp)_i \rangle = u_i(B\nabla \hat{u})_i.$$

In this case, equation (9) reduces to the standard weak formulation of (4) for the density u and the entropy inequalities (10) and (11) take the usual form of entropy inequalities for weak solutions. More generally, the conclusion $\langle \mu, s_i(Bp)_i \rangle = u_i(B\nabla \hat{u})_i$ already holds if for instance μ is only atomic in the density component, i.e. $\mu_{x,t} = \delta_{v(x,t)} \otimes \nu_{x,t}$, where ν denotes the parametrized measure generated by $(\nabla^m \hat{u}_m)_m$. \square

Remark 6 (Full-rank approximation). Let $\Omega \subset \mathbb{R}^d$ be a bounded Lipschitz domain. An alternative to the finite-volume approach is to consider a suitable full-rank symmetric positive definite regularization $(B_\rho) \in \mathbb{R}^{n \times n}$ of B with $\lim_{\rho \rightarrow 0} B_\rho = B$, and to approximate (4) by

$$(27) \quad \partial_t u_i = \text{div}(u_i \nabla(B_\rho u)_i), \quad i = 1, \dots, n.$$

After an appropriate additional regularization, it is possible to apply the entropy method of [31, Sec. 4.4] (using the Rao entropy structure) and to establish the existence of a nonnegative weak solution to (27), (2) that satisfies both the Rao and Shannon entropy inequalities with B replaced by B_ρ . The dissipative measure-valued solution to (4), (2) is then obtained in the limit $\rho \rightarrow 0$.

The statement of Theorem 4 is rather weak, since the Young measure may not be unique. However, we can prove a weak–strong uniqueness result. According to Remark 14, we can assume in the following that Ω is a general bounded domain with Lipschitz boundary.

Theorem 7 (Weak–strong uniqueness). *Let $\Omega \subset \mathbb{R}^d$ be a bounded Lipschitz domain. Let $v \in C^1(\bar{\Omega} \times [0, T]; \mathbb{R}_{\geq}^n)$ be a positive solution to (4), (2) (in the weak sense) with initial*

datum $v(0) = u^{\text{in}} > 0$, and let μ be a dissipative measure-valued solution to (4), (2). Then

$$\mu_{x,t} = \delta_{v(x,t)} \otimes \delta_{\nabla \hat{v}(x,t)} \quad \text{for a.e. } (x,t) \in \Omega \times (0,T).$$

The assertion is deduced from a stability estimate based on the Bregman distance $f(u|v) := f(u) - f(v) - f'(v) \cdot (u-v)$ associated with the convex function $f := h_S + h_R$, which has to be adapted to the measure-valued framework. Loosely speaking, we consider the sum $H_S(u|v) + H_R(u|v)$, where

$$H_k(u|v) = \sum_{i=1}^n \int_{\Omega} (h_k(u) - h_k(v) - h'_k(v) \cdot (u-v)) dx, \quad k = S, R,$$

and compute its time derivative along solutions to (4). Certain error terms arising in this computation need to be estimated from above by $C \int_{\Omega} f(u|v) dx$. For this step and in the absence of $L^\infty(\Omega)$ -bounds on the densities u_i , we take advantage of the better coercivity properties at infinity of the Rao entropy.

As a consequence of Theorem 7, the finite-volume solution converges *strongly* to the classical solution if the latter exists.

Corollary 8. *Let $u \in C^1(\bar{\Omega} \times [0, T]; \mathbb{R}_{\geq}^n)$ be a positive solution to (4), (2). Let (u_m) be a sequence of finite-volume solutions to (15), (17) with $\eta = \eta_m > 0$. Then, as $m \rightarrow \infty$,*

$$(u_m, \nabla^m \hat{u}_m) \rightarrow (u, \nabla \hat{u}) \quad \text{strongly in } L^p(\Omega_T)$$

for all $p \in [1, 2)$ and all $T > 0$.

Indeed, the weak-strong uniqueness implies that the Young measure generated by $(u_m, \nabla^m \hat{u}_m)$ coincides at each point (x, t) with the Dirac measure concentrated at the smooth solution. Since $|(u_m, \nabla^m \hat{u}_m)|^p \subset L^1(\Omega_T)$ is equi-integrable for every $p \in [1, 2)$, the assertion in Corollary 8 thus follows from classical Young measure theory (cf. e.g. [42, Theorem 6.12]).

It is shown in [17, Theorem 2.6] for $\Omega = \mathbb{T}^d$ (with periodic boundary conditions) that problem (4), (2) possesses a positive classical solution on a short time interval if the initial data are positive and smooth. The main results in the present paper should equally be valid in the periodic setting.

If B has a non-trivial kernel, steady states to (4), (2) are not necessarily constant in space and for any fixed mass vector $\mathbf{m} \in (0, \infty)^n$, there exist infinitely many steady states. Given such \mathbf{m} , we define the space of steady states as

$$\mathfrak{S}_{\mathbf{m}} = \left\{ v \in L^2(\Omega; \mathbb{R}_{\geq}^n) : \int_{\Omega} v dx = \mathbf{m} \text{ and } \nabla(Bv) = 0 \text{ in } \Omega \right\}.$$

Theorem 9 (Long-time behavior). *Let μ be a dissipative measure-valued solution to (4), (2). Let $u = \langle \mu, s \rangle$ and set $\mathbf{m} := \int_{\Omega} u^0 dx$. Then $\mathfrak{S}_{\mathbf{m}} \subset L^\infty(\Omega; \mathbb{R}_{\geq}^n)$ and there exists $u^* \in \mathfrak{S}_{\mathbf{m}}$ such that, as $t \rightarrow \infty$,*

$$\hat{u}(t) \rightarrow \hat{u}^* \quad \text{strongly in } L^2(\Omega; \mathbb{R}_{\geq}^n),$$

where $\hat{u}^* = P_{L^\perp} u^*$. We recall that P_{L^\perp} is the projection onto $L^\perp = \text{ran } B$.

For the proof of Theorem 9, we argue as follows. The fact that $\int_0^\infty \|\nabla(B^{1/2}\widehat{u})\|_{L^2(\Omega)}^2 dt$ is finite implies the existence of a sequence $t_k \rightarrow \infty$ such that $k \mapsto (B^{1/2}u)(t_k)$ converges strongly in $L^2(\Omega)$ to $B^{1/2}u^*$ as $k \rightarrow \infty$, where $u^* \in \mathfrak{G}_m$. The monotonicity of $k \mapsto H_R(u(t_k)|u^*)$ then shows that $B^{1/2}\widehat{u}(t)$ converges and consequently, $\widehat{u}(t)$ converges to \widehat{u}^* for all sequences $t \rightarrow \infty$. Such reasoning is classical in degenerate cases, where entropy-entropy dissipation estimates are not available; see for instance [7, 29]. A new aspect in the present situation is the combination of the two functionals H_R and H_S .

3. DISCRETE PROBLEM

In this section, we prove Theorem 3. The existence proof uses a discrete analog of the entropy method for cross-diffusion systems [31]. We first introduce a regularized numerical scheme involving an approximation parameter $\varepsilon > 0$, prove the existence of a solution to this scheme and suitable estimates coming from the Shannon entropy inequality, and apply a topological degree argument. The uniform estimates allow us to perform the limit $\varepsilon \rightarrow 0$.

3.1. Definition and continuity of the fixed-point operator. Let $u^{k-1} \in V_{\mathcal{T}}^n$ be given and let $R > 0$, $\delta > 0$. We set

$$Z_R = \{w = (w_1, \dots, w_n) \in V_{\mathcal{T}}^n : \|w\|_{1,2,\mathcal{T}} < R \text{ for } i = 1, \dots, n\}$$

and define the mapping $F : Z_R \rightarrow \mathbb{R}^{\theta n}$ by $F(w) = w^\varepsilon$, where $\theta = \#\mathcal{T}$ and $w^\varepsilon = (w_1^\varepsilon, \dots, w_n^\varepsilon)$ is the solution to the linear regularized problem

$$(28) \quad \varepsilon \left(- \sum_{\sigma \in \mathcal{E}_K} \tau_\sigma D_{K,\sigma} w_i^\varepsilon + \mathbf{m}(K) w_{i,K}^\varepsilon \right) = - \left(\frac{\mathbf{m}(K)}{\Delta t} (u_{i,K} - u_{i,K}^{k-1}) + \sum_{\sigma \in \mathcal{E}_K} \mathcal{F}_{i,K,\sigma} \right),$$

where $u_{i,K} := \exp(w_{i,K})$ and $\mathcal{F}_{i,K,\sigma}$ is defined as in (17) with $u_{i,K}^k$ replaced by $u_{i,K}$.

To show that F is well defined, we write (28) as

$$(29) \quad \begin{aligned} Mw^\varepsilon &= v, \quad \text{where } v = (v_{i,K})_{i=1,\dots,n, K \in \mathcal{T}}, \\ v_{i,K} &= \frac{\mathbf{m}(K)}{\Delta t} (u_{i,K} - u_{i,K}^{k-1}) + \sum_{\sigma \in \mathcal{E}_K} \mathcal{F}_{i,K,\sigma}, \end{aligned}$$

and $M = \text{diag}(M', \dots, M') \in \mathbb{R}^{\theta n \times \theta n}$ is a block diagonal matrix with $M' \in \mathbb{R}^{\theta \times \theta}$, which has the entries

$$M'_{K,K} = -\varepsilon \mathbf{m}(K) - \varepsilon \sum_{\sigma \in \mathcal{E}_K} \tau_\sigma, \quad M'_{K,L} = \begin{cases} \varepsilon \tau_\sigma & \text{if } K \cap L \neq \emptyset, \sigma = K|L, \\ 0 & \text{if } K \cap L = \emptyset. \end{cases}$$

Therefore, the system $Mw^\varepsilon = v$ can be decomposed into the independent subsystems $M'w_i^\varepsilon = v_i$ for $i = 1, \dots, n$. Since M' is strictly diagonally dominant, these subsystems possess a unique solution w_i^ε . Then $w^\varepsilon = (w_1^\varepsilon, \dots, w_n^\varepsilon)$ is the unique solution to (29). Thus, the mapping F is well defined.

Next, we prove that F is continuous. We multiply (28) for some fixed $i \in \{1, \dots, n\}$ by $w_{i,K}^\varepsilon$ and sum over all $i = 1, \dots, n$ and $K \in \mathcal{T}$:

$$(30) \quad \begin{aligned} & -\varepsilon \sum_{i=1}^n \sum_{K \in \mathcal{T}} \sum_{\sigma \in \mathcal{E}_K} \tau_\sigma (\mathbb{D}_{K,\sigma} w_i^\varepsilon) w_{i,K}^\varepsilon + \varepsilon \sum_{i=1}^n \sum_{K \in \mathcal{T}} \mathbf{m}(K) (w_{i,K}^\varepsilon)^2 \\ & = - \sum_{i=1}^n \sum_{K \in \mathcal{T}} \frac{\mathbf{m}(K)}{\Delta t} (u_{i,K} - u_{i,K}^{k-1}) w_{i,K}^\varepsilon - \sum_{i=1}^n \sum_{K \in \mathcal{T}} \sum_{\sigma \in \mathcal{E}_K} \mathcal{F}_{i,K,\sigma} w_{i,K}^\varepsilon. \end{aligned}$$

Using discrete integration by parts analogous to (19), we can rewrite the left-hand side as

$$\begin{aligned} & -\varepsilon \sum_{i=1}^n \sum_{K \in \mathcal{T}} \sum_{\sigma \in \mathcal{E}_K} \tau_\sigma (\mathbb{D}_{K,\sigma} w_i^\varepsilon) w_{i,K}^\varepsilon + \varepsilon \sum_{i=1}^n \sum_{K \in \mathcal{T}} \mathbf{m}(K) (w_{i,K}^\varepsilon)^2 \\ & = \varepsilon \sum_{i=1}^n \sum_{\sigma \in \mathcal{E}_{\text{int}}} \tau_\sigma (\mathbb{D}_{K,\sigma} w_i^\varepsilon)^2 + \varepsilon \sum_{i=1}^n \sum_{K \in \mathcal{T}} \mathbf{m}(K) (w_{i,K}^\varepsilon)^2 = \varepsilon \sum_{i=1}^n \|w_i^\varepsilon\|_{1,2,\mathcal{T}}^2. \end{aligned}$$

We turn to the terms on the right-hand side of (30). By definition, we have $\|w_i\|_{1,2,\mathcal{T}} < R$ and consequently $\|w_i\|_{0,\infty,\mathcal{T}} \leq C(R, \mathcal{T})$ and $\|u_i\|_{1,2,\mathcal{T}} \leq C(R, \mathcal{T})$ (since the problem is finite-dimensional). This shows that

$$\begin{aligned} & - \sum_{i=1}^n \sum_{K \in \mathcal{T}} \frac{\mathbf{m}(K)}{\Delta t} (u_{i,K} - u_{i,K}^{k-1}) w_{i,K}^\varepsilon \leq \frac{1}{\Delta t} \sum_{i=1}^n \|u_i - u_i^{k-1}\|_{0,2,\mathcal{T}} \|w_i^\varepsilon\|_{0,2,\mathcal{T}} \\ & \leq C(R, \mathcal{T}, \Delta t) \sum_{i=1}^n \|w_i^\varepsilon\|_{1,2,\mathcal{T}}. \end{aligned}$$

Finally, using definition (17) of the flux and discrete integration by parts,

$$\begin{aligned} & - \sum_{i=1}^n \sum_{K \in \mathcal{T}} \sum_{\sigma \in \mathcal{E}_K} \mathcal{F}_{i,K,\sigma} w_{i,K}^\varepsilon = \sum_{i=1}^n \sum_{K \in \mathcal{T}} \sum_{\sigma \in \mathcal{E}_K} \tau_\sigma \left(\sum_{j=1}^n b_{ij} u_{i,\sigma} (\mathbb{D}_{K,\sigma} u_j) + \eta^\alpha \mathbb{D}_{K,\sigma} u_i \right) w_{i,K}^\varepsilon \\ & = - \sum_{i=1}^n \sum_{\sigma \in \mathcal{E}_{\text{int}}} \tau_\sigma \left(\sum_{j=1}^n b_{ij} u_{i,\sigma} (\mathbb{D}_{K,\sigma} u_j) (\mathbb{D}_{K,\sigma} w_i^\varepsilon) + \eta^\alpha (\mathbb{D}_{K,\sigma} u_i) (\mathbb{D}_{K,\sigma} w_i^\varepsilon) \right) \\ & \leq \max_{\sigma \in \mathcal{E}} \|u_{i,\sigma}\|_{0,\infty,\mathcal{T}} \sum_{i,j=1}^n b_{ij} |u_j|_{1,2,\mathcal{T}} |w_i^\varepsilon|_{1,2,\mathcal{T}} + \eta^\alpha \sum_{i=1}^n |u_i|_{1,2,\mathcal{T}} |w_i^\varepsilon|_{1,2,\mathcal{T}} \\ & \leq C(R, \mathcal{T}) \|w_i^\varepsilon\|_{1,2,\mathcal{T}}. \end{aligned}$$

For the last inequality, we used the fact that $u_{i,\sigma}$ depends on $u_{i,K}$ and $u_{i,L}$ for $\sigma = K|L$, and their discrete $L^\infty(\Omega)$ norms can be bounded by the discrete $L^\infty(\Omega)$ norm of w_i , which in turn can be estimated by $C(\mathcal{T}) \|w_i\|_{0,\infty,\mathcal{T}} \leq C(R, \mathcal{T})$.

Inserting these estimates into (30) and dividing by $\|w_i^\varepsilon\|_{1,2,\mathcal{T}}$ if $\|w_i^\varepsilon\|_{1,2,\mathcal{T}} > 0$, it follows that $\varepsilon \|w_i^\varepsilon\|_{1,2,\mathcal{T}} \leq C(R, \mathcal{T}, \Delta t)$. This bound allows us to verify the continuity of F . Indeed, let $w^\ell \rightarrow w$ as $\ell \rightarrow \infty$ and set $w^{\varepsilon,\ell} = F(w^\ell)$. Then $(w^{\varepsilon,\ell})_{\ell \in \mathbb{N}}$ is uniformly bounded in the discrete $H^1(\Omega)$ norm. Therefore, there exists a subsequence, which is not relabeled, such

that $w^{\varepsilon, \ell} \rightarrow w^\varepsilon$ as $\ell \rightarrow \infty$. Passing to the limit $\ell \rightarrow \infty$ in scheme (28), we see that w^ε is a solution to the scheme and $w^\varepsilon = F(w)$. Since the solution to the linear scheme (28) is unique, the entire sequence $(w^{\varepsilon, \ell})_{\ell \in \mathbb{N}}$ converges to w^ε , which shows the continuity of F .

3.2. Existence of a fixed point. We will now show that the map F admits a fixed point by using a topological degree argument. We prove that $\deg(I - F, Z_R, 0) = 1$, where \deg is the Brouwer topological degree [12, Chap. 1]. Since \deg is invariant by homotopy, it is sufficient to verify that any solution $(w^\varepsilon, \rho) \in \overline{Z}_R \times [0, 1]$ to the fixed-point equation $w^\varepsilon = \rho F(w^\varepsilon)$ satisfies $(w^\varepsilon, \rho) \notin \partial Z_R \times [0, 1]$ for sufficiently large values of $R > 0$. Let (w^ε, ρ) be a fixed point. The case $\rho = 0$ being clear, we assume that $\rho \neq 0$. Then w_i^ε solves

$$(31) \quad \varepsilon \left(- \sum_{\sigma \in \mathcal{E}_K} \tau_\sigma D_{K, \sigma} w_i^\varepsilon + \mathbf{m}(K) w_{i, K}^\varepsilon \right) = -\rho \left(\frac{\mathbf{m}(K)}{\Delta t} (u_{i, K}^\varepsilon - u_{i, K}^{k-1}) + \sum_{\sigma \in \mathcal{E}_K} \mathcal{F}_{i, K, \sigma}^\varepsilon \right)$$

for $i = 1, \dots, n$ and $K \in \mathcal{T}$, where $u_{i, K}^\varepsilon = \exp(w_{i, K}^\varepsilon)$ and $\mathcal{F}_{i, K, \sigma}^\varepsilon$ is defined as in (17) with $u_{i, K}^k$ replaced by $u_{i, K}^\varepsilon$. The following inequality is the key argument.

Lemma 10 (Discrete Shannon entropy inequality). *Let w^ε be a solution to (31) and $u_i^\varepsilon := \exp(w_i^\varepsilon)$. Then*

$$(32) \quad \begin{aligned} \rho H_S(u^\varepsilon) + \varepsilon \Delta t \sum_{i=1}^n \|w_i^\varepsilon\|_{1,2,\mathcal{T}}^2 + \rho \Delta t \sum_{i,j=1}^n \sum_{\sigma \in \mathcal{E}_{\text{int}}} \tau_\sigma b_{ij} D_{K, \sigma} u_i^\varepsilon D_{K, \sigma} u_j^\varepsilon \\ + 4\rho \eta^\alpha \Delta t \sum_{i=1}^n |(u_i^\varepsilon)^{1/2}|_{1,2,\mathcal{T}}^2 \leq \rho H_S(u^{k-1}). \end{aligned}$$

Proof. We multiply (31) by $\Delta t w_{i, K}^\varepsilon$, sum over $i = 1, \dots, n$ and $K \in \mathcal{T}$, and use discrete integration by parts (cf. (19)). Then $\varepsilon \Delta t \sum_{i=1}^n \|w_i^\varepsilon\|_{1,2,\mathcal{T}}^2 = I_1 + I_2 + I_3$, where

$$\begin{aligned} I_1 &= -\rho \sum_{i=1}^n \sum_{K \in \mathcal{T}} \mathbf{m}(K) (u_{i, K}^\varepsilon - u_{i, K}^{k-1}) w_{i, K}^\varepsilon, \\ I_2 &= -\rho \Delta t \sum_{i=1}^n \sum_{\sigma \in \mathcal{E}_{\text{int}}} \tau_\sigma u_{i, \sigma}^\varepsilon D_{K, \sigma} p_i(u^\varepsilon) D_{K, \sigma} w_{i, K}^\varepsilon, \\ I_3 &= -\rho \eta^\alpha \Delta t \sum_{i=1}^n \sum_{\sigma \in \mathcal{E}_{\text{int}}} \tau_\sigma D_{K, \sigma} u_i^\varepsilon D_{K, \sigma} w_{i, K}^\varepsilon. \end{aligned}$$

The definition $u_{i, K}^\varepsilon = \exp(w_{i, K}^\varepsilon)$ and the convexity of the Shannon entropy imply that

$$I_1 = -\rho \sum_{i=1}^n \sum_{K \in \mathcal{T}} \mathbf{m}(K) (u_{i, K}^\varepsilon - u_{i, K}^{k-1}) \log u_{i, K}^\varepsilon \leq -\rho (H_S(u^\varepsilon) - H_S(u^{k-1})).$$

For I_2 , we rely on inequality (20):

$$I_2 = -\rho \Delta t \sum_{i=1}^n \sum_{\sigma=K|L \in \mathcal{E}_{\text{int}}} \tau_\sigma u_{i, \sigma}^\varepsilon (p_i(u_L^\varepsilon) - p_i(u_K^\varepsilon)) (\log u_{i, L}^\varepsilon - \log u_{i, K}^\varepsilon)$$

$$\begin{aligned}
&\leq -\rho\Delta t \sum_{i,j=1}^n \sum_{\sigma=K|L\in\mathcal{E}_{\text{int}}} \tau_{\sigma} b_{ij}(u_{j,L}^{\varepsilon} - u_{j,K}^{\varepsilon})(u_{i,L}^{\varepsilon} - u_{i,K}^{\varepsilon}) \\
&= -\rho\Delta t \sum_{i,j=1}^n \sum_{\sigma=K|L\in\mathcal{E}_{\text{int}}} \tau_{\sigma} b_{ij} D_{K,\sigma} u_i^{\varepsilon} D_{K,\sigma} u_j^{\varepsilon}.
\end{aligned}$$

Finally, using the elementary inequality $(a - b)(\log a - \log b) \geq 4(\sqrt{a} - \sqrt{b})^2$,

$$\begin{aligned}
I_3 &= -\rho\eta^{\alpha}\Delta t \sum_{i=1}^n \sum_{\sigma=K|L\in\mathcal{E}_{\text{int}}} \tau_{\sigma}(u_{i,L}^{\varepsilon} - u_{i,K}^{\varepsilon})(\log u_{i,L}^{\varepsilon} - \log u_{i,K}^{\varepsilon}) \\
&\leq -4\rho\eta^{\alpha}\Delta t \sum_{i=1}^n \sum_{\sigma=K|L\in\mathcal{E}_{\text{int}}} \tau_{\sigma} \left((u_{i,L}^{\varepsilon})^{1/2} - (u_{i,K}^{\varepsilon})^{1/2} \right)^2 = -4\rho\eta^{\alpha}\Delta t \sum_{i=1}^n |(u_i^{\varepsilon})^{1/2}|_{1,2,\mathcal{T}}^2.
\end{aligned}$$

Combining these estimates finishes the proof of Lemma 10. \square

We now complete the topological degree argument. Lemma 10 implies that

$$\varepsilon\Delta t \sum_{i=1}^n \|w_i^{\varepsilon}\|_{1,2,\mathcal{T}}^2 \leq \rho H_S(u^{k-1}) \leq H_S(u^{k-1}).$$

With the choice $R := (\varepsilon\Delta t)^{-1/2} H_S(u^{k-1})^{1/2} + 1$ we find that $w^{\varepsilon} \notin \partial Z_R$ and $\deg(I - F, Z_R, 0) = 1$. We conclude that F possesses a fixed point.

3.3. Limit $\varepsilon \rightarrow 0$. By Lemma 10, there exists $C > 0$, independent of ε , such that

$$C \sum_{i=1}^n \sum_{K\in\mathcal{T}} \mathbf{m}(K)(u_{i,K}^{\varepsilon} - 1) \leq H_S(u^{\varepsilon}) \leq H_S(u^{k-1}).$$

This gives a uniform discrete $L^1(\Omega)$ bound for u_i^{ε} . There exists a subsequence (not relabeled) such that $u_{i,K}^{\varepsilon} \rightarrow u_{i,K}$ as $\varepsilon \rightarrow 0$ for all $i = 1, \dots, n$ and $K \in \mathcal{T}$. Moreover, the discrete $H^1(\Omega)$ bound for $\sqrt{\varepsilon}w_i^{\varepsilon}$ implies that $\varepsilon w_{i,K}^{\varepsilon} \rightarrow 0$ for $i = 1, \dots, n$ and $K \in \mathcal{T}$. Then the limit $\varepsilon \rightarrow 0$ in (31) yields the existence of a solution $u^k := (u_{i,K})_{i=1,\dots,n,K\in\mathcal{T}}$ to (16). Observing that

$$\begin{aligned}
\sum_{i,j=1}^n \sum_{\sigma\in\mathcal{E}} \tau_{\sigma} b_{ij} D_{K,\sigma} u_i^{\varepsilon} D_{K,\sigma} u_j^{\varepsilon} &= \sum_{\sigma\in\mathcal{E}} \tau_{\sigma} (D_{K,\sigma} u^{\varepsilon})^T B(D_{K,\sigma} u^{\varepsilon}) \\
&\geq \lambda \sum_{\sigma\in\mathcal{E}} \tau_{\sigma} |B^{1/2} D_{K,\sigma} u^{\varepsilon}|^2 = \lambda |B^{1/2} u^{\varepsilon}|_{1,2,\mathcal{T}}^2,
\end{aligned}$$

the same limit in the regularized entropy inequality (32) directly leads to the discrete entropy inequality (24).

3.4. Discrete Rao entropy inequality. To verify (25), we multiply (16) by $\Delta t p_i(u_K^k)$, sum over $i = 1, \dots, n$ and $K \in \mathcal{T}$, and use discrete integration by parts:

$$\begin{aligned}
& \sum_{i=1}^n \sum_{K \in \mathcal{T}} \mathbf{m}(K) (u_{i,K}^k - u_{i,K}^{k-1}) p_i(u_K^k) \\
&= \Delta t \sum_{i=1}^n \sum_{K \in \mathcal{T}} \sum_{\sigma \in \mathcal{E}_K} \tau_\sigma (u_{i,\sigma}^k D_{K,\sigma} p_i(u^k) + \eta^\alpha D_{K,\sigma} u_i^k) p_i(u_K^k) \\
&= -\Delta t \sum_{i=1}^n \sum_{\sigma \in \mathcal{E}_{\text{int}}} \tau_\sigma u_{i,\sigma}^k (D_\sigma p_i(u^k))^2 - \eta^\alpha \Delta t \sum_{i,j=1}^n \sum_{\sigma \in \mathcal{E}_{\text{int}}} \tau_\sigma b_{ij} D_{K,\sigma} u_i^k D_{K,\sigma} u_j^k \\
&= -\Delta t \sum_{i=1}^n \sum_{\sigma \in \mathcal{E}_{\text{int}}} \tau_\sigma u_{i,\sigma}^k (D_\sigma p_i(u^k))^2 - \eta^\alpha \Delta t \sum_{i=1}^n |(B^{1/2} u^k)_i|_{1,2,\mathcal{T}}^2.
\end{aligned}$$

By the definition of $p_i(u^k)$ and the symmetry and positive semidefiniteness of B , the left-hand side becomes

$$\begin{aligned}
& \sum_{i=1}^n \sum_{K \in \mathcal{T}} \mathbf{m}(K) (u_{i,K}^k - u_{i,K}^{k-1}) p_i(u_K^k) = \sum_{i,j=1}^n \sum_{K \in \mathcal{T}} \mathbf{m}(K) b_{ij} (u_{i,K}^k - u_{i,K}^{k-1}) u_{j,K}^k \\
&= \frac{1}{2} \sum_{i,j=1}^n \sum_{K \in \mathcal{T}} \mathbf{m}(K) b_{ij} (u_{i,K}^k u_{j,K}^k - u_{i,K}^{k-1} u_{j,K}^{k-1} + (u_{i,K}^k - u_{i,K}^{k-1})(u_{j,K}^k - u_{j,K}^{k-1})) \\
&\geq H_R(u^k) - H_R(u^{k-1}).
\end{aligned}$$

We infer the monotonicity of $k \mapsto H_R(u^k)$. After summation over $k = 1, \dots, j$ and a renaming of the indices k and j , this shows (25) and thus completes the proof of Theorem 3.

3.5. Positivity. Thanks to the artificial diffusion, the discrete solution $u_{i,K}^k$ is positive for $i = 1, \dots, n$ and $K \in \mathcal{T}$. Indeed, let $i \in \{1, \dots, n\}$ be fixed and assume that there exists $K \in \mathcal{T}$ such that $u_{i,K}^k = 0$. We infer from I_3 in Section 3.2 that

$$\eta^\alpha (u_{i,L}^\varepsilon - u_{i,K}^\varepsilon) (\log u_{i,L}^\varepsilon - \log u_{i,K}^\varepsilon) \leq C(\Delta t, u^0),$$

where $L \in K$ is a neighboring cell of K . If $u_{i,L}^k > 0$, the limit $\varepsilon \rightarrow 0$ in the previous estimate leads to a contradiction since $\log u_{i,K}^\varepsilon$ diverges. Therefore, $u_{i,L}^k = 0$. Let $L' \in \mathcal{T}$ be a neighboring cell of L . Arguing in a similar way as before, it follows that $u_{i,L'}^k = 0$. Repeating this argument for all cells in \mathcal{T} , we find that $u_{i,K}^k = 0$ for all $K \in \mathcal{T}$. This implies that $\sum_{K \in \mathcal{T}} \mathbf{m}(K) u_{i,K}^k = 0$ and, by mass conservation, $\sum_{K \in \mathcal{T}} \mathbf{m}(K) u_{i,K}^0 = 0$, which contradicts the positivity of the $L^1(\Omega)$ norm of u^0 in Hypothesis (H1).

4. CONVERGENCE

In this section, we prove Theorem 4, that is, we show the asserted convergence of the numerical scheme. Uniform estimates are derived from the entropy inequalities (24) and (25). Lemma 16 in the appendix shows that $|\hat{u}^k| \leq \lambda^{-1} |B^{1/2} u^k|$, where we recall that $\hat{u}^k = P_{L^\perp} u^k$.

Thus, we obtain a uniform estimate for \widehat{u}^k in the seminorm $|\cdot|_{1,2,\mathcal{T}}$. Moreover, since $b_{ii} > 0$ and $b_{ij} \geq 0$ for all i, j (cf. Hypothesis (H3)), estimate (25) provides a uniform bound for u^k in the discrete $L^2(\Omega)$ norm. Hence, there exists a constant $C > 0$ which is independent of $\eta = \max\{\Delta x, \Delta t\}$ such that

$$(33) \quad \sum_{k=1}^{N_T} \Delta t (\|\widehat{u}^k\|_{1,2,\mathcal{T}}^2 + \|B^{1/2}u^k\|_{1,2,\mathcal{T}}^2) + \eta^\alpha \sum_{k=1}^{N_T} \Delta t |(u^k)^{1/2}|_{1,2,\mathcal{T}}^2 \leq C,$$

$$(34) \quad \max_{k=1,\dots,N_T} \|u^k\|_{0,2,\mathcal{T}} + \sum_{j=1}^k \Delta t \sum_{i=1}^n \sum_{\sigma \in \mathcal{E}_{\text{int}}} \tau_\sigma u_{i,\sigma}^j |D_\sigma(Bu^j)_i|^2 \leq C.$$

4.1. Compactness properties. We first prove a full gradient bound with a negative power of η on the right-hand side.

Lemma 11. *There exists $C = C(\zeta) > 0$ independent of η such that*

$$\sum_{k=1}^N \Delta t |u_i^k|_{1,4/3,\mathcal{T}}^2 \leq C\eta^{-\alpha}, \quad \sum_{k=1}^N \Delta t |u_i^k|_{1,1,\mathcal{T}}^2 \leq C\eta^{-\alpha}.$$

Proof. By the mesh regularity (13) and property (14),

$$(35) \quad \sum_{\sigma \in \mathcal{E}_K} \frac{\widetilde{\mathbf{m}}(\sigma) d_\sigma}{\mathbf{m}(K)} \leq \sum_{\sigma \in \mathcal{E}_K} \frac{\widetilde{\mathbf{m}}(\sigma) d(x_K, \sigma)}{\zeta \mathbf{m}(K)} \leq \frac{d}{\zeta}.$$

This yields, using Hölder's inequality and the $L^2(\Omega)$ bound (34) for u_i^k ,

$$\begin{aligned} |u_i^k|_{1,4/3,\mathcal{T}}^{4/3} &= \sum_{\sigma \in \mathcal{E}_{\text{int}}} \widetilde{\mathbf{m}}(\sigma) d_\sigma \left| \frac{u_{i,L}^k - u_{i,K}^k}{d_\sigma} \right|^{4/3} \\ &= \sum_{\sigma \in \mathcal{E}_{\text{int}}} \widetilde{\mathbf{m}}(\sigma) d_\sigma^{-1/3} |(u_{i,L}^k)^{1/2} - (u_{i,K}^k)^{1/2}|^{4/3} |(u_{i,L}^k)^{1/2} + (u_{i,K}^k)^{1/2}|^{4/3} \\ &\leq \left(\sum_{\sigma \in \mathcal{E}_{\text{int}}} \widetilde{\mathbf{m}}(\sigma) d_\sigma^{-1} ((u_{i,L}^k)^{1/2} - (u_{i,K}^k)^{1/2})^2 \right)^{2/3} \\ &\quad \times \left(\sum_{\sigma \in \mathcal{E}_{\text{int}}} \widetilde{\mathbf{m}}(\sigma) d_\sigma ((u_{i,L}^k)^{1/2} + (u_{i,K}^k)^{1/2})^4 \right)^{1/3} \\ &\leq C |(u_i^k)^{1/2}|_{1,2,\mathcal{T}}^{4/3} \left(\sum_{K \in \mathcal{T}} \mathbf{m}(K) (u_{i,K}^k)^2 \sum_{\sigma \in \mathcal{E}_K} \frac{\widetilde{\mathbf{m}}(\sigma) d_\sigma}{\mathbf{m}(K)} \right)^{1/3} \\ &\leq C(\zeta) |(u_i^k)^{1/2}|_{1,2,\mathcal{T}}^{4/3} \|u_i^k\|_{0,2,\mathcal{T}}^{2/3}. \end{aligned}$$

Taking the exponent $3/2$, multiplying by Δt , and summing over $k = 1, \dots, N$ proves the first inequality. The second inequality follows along the same lines (or by Hölder's inequality). \square

Lemma 12. *There exists $C = C(\zeta) > 0$ independent of η such that*

$$\sum_{k=1}^N \Delta t \|u_{i,\sigma}^k (B\nabla^{\mathcal{D}} \hat{u}^k)_i\|_{0,4/3,\mathcal{T}}^2 \leq C.$$

Proof. We infer from the definition of the discrete gradient and Hölder's inequality that

$$\begin{aligned} (36) \quad & \|u_{i,\sigma}^k (B\nabla^{\mathcal{D}} \hat{u}^k)_i\|_{0,4/3,\mathcal{T}}^{4/3} = \sum_{K \in \mathcal{T}} \sum_{\sigma \in \mathcal{E}_{\text{int},K}} \mathbf{m}(T_{K,\sigma}) (u_{i,\sigma}^k)^{4/3} \left| \frac{\tilde{\mathbf{m}}(\sigma)}{\mathbf{m}(T_{K,\sigma})} \mathbf{D}_{K,\sigma}(B\hat{u}^k)_i \right|^{4/3} \\ & = \sum_{K \in \mathcal{T}} \sum_{\sigma \in \mathcal{E}_{\text{int},K}} \mathbf{m}(T_{K,\sigma})^{1/3} (u_{i,\sigma}^k)^{2/3} \frac{\tilde{\mathbf{m}}(\sigma)^{4/3}}{\mathbf{m}(T_{K,\sigma})^{2/3}} |(u_{i,\sigma}^k)^{1/2} \mathbf{D}_{K,\sigma}(B\hat{u}^k)_i|^{4/3} \\ & \leq \left(\sum_{K \in \mathcal{T}} \sum_{\sigma \in \mathcal{E}_{\text{int},K}} \mathbf{m}(T_{K,\sigma}) (u_{i,\sigma}^k)^2 \right)^{1/3} \left(\sum_{K \in \mathcal{T}} \sum_{\sigma \in \mathcal{E}_{\text{int},K}} \frac{\tilde{\mathbf{m}}(\sigma)^2}{\mathbf{m}(T_{K,\sigma})} u_{i,\sigma}^k |\mathbf{D}_{K,\sigma}(B\hat{u}^k)_i|^2 \right)^{2/3}. \end{aligned}$$

Because of $\mathbf{m}(T_{K,\sigma}) = \tilde{\mathbf{m}}(\sigma) d_\sigma / d$ for $\sigma \in \mathcal{E}_{\text{int},K}$, mesh regularity (13), and property (14), we find for the first factor that

$$\begin{aligned} (37) \quad & \sum_{K \in \mathcal{T}} \sum_{\sigma \in \mathcal{E}_{\text{int},K}} \mathbf{m}(T_{K,\sigma}) (u_{i,\sigma}^k)^2 \leq C(\zeta) \sum_{K \in \mathcal{T}} \left(\sum_{\sigma \in \mathcal{E}_{\text{int},K}} \tilde{\mathbf{m}}(\sigma) d(x_K, \sigma) \right) (u_{i,K}^k)^2 \\ & \leq C(\zeta) \sum_{K \in \mathcal{T}} \mathbf{m}(K) (u_{i,K}^k)^2 = C(\zeta) \|u_i\|_{0,2,\mathcal{T}}^2, \end{aligned}$$

where we also used (23). The second factor on the right-hand side of (36) becomes

$$\sum_{K \in \mathcal{T}} \sum_{\sigma \in \mathcal{E}_{\text{int},K}} \frac{\tilde{\mathbf{m}}(\sigma)^2}{\mathbf{m}(T_{K,\sigma})} u_{i,\sigma}^k |\mathbf{D}_{K,\sigma}(B\hat{u}^k)_i|^2 = d \sum_{K \in \mathcal{T}} \sum_{\sigma \in \mathcal{E}_{\text{int},K}} \tau_\sigma u_{i,\sigma}^k |\mathbf{D}_{K,\sigma}(B\hat{u}^k)_i|^2.$$

We take (36) to the power 3/2, multiply by Δt , and sum over $k = 1, \dots, N$:

$$\sum_{k=1}^N \Delta t \|u_{i,\sigma}^k (B\nabla^{\mathcal{D}} \hat{u}^k)_i\|_{0,4/3,\mathcal{T}}^2 \leq C \max_{k=1,\dots,N} \|u_i^k\|_{0,2,\mathcal{T}}^2 \sum_{k=1}^N \Delta t \sum_{\sigma \in \mathcal{E}_{\text{int}}} \tau_\sigma u_{i,\sigma}^k |\mathbf{D}_\sigma(B\hat{u}^k)_i|^2 \leq C,$$

where the uniform bound follows from (34). \square

For the compactness argument, we need an estimate for the discrete time derivative, which is defined by

$$\partial_t^{\Delta t} v^k = \frac{v^k - v^{k-1}}{\Delta t} \quad \text{for } v \in V_{\mathcal{T},\Delta t}, \quad k = 1, \dots, N.$$

Lemma 13 (Discrete time derivative). *There exists a constant $C = C(\zeta) > 0$ independent of η such that*

$$\sum_{k=1}^N \Delta t \|\partial_t^{\Delta t} u^k\|_{-1,4,\mathcal{T}}^2 \leq C.$$

Proof. Let $\phi \in V_{\mathcal{T}}$ be such that $\|\phi\|_{1,4,\mathcal{T}} = 1$. We multiply (16) by ϕ_K , sum over $K \in \mathcal{T}$, apply discrete integration by parts, and use Hölder's inequality:

$$\begin{aligned} & \left| \sum_{K \in \mathcal{T}} \frac{\mathbf{m}(K)}{\Delta t} (u_{i,K}^k - u_{i,K}^{k-1}) \phi_K \right| \\ &= \left| - \sum_{\sigma \in \mathcal{E}_{\text{int}}} \tau_{\sigma} u_{i,\sigma}^k D_{K,\sigma} p_i(u^k) D_{K,\sigma} \phi - \eta^{\alpha} \sum_{\sigma \in \mathcal{E}_{\text{int}}} \tau_{\sigma} D_{K,\sigma} u_{i,\sigma}^k D_{K,\sigma} \phi \right| \\ &\leq C \|u_{i,\sigma}^k (B \nabla^{\mathcal{D}} \hat{u}^k)_i\|_{0,4/3,\mathcal{T}} \|\phi\|_{1,4,\mathcal{T}} + \eta^{\alpha} \|u_{i,\sigma}^k\|_{1,4/3,\mathcal{T}} \|\phi\|_{1,4,\mathcal{T}}. \end{aligned}$$

Then we infer from Lemmas 11 and 12 that

$$\sum_{k=1}^N \Delta t \left\| \frac{u_i^k - u_i^{k-1}}{\Delta t} \right\|_{-1,4,\mathcal{T}}^2 \leq C(\zeta) + C(\zeta) \eta^{\alpha},$$

which concludes the proof. \square

The solution $u^k \in V_{\mathcal{T}}$ to (16) refers to a fixed mesh. Let $(\mathcal{D}_m)_{m \in \mathbb{N}}$ be a sequence of meshes satisfying (13) such that the mesh size $\eta_m = \max\{\Delta x_m, \Delta t_m\}$ converges to zero as $m \rightarrow \infty$ and set $N_m = T/\Delta t_m$. Let $u_m = (u_{m,1}, \dots, u_{m,n})$ be defined as the piecewise constant function $u_m(x, t) = u_K^k$ for $(x, t) \in K \times [t_{k-1}, t_k)$, where u^k is a solution to (16) on the mesh \mathcal{D}_m , $K \in \mathcal{T}$, and $k = 1, \dots, N$, and set $u_m^0 = (u_{m,i}^0)_{i=1}^n$, where $u_{m,i}^0(x) := u_{i,K}^0(x)$ for $x \in K$. Notice that $u_m^0 \rightarrow u^{\text{in}}$ in $L^2(\Omega)$ as $m \rightarrow \infty$. Furthermore, we introduce the function $u_{m,\sigma} := (u_{m,i,\sigma})_{i=1}^n$ defined by $u_{m,i,\sigma}(x, t) = u_{i,\sigma}^k$ for $(x, t) \in T_{K,\sigma} \times [t_{k-1}, t_k)$, where $K \in \mathcal{T}$, $\sigma \in \mathcal{E}$, and $k = 1, \dots, N$. This function is piecewise constant on the dual mesh.

Let $\phi \in V_{\mathcal{T}}$ be such that $\|\phi\|_{1,4,\mathcal{T}} = 1$ and let $\hat{u}_m = P_{L^{\perp}} u_m$. We write (P_{ij}) for the matrix associated to $P_{L^{\perp}}$. Then

$$\begin{aligned} & \left| \sum_{K \in \mathcal{T}} \mathbf{m}(K) \partial_t^{\Delta t} \hat{u}_{m,i,K} \phi_K \right| = \left| \sum_{K \in \mathcal{T}} \sum_{j=1}^n \frac{\mathbf{m}(K)}{\Delta t} P_{ij} (u_{j,K}^k - u_{j,K}^{k-1}) \phi_K \right| \\ &\leq C \|\partial_t^{\Delta t} u_m^k\|_{-1,4,\mathcal{T}} \|\phi\|_{1,4,\mathcal{T}} \leq C. \end{aligned}$$

Together with estimate (33), this implies that

$$\sum_{k=1}^{N_m} \Delta t \|\partial_t^{\Delta t} \hat{u}_m^k\|_{-1,4,\mathcal{T}}^2 \leq C, \quad \sum_{k=1}^{N_m} \Delta t \|\hat{u}_m^k\|_{1,2,\mathcal{T}}^2 \leq C.$$

It is shown in [35, Sec. 6.1] that the discrete norms $\|\cdot\|_{1,2,\mathcal{T}}$ and $\|\cdot\|_{-1,4,\mathcal{T}}$ satisfy the assumptions of the compactness result in [26, Theorem 3.4]. Therefore, there exists a subsequence, which is not relabeled, such that $\hat{u}_m \rightarrow v$ strongly in $L^2(\Omega_T)$ as $m \rightarrow \infty$ for some $v \in L^2(\Omega_T)$. Moreover, up to a subsequence, we have $u_m \rightharpoonup u$ weakly in $L^2(\Omega_T)$ and consequently $\hat{u}_m = P_{L^{\perp}} u_m \rightharpoonup P_{L^{\perp}} u = \hat{u}$ weakly in $L^2(\Omega_T)$. This shows that $\hat{u} = v$.

Estimate (33) implies that $y_m := \nabla^m \hat{u}_m$ is uniformly bounded in $L^2(\Omega_T)$. Hence, there exists a subsequence (not relabeled) such that $y_m \rightharpoonup y$ weakly in $L^2(\Omega_T)$. We conclude as

in [8, Lemma 4.4] that $y = \nabla \widehat{u}$. We summarize:

$$(38) \quad u_m \rightharpoonup u, \quad y_m \rightharpoonup y = \nabla \widehat{u} \quad \text{weakly in } L^2(\Omega_T).$$

These convergences are not sufficient to pass to the limit in the term $u_{m,i,\sigma} \nabla^m (B u_m)_i$. The idea is to embed the problem in the larger space of Young measures. Let $\mathcal{P}(W)$ be the space of probability measures on $W := \mathbb{R}_{\geq}^n \times (L^\perp)^d$. Since the sequences (u_m) and (y_m) are bounded in $L^2(\Omega_T)$, we can apply [42, Theorem 6.2] to conclude the existence of a subsequence (not relabeled) and a family of probability measures $\mu = (\mu_{x,t})$ with $\mu_{x,t} \in \mathcal{P}(W)$ for a.e. $(x,t) \in \Omega_T$ such that the following holds:

If f is a continuous function on W vanishing at infinity and if the sequence $(f(u_m, y_m))$ converges weakly in $L^1(\Omega_T)$, then its weak limit, which we denote by $\overline{f(u_m, y_m)}$, satisfies

$$\overline{f(u_m, y_m)}(x, t) = \langle \mu_{x,t}, f(s, p) \rangle \quad \text{for a.e. } (x, t) \in \Omega_T.$$

In the above reasoning $T \in (0, \infty)$ was arbitrary. Hence, a diagonal argument allows us to choose μ independent of $T \in (0, \infty)$ such that $\mu \in L_w^\infty(\Omega \times (0, \infty); \mathcal{P}(W))$ and the weak convergences (38) hold for all $T > 0$. As a consequence,

$$u = \langle \mu, s \rangle, \quad \widehat{u} = \langle \mu, \widehat{s} \rangle, \quad y = \langle \mu, p \rangle \quad \text{a.e. in } \Omega \times (0, \infty),$$

where $\widehat{s} = P_{L^\perp} s$.

4.2. Convergence of the scheme. We show that μ is a dissipative measure-valued solution in the sense of Definition 1 satisfying (26). The proof adapts the strategy of [8] to the present situation, where only a weaker form of convergence is known to hold. Let $T \in (0, \infty)$, let $i \in \{1, \dots, n\}$, $\psi \in C_0^\infty(\Omega \times [0, T])$, and let $\eta_m = \max\{\Delta x_m, \Delta t_m\}$ be small enough such that $\text{supp}(\psi) \subset \{x \in \Omega : d(x, \partial\Omega) > \eta_m\} \times [0, T]$. We introduce

$$\begin{aligned} F_{10}^m &= - \int_0^T \int_\Omega u_{m,i} \partial_t \psi \, dx \, dt - \int_\Omega u_{m,i}^0(x) \psi(x, 0) \, dx, \\ F_{20}^m &= \int_0^T \int_\Omega u_{m,i,\sigma} \nabla^m (B \widehat{u}_m)_i \cdot \nabla \psi \, dx \, dt. \end{aligned}$$

The convergence results established above imply that, as $m \rightarrow \infty$,

$$F_{10}^m \rightarrow - \int_0^T \int_\Omega u_i \partial_t \psi \, dx \, dt - \int_\Omega u_i^{\text{in}}(x) \psi(x, 0) \, dx.$$

The limit in F_{20}^m is more involved. First, Lemma 12 implies that the term $u_{m,i,\sigma} (B \nabla^m \widehat{u}_m)_i$ is weakly relatively compact in $L^1(\Omega_T)$ and thus weakly convergent in $L^1(\Omega_T)$ along a subsequence. Second, we assert that

$$(39) \quad u_{m,\sigma} - u_m \rightarrow 0 \quad \text{in } L^1(\Omega_T) \text{ as } m \rightarrow \infty.$$

We proceed as in [41, Section 4.2], but since we cannot control the full gradient, we need to rely on the artificial diffusion. It follows from $\mathbf{m}(T_{K,\sigma}) = d_\sigma^2 \tau_\sigma / d$ that

$$\|u_{m,i,\sigma}^k - u_{m,i}^k\|_{0,1,\mathcal{T}_m} \leq C \sum_{K \in \mathcal{T}_m} \sum_{\sigma \in \mathcal{E}_{\text{int},K}} \mathbf{m}(T_{K,\sigma}) |u_{m,i,\sigma}^k - u_{m,i,K}^k|$$

$$\begin{aligned}
&\leq C \sum_{K \in \mathcal{T}_m} \sum_{\sigma \in \mathcal{E}_{\text{int},K}} \mathbf{m}(T_{K,\sigma}) |u_{m,i,L}^k - u_{m,i,K}^k| \\
&\leq C \sum_{\sigma=K|L \in \mathcal{E}_{\text{int}}} \mathbf{m}(T_{K,\sigma}) |u_{m,i,L}^k - u_{m,i,K}^k| \\
&\leq C \sum_{\sigma=K|L \in \mathcal{E}_{\text{int}}} d_\sigma^2 \tau_\sigma |u_{m,i,L}^k - u_{m,i,K}^k| \leq C \eta_m |u_{m,i}^k|_{1,1,\mathcal{T}_m},
\end{aligned}$$

where the constant $C > 0$ may change from line to line. We take the square, multiply by Δt_m , sum over $k = 1, \dots, N_m$, and use Lemma 11:

$$\sum_{k=1}^{N_m} \Delta t_m \|u_{m,i,\sigma}^k - u_{m,i}^k\|_{0,1,\mathcal{T}_m}^2 \leq C \eta_m^{2-\alpha}.$$

The right-hand side goes to zero as soon as $\alpha < 2$. Hence, $u_m - u_{m,\sigma} \rightarrow 0$ strongly in $L^2(0, T; L^1(\Omega))$, which implies (39). We note that, by interpolation, the strong convergence (39) together with the fact that the sequence $(u_m - u_{m,\sigma})_m$ is uniformly bounded in $L^2(\Omega_T)$ implies that

$$u_m - u_{m,\sigma} \rightarrow 0 \quad \text{strongly in } L^p(\Omega_T) \text{ for every } p < 2.$$

We now assert that, as a consequence of (39), the sequence $(u_{m,\sigma}, \nabla^m \widehat{u}_m)$ generates the same Young measure μ as $(u_m, \nabla^m \widehat{u}_m)$ (after possibly passing to another subsequence). Indeed, since μ is uniquely determined by its action on C_0 -functions, to verify the assertion, it suffices to show that

$$\lim_{m \rightarrow \infty} \int_{\Omega_T} (f(u_{m,\sigma}, \nabla^m \widehat{u}_m) - f(u_m, \nabla^m \widehat{u}_m)) \phi \, dx \, dt = 0$$

for all $f \in C_0(W)$ and $\phi \in L^1(\Omega_T)$. This follows from (39) and the dominated convergence theorem, because functions $f \in C_0(W)$ are uniformly continuous. Since $u_{m,i,\sigma}(B \nabla^m \widehat{u}_m)_i$ is weakly convergent in $L^1(\Omega_T)$, we thus infer that

$$\overline{u_{m,i,\sigma}(B \nabla^m \widehat{u}_m)_i}(x, t) = \int_W s_i(Bp)_i \, d\mu_{x,t}(s, p) = \langle \mu_{x,t}, s_i(Bp)_i \rangle.$$

We conclude that

$$F_{20}^m \rightarrow \int_0^T \int_\Omega \langle \mu_{x,t}, s_i(Bp)_i \rangle \, dx \, dt.$$

Let $\psi_K^k = \psi(x_K, t_k)$ and multiply (16) by $\Delta t \psi_K^{k-1}$ and sum over $K \in \mathcal{T}_m$, $k = 1, \dots, N_m$. This gives $F_1^m + F_2^m + F_3^m = 0$, where

$$\begin{aligned}
F_1^m &= \sum_{k=1}^{N_m} \sum_{K \in \mathcal{T}_m} \mathbf{m}(K) (u_{i,K}^k - u_{i,K}^{k-1}) \psi_K^{k-1}, \\
F_2^m &= - \sum_{k=1}^{N_m} \Delta t_m \sum_{K \in \mathcal{T}_m} \sum_{\sigma \in \mathcal{E}_{\text{int},K}} \tau_\sigma u_{i,\sigma}^k D_{K,\sigma} (B u^k)_i \psi_K^{k-1},
\end{aligned}$$

$$F_3^m = -\eta_m^\alpha \sum_{k=1}^{N_m} \Delta t_m \sum_{K \in \mathcal{T}_m} \sum_{\sigma \in \mathcal{E}_{\text{int},K}} \tau_\sigma \mathbf{D}_{K,\sigma} u_i^k \psi_K^{k-1}.$$

We infer from the Cauchy–Schwarz inequality and Lemma 11 that

$$|F_3^m| \leq \eta_m^\alpha \left(\sum_{k=1}^{N_m} \Delta t_m |u_i^k|_{1,4/3,\mathcal{T}_m}^2 \right)^{1/2} \left(\sum_{k=1}^{N_m} \Delta t_m |\psi^{k-1}|_{1,4,\mathcal{T}_m}^2 \right)^{1/2} \leq C \eta_m^{\alpha/2} \rightarrow 0$$

as $m \rightarrow \infty$. We claim that $F_{j0}^m - F_j^m \rightarrow 0$ for $j = 1, 2$.

For the limit of $F_{10}^m - F_1^m$, we use as in the proof of [8, Theorem 5.2] discrete integration by parts in time:

$$\begin{aligned} F_1^m &= - \sum_{k=1}^{N_m} \sum_{K \in \mathcal{T}_m} \mathbf{m}(K) u_{i,K}^k (\psi_K^k - \psi_K^{k-1}) - \sum_{K \in \mathcal{T}_m} \mathbf{m}(K) u_{i,K}^0 \psi_K^0 \\ &= - \sum_{k=1}^{N_m} \sum_{K \in \mathcal{T}_m} \int_{t_{k-1}}^{t_k} \int_K u_{i,K}^k \partial_t \psi(x_K, t) dx dt - \sum_{K \in \mathcal{T}_m} \int_K u_{i,K}^0 \psi(x_K, 0) dx, \\ F_{10}^m &= - \sum_{k=1}^{N_m} \sum_{K \in \mathcal{T}_m} \int_{t_{k-1}}^{t_k} \int_K u_{i,K}^k \partial_t \psi(x, t) dx dt - \sum_{K \in \mathcal{T}_m} \int_K u_{i,K}^0 \psi(x, 0) dx. \end{aligned}$$

It follows from the regularity of ψ that

$$|F_{10}^m - F_1^m| \leq C(\Omega_T) \|u_i^k\|_{L^\infty(0,T;L^2(\Omega))} \|\psi\|_{C^2(\bar{\Omega}_T)} \Delta t_m \rightarrow 0 \quad \text{as } m \rightarrow \infty.$$

We deduce from the definition of the discrete gradient that

$$\begin{aligned} F_{20}^m &= \sum_{k=1}^{N_m} \int_{t_k}^{t_{k-1}} \sum_{\sigma \in \mathcal{E}_{\text{int}}} \frac{\tilde{\mathbf{m}}(\sigma)}{\mathbf{m}(T_{K,\sigma})} u_{i,\sigma}^k \mathbf{D}_{K,\sigma}(B\hat{u}_m)_i \int_{T_{K,\sigma}} \nabla \psi \cdot \nu_{K,\sigma} dx dt, \\ F_2^m &= \sum_{k=1}^{N_m} \int_{t_k}^{t_{k-1}} \sum_{\sigma \in \mathcal{E}_{\text{int}}} \frac{\tilde{\mathbf{m}}(\sigma)}{d_\sigma} u_{i,\sigma}^k \mathbf{D}_{K,\sigma}(B\hat{u}_m)_i \mathbf{D}_{K,\sigma} \psi^{k-1} dt. \end{aligned}$$

This gives

$$\begin{aligned} |F_{20}^m - F_2^m| &\leq \sum_{k=1}^{N_m} \sum_{\sigma \in \mathcal{E}_{\text{int}}} \tilde{\mathbf{m}}(\sigma) u_{i,\sigma}^k |\mathbf{D}_{K,\sigma}(B\hat{u}_m)_i| \\ &\quad \times \left| \int_{t_{k-1}}^{t_k} \left(\frac{\mathbf{D}_{K,\sigma} \psi^{k-1}}{d_\sigma} - \frac{1}{\mathbf{m}(T_{K,\sigma})} \int_{T_{K,\sigma}} \nabla \psi \cdot \nu_{K,\sigma} dx \right) dt \right|. \end{aligned}$$

By the proof of Theorem 5.1 in [8], there exists $C > 0$, independent of η_m , such that

$$\left| \int_{t_{k-1}}^{t_k} \left(\frac{\mathbf{D}_{K,\sigma} \psi^{k-1}}{d_\sigma} - \frac{1}{\mathbf{m}(T_{K,\sigma})} \int_{T_{K,\sigma}} \nabla \psi \cdot \nu_{K,\sigma} dx \right) dt \right| \leq C \Delta t_m \eta_m,$$

which shows, using the Cauchy-Schwarz inequality, that

$$\begin{aligned} |F_{20}^m - F_2^m| &\leq C\eta_m \sum_{k=1}^{N_m} \Delta t_m \sum_{\sigma \in \mathcal{E}_{\text{int}}} \tilde{\mathbf{m}}(\sigma) u_{i,\sigma}^k |D_\sigma(B\hat{u}_m)_i| \\ &\leq C\eta_m \sum_{k=1}^{N_m} \Delta t_m |(Bu_m^k)_i|_{1,2,\mathcal{T}_m} \left(\sum_{K \in \mathcal{T}_m} \sum_{\sigma \in \mathcal{E}_{\text{int},K}} \tilde{\mathbf{m}}(\sigma) d_\sigma (u_{i,\sigma}^k)^2 \right)^{1/2}. \end{aligned}$$

We conclude from the Cauchy-Schwarz inequality, estimate (37), and the uniform bounds (33)–(34) that

$$\begin{aligned} |F_{20}^m - F_2^m| &\leq C(\zeta)\eta_m \left(\sum_{k=1}^{N_m} \Delta t_m |(Bu^k)_i|_{1,2,\mathcal{T}_m}^2 \right)^{1/2} \left(\sum_{k=1}^{N_m} \Delta t_m \|u_i^k\|_{0,2,\mathcal{T}_m}^2 \right)^{1/2} \\ &\leq C(\zeta)\eta_m \rightarrow 0 \quad \text{as } m \rightarrow \infty. \end{aligned}$$

We deduce that $F_{10}^m + F_{20}^m \rightarrow 0$ as $m \rightarrow \infty$. Then, because of $F_1^m + F_2^m + F_3^m = 0$,

$$F_{10}^m + F_{20}^m = (F_{10}^m - F_1^m) + (F_{20}^m - F_2^m) - F_3^m \rightarrow 0 \quad \text{as } m \rightarrow \infty,$$

which proves that u_i satisfies

$$\int_0^T \int_\Omega u_i \partial_t \psi dx dt + \int_\Omega u_i^{\text{in}} \psi(0) dx = \int_0^T \int_\Omega \langle \mu_{x,t}, s_i(Bp)_i \rangle \cdot \nabla \psi dx dt.$$

Hence, in the sense of distributions,

$$(40) \quad \partial_t u_i = \text{div} \langle \mu, s_i(Bp)_i \rangle, \quad u_i(0) = u_i^{\text{in}}, \quad i = 1, \dots, n.$$

4.3. Entropy inequalities. We verify the entropy inequalities (10) and (11). The definition of u_m^0 and the regularity $u^{\text{in}} \in L^2(\Omega)$ imply the strong convergence $u_m^0 \rightarrow u^{\text{in}}$ in $L^2(\Omega)$ as $m \rightarrow \infty$.

Re Shannon: Since $(u_m)_m$ is bounded in $L^2(\Omega_T)$, the sequence $(h_S(u_m))_m \subset L^1(\Omega_T)$ is equi-integrable. After passing to a subsequence, we can therefore assume that $(h_S(u_m))_m$ is weakly convergent in $L^1(\Omega_T)$, which implies that for a.e. $(x, t) \in \Omega_T$,

$$\langle \mu_{x,t}, h_S(s) \rangle = \overline{h_S(u_m)}(x, t).$$

The dual mesh allows us to rewrite the Shannon entropy dissipation in (24) as

$$\sum_{j=1}^k \Delta t_m |B^{1/2} u_m^j|_{1,2,\mathcal{T}_m}^2 = \int_0^{t_k} \int_\Omega |\nabla^m(B^{1/2} u_m)|^2 dx d\tau.$$

Given $0 < \delta \ll 1$, let m be large enough such that $\Delta t_m < \delta$. Then (24) entails for all $t \in [\delta, T]$ that

$$(41) \quad H_S(u_m(t)) + \int_0^{t-\delta} \int_\Omega |\nabla^m(B^{1/2} u_m)|^2 dx d\tau \leq H_S(u_m^0).$$

Next, let $\xi \in C_c^1([0, T]; \mathbb{R}_{\geq})$ with $\xi(0) = 1$ and $\xi' \leq 0$. We multiply the last inequality by the non-negative function $-\xi'(t)$ and integrate over $t \in [\delta, T]$:

$$\int_{\delta}^T \int_{\Omega} (-\xi'(t)) h_S(u_m(t)) dx dt + \int_{\delta}^T (-\xi'(t)) \int_0^{t-\delta} \int_{\Omega} |\nabla^m(B^{1/2}u_m)|^2 dx d\tau dt \leq \xi(\delta) H_S(u_m^0).$$

Using the weak convergence $\nabla^m(B^{1/2}u_m) = \nabla^m(B^{1/2}\hat{u}_m) \rightharpoonup \nabla(B^{1/2}\hat{u})$ in $L^2(\Omega_T)$ and the convexity of the continuous functional $L^2(\Omega_T) \ni v \mapsto \int_{\delta}^T (-\xi'(t)) \int_0^{t-\delta} \int_{\Omega} |v(\tau, x)|^2 dx d\tau dt$, we take the $\liminf_{m \rightarrow \infty}$ in the above inequality. This yields

$$\int_{\delta}^T (-\xi'(t)) \int_{\Omega} \langle \mu_{x,t}, h_S(s) \rangle dx dt + \int_{\delta}^T (-\xi'(t)) \int_0^{t-\delta} \int_{\Omega} |\nabla(B^{1/2}\hat{u})|^2 dx d\tau dt \leq \xi(\delta) H_S(u^{\text{in}}).$$

As $\delta \downarrow 0$, we infer

$$\int_0^T (-\xi'(t)) \int_{\Omega} \langle \mu_{x,t}, h_S(s) \rangle dx dt + \int_0^T (-\xi'(t)) \int_0^t \int_{\Omega} |\nabla(B^{1/2}\hat{u})|^2 dx d\tau dt \leq H_S(u^{\text{in}}).$$

This is true for all $\xi \in C_c^1([0, T]; \mathbb{R}_{\geq})$ with $\xi(0) = 1$ and $\xi' \leq 0$. We then choose $\xi = \xi_{\ell}$ with $(\xi_{\ell})_{\ell}$ a suitable approximation of the Heaviside-type function $1_{[0, t_0]}$ and let $\ell \rightarrow \infty$ to deduce (10) at time $t = t_0$ for a.e. $t_0 \in (0, T]$.

Re Rao: Next, we verify (11) and the time monotonicity of $H_R(u)$. Since (\hat{u}_m) converges strongly to \hat{u} in $L^2(\Omega_T)$, we find that

$$H_R(u(t)) = \frac{1}{2} \int_{\Omega} |B^{1/2}\hat{u}(t)|^2 dx = \frac{1}{2} \lim_{m \rightarrow \infty} \sum_{K \in \mathcal{T}_m} \mathbf{m}(K) |B^{1/2}\hat{u}_m(t)|^2 = \lim_{m \rightarrow \infty} H_R(u_m(t)).$$

Together with the non-increase of $[0, \infty) \ni t \mapsto H_R(u_m(t))$ (cf. Theorem 3), this implies that the mapping $t \mapsto H_R(u(t))$ is nonincreasing. It remains to show (11). To this end, we let $0 < \delta \ll 1$ and take m large enough so that $\Delta t_m < \delta$. Then it follows from the discrete Rao entropy inequality (25) that

$$H_R(u_m(t)) + \sum_{i=1}^n \int_0^{t-\delta} \int_{\Omega} u_{m,i,\sigma} |(B\nabla^m \hat{u}_m)_i|^2 dx d\tau \leq H_R(u_m^0).$$

To estimate below the $\liminf_{m \rightarrow \infty}$ of the second term on the left-hand side, we recall that μ is also the Young measure associated with $(u_{m,\sigma}, \nabla^m \hat{u}_m)$. We therefore infer from [42, Theorem 6.11] for every $i \in \{1, \dots, n\}$

$$\int_0^{t-\delta} \int_{\Omega} \langle \mu_{x,\tau}, s_i |(Bp)_i|^2 \rangle dx d\tau \leq \liminf_{m \rightarrow \infty} \int_0^{t-\delta} \int_{\Omega} u_{m,i,\sigma} |(B\nabla^m \hat{u}_m)_i|^2 dx d\tau.$$

Thus, in the limit $m \rightarrow \infty$ we deduce

$$H_R(u(t)) + \sum_{i=1}^n \int_0^{t-\delta} \int_{\Omega} \langle \mu_{x,\tau}, s_i |(Bp)_i|^2 \rangle dx d\tau \leq H_R(u^{\text{in}}),$$

and sending $\delta \downarrow 0$ we obtain (11).

4.4. Separation of the \widehat{s} -component. For simplicity, we only prove identity (26) in the case where $f = f(s) \in C_0(\mathbb{R}_{\geq}^n)$. Let $g(s_1, s_2) = f(s_1 + s_2)$, defined on the convex set

$$Q = \{(s_1, s_2) \in L^\perp \times L : s_1 + s_2 \in \mathbb{R}_{\geq}^n\}.$$

Since the sequence (\widehat{u}_m) converges strongly in $L^2(\Omega_T)$, the Young measure $\widetilde{\mu}$, generated by $(P_{L^\perp} u_m, P_L u_m)$, has the form $\widetilde{\mu}_{x,t} = \delta_{\widehat{u}(x,t)} \otimes \nu_{x,t}$, where $\nu = (\nu_{x,t})$ is the Young measure generated by the sequence $(P_L u_m)$ [42, Prop. 6.13]. Hence, by construction of μ and $\widetilde{\mu}$,

$$\begin{aligned} \int_{\mathbb{R}_{\geq}^n} f(s) d\mu_{x,t}(s) &= \int_Q g(s_1, s_2) d\widetilde{\mu}_{x,t}(s_1, s_2) = \int_Q g(\widehat{u}(x, t), s_2) d\widetilde{\mu}_{x,t}(s_1, s_2) \\ &= \int_Q f(\widehat{u}(x, t) + s_2) d\widetilde{\mu}_{x,t}(s_1, s_2). \end{aligned}$$

It follows that $\langle \mu_{x,t}, f(s) \rangle = \langle \mu_{x,t}, f(\widehat{u}(x, t) + s_2) \rangle$ for all $f = f(s) \in C_0(\mathbb{R}_{\geq}^n)$ and a.a. (x, t) .

4.5. Time regularity. The time regularity for the density part $u = \langle \mu, s \rangle$ of the barycenter of μ follows from the continuity equation (40). To see this, we first note that due to $b_{ii} > 0, b_{ij} \geq 0$, and property (26),

$$(42) \quad \left\langle \mu_{x,t}, \sum_{i=1}^n s_i^2 \right\rangle \leq C \langle \mu_{x,t}, |B^{1/2} \widehat{s}|^2 \rangle = C |B^{1/2} \widehat{u}(x, t)|^2 = Ch_R(u(x, t))$$

for a.e. $(x, t) \in \Omega \times (0, \infty)$. Then we use Jensen's inequality to estimate for $i = 1, \dots, n$,

$$\begin{aligned} \|\langle \mu, s_i(Bp)_i \rangle\|_{L^2(0, \infty; L^{4/3}(\Omega))}^2 &= \int_0^\infty \left(\int_\Omega \langle \mu_{x,t}, |s_i(Bp)_i|^{4/3} \rangle dx \right)^{3/2} dt \\ &\leq \int_0^\infty \left(\int_\Omega \langle \mu_{x,t}, s_i^2 \rangle^{1/3} \langle \mu_{x,t}, |s_i(Bp)_i|^2 \rangle^{2/3} dx \right)^{3/2} dt \\ &\leq \int_0^\infty \left(\int_\Omega \langle \mu_{x,t}, s_i^2 \rangle dx \right)^{1/2} \int_\Omega \langle \mu_{x,t}, |s_i(Bp)_i|^2 \rangle dx dt \\ &\leq \left(\operatorname{ess\,sup}_{0 < t < \infty} \int_\Omega \langle \mu_{x,t}, s_i^2 \rangle dx \right) \left(\int_0^\infty \int_\Omega \langle \mu_{x,t}, |s_i(Bp)_i|^2 \rangle dx dt \right), \end{aligned}$$

where Hölder's inequality was applied several times. It therefore follows from (40) that

$$\|\partial_t u_i\|_{L^2(0, \infty; W^{1,4}(\Omega)^*)} \leq \|\langle \mu, s_i(Bp)_i \rangle\|_{L^2(0, \infty; L^{4/3}(\Omega))} \leq CH_R(u^{\text{in}}),$$

where the last step also uses (11) and (42). This finishes the proof of Theorem 4.

Remark 14 (Curved domains). We claim that Theorems 3 and 4 also hold for curved Lipschitz domains $\Omega \subset \mathbb{R}^d$. The triangulation then contains control volumes with curved segments that are part of $\partial\Omega$. The analysis of this section is still possible, since we consider no-flux boundary conditions and no boundary values need to be defined. The analysis has to be adapted in two points. First, the convergence of the scheme is typically proved on polygonal meshes and the error between the curved cell and the polygonal cell (which is of order $(\Delta x)^{d+1}$) needs to be taken into account. Second, as the compactness of the

approximate sequence has been established for polygonal domains [26], the error between the approximate sequence and its extension by zero to the polygonal domain has to be estimated. In two space dimensions, it is of order Δx ; see [40, Prop. 4.14] for details. The drawback of this approach is that one has to perform numerical integrations over the curved elements, which may be cumbersome in particular in three space dimensions.




FIGURE 1. Triangulation of a curved domain.

Here we report on the simple approach of [19]. The idea is to cover Ω by additional control volumes and to estimate the integral error. To simplify the presentation, let $\Omega \subset \mathbb{R}^2$ and let \mathcal{T} be a sufficiently fine triangulation of Ω into triangles. To each cell with two vertices on $\partial\Omega$, we add the reflected triangle to the triangulation such that $\Omega \subset \cup_{K \in \mathcal{T}^*} K$, where \mathcal{T}^* consists of all cells $K \in \mathcal{T}$ and the associated reflected cells K_r with nonempty intersection with Ω ; see Figure 1. Denoting by $\omega_r = K_r \cap \Omega$ if $K_r \cap \Omega \neq \emptyset$ and $\omega_c = K \setminus \Omega$ if $K_r \cap \Omega = \emptyset$, the domain splits into

$$\Omega = \Omega_h \cup \Omega_r \setminus \Omega_c := \left(\bigcup_{K \in \mathcal{T}} K \right) \cup \left(\bigcup_{\omega_r} \omega_r \right) \setminus \left(\bigcup_{\omega_c} \omega_c \right).$$

We can perform the numerical analysis on $V_{\mathcal{T}^*}$ as in Sections 3 and 4. For the convergence of the scheme, we need to show that the difference of the integrals over Ω_h and Ω vanishes when $\eta_m \rightarrow 0$. The difference consists of two contributions: the integral over Ω_r and the integral over Ω_c . We illustrate the convergence for the integral

$$\left| \int_{\Omega_r} u_{m,i,\sigma} \nabla^m (B\hat{u}_m)_i \cdot \nabla \psi dx \right| \leq C \sum_{\omega_r} \mathbf{m}(\omega_r) \|u_{m,i,\sigma}\|_{0,\infty,\omega_r} \|\nabla^m (B\hat{u}_m)_i\|_{0,\infty,\omega_r},$$

where ψ is a smooth test function. By the inverse inequality [13, Section 21.1]

$$\|v\|_{0,\infty,\omega_r} \leq \|v\|_{0,\infty,K_r} \leq C(\Delta x)^{-d/2} \|v\|_{0,2,K_r},$$

the bound $\mathbf{m}(\omega_r) \leq C(\Delta x)^{d+1}$ (which is valid under certain regularity conditions on the mesh), and the Cauchy–Schwarz inequality, we have

$$\left| \int_{\Omega_r} u_{m,i,\sigma} \nabla^m (B\hat{u}_m)_i \cdot \nabla \psi dx \right| \leq C\Delta x \left(\sum_{K_r} \|u_{m,i,\sigma}\|_{0,2,K_r}^2 \right)^{1/2} \left(\sum_{K_r} \|\nabla^m (B\hat{u}_m)_i\|_{0,2,K_r}^2 \right)^{1/2}$$

$$\leq C\Delta x \rightarrow 0 \quad \text{as } \eta \rightarrow 0,$$

taking into account the uniform bounds from (24) and (25). In a similar way, the integral over Ω_c tends to zero as $\eta \rightarrow 0$. \square

5. STABILITY

In this section, we prove Theorem 7. Let μ be a dissipative measure-valued solution and let $v \in C^1(\overline{\Omega}_T)$ be a positive solution satisfying (4), (2). We introduce the relative Shannon and Rao entropies by, respectively,

$$\begin{aligned} H_S^{\text{mv}}(u(t)|v(t)) &= \sum_{i=1}^n \int_{\Omega} (\langle \mu_{x,t}, \mathfrak{h}(s_i) \rangle - \mathfrak{h}(v_i(x,t)) - \mathfrak{h}'(v_i(x,t)) \cdot (u_i - v_i)(x,t)) dx, \\ &= \int_{\Omega} \sum_{i=1}^n (\langle \mu_{x,t}, s_i \log s_i \rangle - u_i \log v_i - (u_i - v_i)) dx \geq 0, \\ H_R(u(t)|v(t)) &= \frac{1}{2} \int_{\Omega} |B^{1/2}(u - v)(x,t)|^2 dx \geq 0, \end{aligned}$$

where $\mathfrak{h}(z) = z(\log z - 1) + 1$ for $z \geq 0$. We further define the usual relative Shannon entropy $H_S(u|v) = \int_{\Omega} \sum_{i=1}^n (u_i \log u_i - u_i \log v_i - (u_i - v_i)) dx$. Furthermore, we set

$$\begin{aligned} H_{\text{rel}}^{\text{mv}}(u|v) &= H_S^{\text{mv}}(u|v) + H_R(u|v), \\ H_{\text{rel}}(u|v) &= H_S(u|v) + H_R(u|v). \end{aligned}$$

We first compute the relative entropy inequalities.

Lemma 15 (Relative entropy inequalities). *Suppose that Ω has a Lipschitz boundary. Let μ be a dissipative measure-valued solution, $u := \langle \mu, s \rangle$, and let $v \in C^1(\overline{\Omega}_T)$ be a positive solution to (4), (2) for $t \in (0, T)$ (in the weak sense). Then, for a.e. $t \in (0, T)$,*

$$(43) \quad \begin{aligned} H_S^{\text{mv}}(u(t)|v(t)) &+ \int_0^t \int_{\Omega} \langle \mu_{x,\tau}, |B^{1/2}(p - \nabla v)|^2 \rangle dx d\tau \\ &+ \int_0^t \int_{\Omega} \left\langle \mu_{x,\tau}, \sum_{i=1}^n (v_i - s_i) \nabla \log v_i \cdot (B(p - \nabla v))_i \right\rangle dx d\tau \leq H_S(u^{\text{in}}|v(0)), \end{aligned}$$

$$(44) \quad \begin{aligned} H_R(u(t)|v(t)) &+ \int_0^t \int_{\Omega} \sum_{i=1}^n \langle \mu_{x,\tau}, s_i | (B(p - \nabla v))_i|^2 \rangle dx d\tau \\ &+ \int_0^t \int_{\Omega} \left\langle \mu_{x,\tau}, \sum_{i=1}^n (s_i - v_i) \nabla (Bv)_i \cdot (B(p - \nabla v))_i \right\rangle dx d\tau \leq H_R(u^{\text{in}}|v(0)). \end{aligned}$$

Proof. It follows from (9) that for all $i = 1, \dots, n$ and $\phi \in L^2(0, T; W^{1,4}(\Omega))$

$$(45) \quad \int_0^T (\partial_t u_i, \phi)_{W^{1,4}(\Omega)^*} dt = - \int_0^T \int_{\Omega} \langle \mu_{x,t}, s_i (Bp)_i \rangle \cdot \nabla \phi dx dt,$$

where $(\cdot, \cdot)_{W^{1,4}(\Omega)^*}$ denotes the duality pairing between $W^{1,4}(\Omega)^*$ and $W^{1,4}(\Omega)$.

Re Shannon: The solution property and positivity of v imply that for every $\psi \in C^1(\overline{\Omega_T}; \mathbb{R}^n)$,

$$\begin{aligned} - \sum_{i=1}^n \int_{\Omega} (\partial_t \log v_i) \psi_i dx &= \int_{\Omega} \sum_{i=1}^n v_i \nabla (Bv)_i \cdot \nabla \left(\frac{\psi_i}{v_i} \right) dx \\ &= \int_{\Omega} \nabla v : \nabla (B\widehat{\psi}) dx - \sum_{i=1}^n \int_{\Omega} \nabla (Bv)_i \cdot (\nabla \log v_i) \psi_i dx. \end{aligned}$$

Let $t \in (0, T)$ be arbitrary. An integration over $\tau \in (0, t)$ and an approximation argument imply that for all $\psi \in L^2(\Omega_T; \mathbb{R}^n)$ with $\nabla B\widehat{\psi} \in L^2(\Omega_T)$,

$$\begin{aligned} - \sum_{i=1}^n \int_0^t \int_{\Omega} (\partial_t \log v_i) \psi_i dx d\tau &= \int_0^t \int_{\Omega} \nabla v : \nabla B\widehat{\psi} dx d\tau \\ &\quad - \sum_{i=1}^n \int_0^t \int_{\Omega} \nabla (Bv)_i \cdot (\nabla \log v_i) \psi_i dx d\tau. \end{aligned}$$

The choice $\psi = u = \langle \mu, s \rangle$ and the property $\nabla B\widehat{u} = B\langle \mu, p \rangle = \langle \mu, Bp \rangle$ lead to

$$\begin{aligned} - \sum_{i=1}^n \int_0^t \int_{\Omega} (\partial_t \log v_i) u_i dx d\tau &= \\ &= \int_0^t \int_{\Omega} \nabla v : \nabla B\widehat{u} dx d\tau - \sum_{i=1}^n \int_0^t \int_{\Omega} \nabla (Bv)_i \cdot (\nabla \log v_i) u_i dx d\tau \\ &= \int_0^t \int_{\Omega} \langle \mu_{x,t}, B^{1/2} \nabla v : B^{1/2} p \rangle dx d\tau - \sum_{i=1}^n \int_0^t \int_{\Omega} \langle \mu_{x,t}, s_i \nabla \log v_i \cdot \nabla (Bv)_i \rangle dx d\tau. \end{aligned}$$

Next, we use $\phi_i = 1_{[0,t]} \log v_i$ as a test function in the weak formulation (45), multiply by -1 , and sum over $i = 1, \dots, n$:

$$- \sum_{i=1}^n \int_0^t (\partial_t u_i, \log v_i)_{W^{1,4}(\Omega)^*} d\tau = \sum_{i=1}^n \int_0^t \int_{\Omega} \langle \mu_{x,t}, s_i (Bp)_i \rangle \cdot \nabla \log v_i dx d\tau.$$

We add the previous two equations:

$$\begin{aligned} - \int_0^t \frac{d}{dt} \int_{\Omega} \sum_{i=1}^n (\log v_i) u_i dx d\tau &= \int_0^t \int_{\Omega} \langle \mu_{x,t}, B^{1/2} \nabla v : B^{1/2} p \rangle dx d\tau \\ &\quad + \int_0^t \int_{\Omega} \langle \mu_{x,t}, \sum_{i=1}^n s_i \nabla \log v_i \cdot (B(p - \nabla v))_i \rangle dx d\tau. \end{aligned}$$

Combined with the identity

$$\int_0^t \int_{\Omega} \langle \mu_{x,t}, B^{1/2} \nabla v : B^{1/2} p \rangle dx d\tau - \int_0^t \int_{\Omega} \langle \mu_{x,t}, |B^{1/2} \nabla v|^2 \rangle dx d\tau$$

$$- \int_0^t \int_{\Omega} \left\langle \mu_{x,t}, \sum_{i=1}^n v_i \nabla \log v_i \cdot (B(p - \nabla v))_i \right\rangle dx d\tau = 0,$$

the Shannon entropy inequality (10), and mass conservation $(d/dt) \int_{\Omega} v_i dx = 0$, this gives (43).

Re Rao: Since $v_i \nabla (Bv)_i \in L^2(\Omega_T)$, we can test the equation for v with $1_{[0,t]} B(v - u) \in L^2(0, T; H^1(\Omega))$. This yields

$$\int_0^t \int_{\Omega} \partial_t v^T B(v - u) dx d\tau = - \int_0^t \int_{\Omega} \sum_{i=1}^n v_i \nabla (Bv)_i \cdot \nabla (B(v - u))_i dx d\tau.$$

Next, we choose $\phi = 1_{[0,t]} (Bv)_i$ in equation (45) for u and sum over $i = 1, \dots, n$:

$$- \int_0^t (\partial_t u, Bv)_{W^{1,4}(\Omega)^*} d\tau = \int_0^t \int_{\Omega} \sum_{i=1}^n \langle \mu_{x,t}, s_i (Bp)_i \rangle \cdot (B\nabla v)_i dx d\tau.$$

Adding to these identities the Rao entropy inequality (11) and rearranging terms gives

$$\begin{aligned} \frac{1}{2} \int_0^t \frac{d}{dt} \int_{\Omega} (u - v)^T B(u - v) dx d\tau &\leq - \int_0^t \int_{\Omega} \sum_{i=1}^n \langle \mu_{x,t}, s_i | (B(p - \nabla v))_i |^2 \rangle dx d\tau \\ &\quad - \int_0^t \int_{\Omega} \sum_{i=1}^n \langle \mu_{x,t}, (s_i - v_i) (B\nabla v)_i \cdot (B(p - \nabla v))_i \rangle dx d\tau, \end{aligned}$$

which implies (44), concluding the proof. \square

We proceed with the proof of Theorem 7. To this end, we estimate the last integrals on the left-hand sides of (43) and (44). We infer from Young's inequality that

$$\begin{aligned} (46) \quad &\left| \sum_{i=1}^n (v_i - s_i) \nabla \log v_i \cdot (B(p - \nabla v))_i \right| \\ &\leq \frac{1}{4} |B^{1/2}(p - \nabla v)|^2 + C \sum_{i=1}^n |\nabla \log v_i|^2 (s_i - v_i)^2 \\ &\leq \frac{1}{4} |B^{1/2}(p - \nabla v)|^2 + C |s - v|^2, \end{aligned}$$

$$\begin{aligned} (47) \quad &\left| \sum_{i=1}^n (s_i - v_i) \nabla (Bv)_i \cdot (B(p - \nabla v))_i \right| \\ &\leq \frac{1}{4} |B^{1/2}(p - \nabla v)|^2 + C \sum_{i=1}^n |\nabla (Bv)_i|^2 (s_i - v_i)^2 \\ &\leq \frac{1}{4} |B^{1/2}(p - \nabla v)|^2 + C |s - v|^2, \end{aligned}$$

where $C > 0$ depends on the $L^\infty(\Omega_T)$ norms of $|\nabla \log v_i|$ and $\nabla (Bv)_i$. Thus, adding the relative entropy inequalities (43) and (44), the first terms on the right-hand sides of (46)

and (47) can be absorbed by the left-hand side of (43) such that

$$(48) \quad \begin{aligned} H_{\text{rel}}^{\text{mv}}(u(t)|v(t)) + \int_0^t \int_{\Omega} \left\langle \mu_{x,t}, \frac{1}{2} |B^{1/2}(p - \nabla v)|^2 \right\rangle dx d\tau \\ \leq C \int_0^t \int_{\Omega} \langle \mu_{x,t}, |s - v|^2 \rangle dx d\tau + H_{\text{rel}}(u^{\text{in}}|v(0)). \end{aligned}$$

The coercivity estimate from Lemma 17 in Appendix A implies that

$$\int_{\Omega} \langle \mu_{x,t}, |s - v(x,t)|^2 \rangle dx \leq C H_{\text{rel}}^{\text{mv}}(u(t)|v(t)).$$

We insert this bound into (48) and invoke Gronwall's inequality to deduce that

$$H_{\text{rel}}^{\text{mv}}(u(t)|v(t)) + \int_0^t \int_{\Omega} \left\langle \mu_{x,\tau}, \frac{1}{2} |B^{1/2}(p - \nabla \hat{v}(x,\tau))|^2 \right\rangle dx d\tau \leq e^{Ct} H_{\text{rel}}(u^{\text{in}}|v(0)) = 0,$$

where the last equality follows from $v(0) = u^{\text{in}}$. Hence, $\mu_{x,t} = \delta_{v(x,t)} \otimes \delta_{\nabla \hat{v}(x,t)}$ for a.e. $(t,x) \in \Omega \times (0,T)$, which finishes the proof of Theorem 7.

6. LONG-TIME ASYMPTOTICS

In this section, we prove Theorem 9. First, we verify that $\mathfrak{S}_{\mathbf{m}} \subset L^{\infty}(\Omega)$. Indeed, if $v \in \mathfrak{S}_{\mathbf{m}}$, the vector Bv is constant and $\int_{\Omega} Bv dx = B\mathbf{m}$, which implies that $Bv = (B\mathbf{m})/|\Omega|$. Since the entries of B and the components of v are nonnegative, $v_i \leq (B\mathbf{m})_i/(b_{ii}|\Omega|)$ for all $i = 1, \dots, n$. This proves the claim.

The entropy inequalities (10)–(11) show that

$$\int_0^{\infty} \|\nabla(B^{1/2}u)\|_{L^2(\Omega)}^2 dt < \infty, \quad \sup_{0 < t < \infty} \|u(t)\|_{L^2(\Omega)} < \infty.$$

Thus, there exists a sequence $(t_k) \subset (0, \infty)$ with $t_k \rightarrow \infty$ such that $u(t_k) \rightharpoonup u^*$ weakly in $L^2(\Omega)$ and $B^{1/2}u(t_k) \rightarrow B^{1/2}u^*$ strongly in $L^2(\Omega)$ as $k \rightarrow \infty$. Since $\int_{\Omega} u(t_k) dx = \mathbf{m}$ and the sequence $(\nabla(B^{1/2}u(t_k)))$ converges to zero in the $L^2(\Omega)$ norm, we find that $\int_{\Omega} u^* dx = \mathbf{m}$ and $\nabla(B^{1/2}u^*) = 0$. This implies that $u^* \in \mathfrak{S}_{\mathbf{m}}$. Moreover, we deduce from the strong convergence that

$$\lim_{k \rightarrow \infty} H_R(u(t_k)|u^*) = \frac{1}{2} \lim_{k \rightarrow \infty} \|B^{1/2}(u(t_k) - u^*)\|_{L^2(\Omega)}^2 = 0.$$

We assert that $t \mapsto H_R(u(t)|u^*)$ is nonincreasing for a.e. $t > 0$. Indeed, we know from Section 4.3 that $t \mapsto H_R(u(t))$ is nonincreasing. Furthermore, since $\int_{\Omega} u(t) dx = \int_{\Omega} u^* dx$ and Bu^* is a constant vector, we have $\int_{\Omega} u(t)^T Bu^* dx = \int_{\Omega} u(s)^T Bu^* dx$ for $t \geq s$. Hence, for $t \geq s$,

$$\begin{aligned} H_R(u(t)|u^*) &= H_R(u(t)) + H_R(u^*) - \int_{\Omega} u(t)^T Bu^* dx \\ &\leq H_R(u(s)) + H_R(u^*) - \int_{\Omega} u(s)^T Bu^* dx = H_R(u(s)|u^*), \end{aligned}$$

proving the claim.

We conclude that $H_R(u(t)|u^*) \leq H_R(u(t_k)|u^*) \rightarrow 0$ for $t \geq t_k \rightarrow \infty$. It follows from the positive definiteness of $B^{1/2}$ on L^\perp that

$$\|\widehat{u}(t) - \widehat{u}^*\|_{L^2(\Omega)} \leq C\|B^{1/2}(\widehat{u}(t) - \widehat{u}^*)\|_{L^2(\Omega)} \leq 2H_R(u(t)|u^*) \rightarrow 0$$

as $t \rightarrow \infty$. This finishes the proof of Theorem 9.

APPENDIX A. AUXILIARY RESULTS

Let the matrix $B = (b_{ij}) \in \mathbb{R}^{n \times n}$ be symmetric positive semidefinite. Then the square root of $B^{1/2}$ exists and $z^T B z = |B^{1/2} z|^2$ for $z \in \mathbb{R}^n$. Let P_L and P_{L^\perp} be the projection matrices onto $L = \ker B = \ker B^{1/2} \neq \{0\}$ and $L^\perp = \text{ran } B$, respectively.

Lemma 16. *Let $\lambda > 0$ be the smallest positive eigenvalue of $B^{1/2}$. Then*

$$|P_{L^\perp} z| \leq \lambda^{-1} |B^{1/2} z| \quad \text{for } z \in \mathbb{R}^n.$$

Proof. Let $z \in \mathbb{R}^n$ and $\widehat{z} = P_{L^\perp} z$. By definition of λ , $|B^{1/2} \widehat{z}|^2 \geq \lambda |\widehat{z}|^2$. Then the conclusion follows from $B^{1/2} \widehat{z} = B^{1/2} z - B^{1/2} P_L z = B^{1/2} z$. \square

We introduce the relative entropy densities

$$h_S(u|v) = \sum_{i=1}^n (\mathfrak{h}(u_i) - \mathfrak{h}(v_i) - \mathfrak{h}'(v_i)(u_i - v_i)) = \sum_{i=1}^n \left(u_i \log \frac{u_i}{v_i} - (u_i - v_i) \right),$$

$$h_R(u|v) = \frac{1}{2} (u - v)^T B (u - v) = \frac{1}{2} |B^{1/2}(u - v)|^2, \quad u, v \in [0, \infty)^n,$$

where $\mathfrak{h}(z) = z(\log z - 1) + 1$. We denote by $\|A\|_2$ the norm of A induced by the Euclidean norm $|\cdot|$ in \mathbb{R}^n .

Lemma 17 (Coercivity). *Let $a_0 = \frac{1}{2} \min_{i \in \{1, \dots, n\}} b_{ii} > 0$, $a_1 = \|B\|_2$, and let $K \geq 1$. Then there exists a constant $c_* > 0$, only depending on a_0 , a_1/a_0 , and M , such that for all $u, v \in \mathbb{R}_{\geq}^n$ with $0 < |v| \leq M$,*

$$h_S(u|v) + h_R(u|v) \geq c_* |u - v|^2.$$

Proof. By assumption, we have $\frac{1}{2} u^T B u \geq \frac{1}{2} \sum_{i=1}^n b_{ii} u_i^2 \geq a_0 |u|^2$ for all $u \in \mathbb{R}_{\geq}^n$. If $(a_0/2)|u| \geq a_1|v|$ then

$$\begin{aligned} h_R(u|v) &= \frac{1}{2} u^T B v - v^T B u + \frac{1}{2} v^T B v \geq a_0 |u|^2 - a_1 |u| |v| + a_0 |v|^2 \\ &\geq a_0 |u|^2 - \frac{a_0}{2} |u|^2 + a_0 |v|^2 = \frac{a_0}{2} |u|^2 + a_0 |v|^2 \geq \frac{a_0}{3} |u - v|^2. \end{aligned}$$

Next let $(a_0/2)|u| < a_1|v|$. We find for $f(z) = z \log z$ that

$$\begin{aligned} u_i \log \frac{u_i}{v_i} - (u_i - v_i) &= f(u_i) - f(v_i) - f'(v_i)(u_i - v_i) \\ &= (u_i - v_i) \int_0^1 (f'(s(u_i - v_i) + v_i) - f'(v_i)) \Big|_{s=0}^\theta d\theta \end{aligned}$$

$$= (u_i - v_i)^2 \int_0^1 \int_0^\theta f''(s(u_i - v_i) + v_i) ds d\theta.$$

Then we infer from $|u_i/v_i| < 2a_1/a_0$ that

$$f''(s(u_i - v_i) + v_i) = \frac{1}{v_i(s(u_i/v_i - 1) + 1)} > \frac{1}{M(s(2a_1/a_0 - 1) + 1)}$$

and consequently,

$$u_i \log \frac{u_i}{v_i} - (u_i - v_i) \geq \frac{(u_i - v_i)^2}{M} \int_0^1 \int_0^\theta \frac{ds d\theta}{s(2a_1/a_0 - 1) + 1},$$

which shows that $h_S(u|v) \geq c_1|u - v|^2$, where

$$c_1 = \frac{1}{M} \min_{i=1, \dots, n} \int_0^1 \int_0^\theta \frac{ds d\theta}{s(2a_1/a_0 - 1) + 1}.$$

Putting these estimates together and observing that $h_S(u|v) \geq 0$, $h_R(u|v) \geq 0$, we conclude the proof with $c_* = \min\{a_0/3, c_1\}$. \square

REFERENCES

- [1] H. Amann. Nonhomogeneous linear and quasilinear elliptic and parabolic boundary value problems. In: H. J. Schmeisser and H. Triebel (eds.), *Funct. Spaces Diff. Oper. Nonlin. Anal.*, pp. 9–126. Teubner, Wiesbaden, 1993.
- [2] M. Bertsch, M. Gurtin, D. Hilhorst, and L. Peletier. On interacting populations that disperse to avoid crowding: preservation of segregation. *J. Math. Biol.* 23 (1985), 1–13.
- [3] M. Bertsch, D. Hilhorst, H. Izuhara, and M. Mumura. A nonlinear parabolic-hyperbolic system for contact inhibition of cell-growth. *Diff. Eqs. Appl.* 4 (2012), 137–157.
- [4] D. Bothe. On the Maxwell–Stefan equations to multicomponent diffusion. In: *Progress in Nonlinear Differential Equations and their Applications*, pp. 81–93. Springer, Basel, 2011.
- [5] Y. Brenier, C. De Lellis, L. Székelyhidi. Weak–strong uniqueness for measure-valued solutions. *Commun. Math. Phys.* 305 (2011), 351–361.
- [6] M. Burger, J. A. Carrillo, J.-F. Pietschmann, and M. Schmidtchen. Segregation effects and gap formation in cross-diffusion models. *Interfaces Free Bound.* 22 (2020), 175–203.
- [7] J. A. Cañizo, J. A. Carrillo, P. Laurençot, and J. Rosado. The Fokker–Planck equation for bosons in 2D: well-posedness and asymptotic behavior. *Nonlin. Anal.* 137 (2016), 291–305.
- [8] C. Chainais-Hillairet, J.-G. Liu, and Y.-J. Peng. Finite volume scheme for multi-dimensional drift-diffusion equations and convergence analysis. *ESAIM Math. Model. Numer. Anal.* 37 (2003), 319–338.
- [9] L. Chen, E. Daus, and A. Jüngel. Rigorous mean-field limit and cross diffusion. *Z. Angew. Math. Phys.* 70 (2019), no. 122, 21 pages.
- [10] C. Christoforou and A. Tzavaras. Relative entropy for hyperbolic–parabolic systems and application to the constitutive theory of thermoviscoelasticity. *Arch. Ration. Mech. Anal.* 229 (2018), 1–52.
- [11] S. Demoulini, D. Stuart, and A. Tzavaras. Weak–strong uniqueness of dissipative measure-valued solutions for polyconvex elastodynamics. *Arch. Ration. Mech. Anal.* 205 (2012), 927–961.
- [12] K. Deimling. *Nonlinear Functional Analysis*. Springer, Berlin, 1985.
- [13] A. Ern and J.-L. Guermond. *Finite Elements I: Approximation and Interpolation*. Springer, Cham, 2021.
- [14] J. Escher, A.-V. Matioc, and B.-V. Matioc. Modelling and analysis of the Muskat problem for thin fluid layers. *J. Math. Fluid Mech.* 14 (2012), 267–277.

- [15] M. Di Francesco, A. Esposito, and S. Fagioli. Nonlinear degenerate cross-diffusion systems with non-local interaction. *Nonlin. Anal.* 169 (2018), 94–117.
- [16] R. DiPerna. Measure-valued solutions to conservation laws. *Arch. Ration. Mech. Anal.* 88 (1985), 223–270.
- [17] P.-E. Druet, K. Hopf, and A. Jüngel. Hyperbolic–parabolic normal form and local classical solutions for cross-diffusion systems with incomplete diffusion. Submitted for publication, 2022. arXiv:2210.17244.
- [18] P.-E. Druet and A. Jüngel. Analysis of cross-diffusion systems for fluid mixtures driven by a pressure gradient. *SIAM J. Math. Anal.* 52 (2020), 2179–2197.
- [19] C. Elliott and V. Janovskiy. An error estimate for a finite-element approximation of an elliptic variational inequality formulation of a Hele–Shaw moving-boundary problem. *IMA J. Numer. Anal.* 3 (1983), 1–9.
- [20] R. Eymard, T. Gallouët, and R. Herbin. Finite volume methods. In: P. G. Ciarlet and J.-L. Lions (eds.). *Handbook of Numerical Analysis* 7 (2000), 713–1018.
- [21] E. Feireisl, P. Gwiazda, A. Świerczewska-Gwiazda, and E. Wiedemann. Dissipative measure-valued solutions to the compressible Navier–Stokes system. *Calc. Var. Partial Diff. Eqs.* 55 (2016), no. 141, 20 pages.
- [22] E. Feireisl and M. Lukáčová-Medvid’ová. Convergence of a mixed finite element–finite volume scheme for the isentropic Navier–Stokes system via dissipative measure-valued solutions. *Found. Comput. Math.* 18 (2018), 703–730.
- [23] E. Feireisl, M. Lukáčová-Medvid’ová, and H. Mizerová. Convergence of finite volume schemes for the Euler equations via dissipative measure-valued solutions. *Found. Comput. Math.* 20 (2020), 923–966.
- [24] U. Fjordholm, R. Käppeli, S. Mishra, and E. Tadmor. On the computation of measure-valued solutions. *Acta Numerica* 25 (2016), 567–679.
- [25] K. Friedrichs and P. Lax. Systems of conservation equations with a convex extension. *Proc. Nat. Acad. Sci. USA* 68 (1971), 1686–1688.
- [26] T. Gallouët and J.-C. Latché. Compactness of discrete approximate solutions to parabolic PDEs – Application to a turbulence model. *Commun. Pure Appl. Anal.* 11 (2012), 2371–2391.
- [27] P. Gwiazda, B. Perthame, and A. Świerczewska-Gwiazda. A two-species hyperbolic–parabolic model of tissue growth. *Commun. Partial Diff. Eqs.* 44 (2019), 1605–1618.
- [28] P. Gwiazda, A. Świerczewska-Gwiazda, and E. Wiedemann. Weak–strong uniqueness for measure-valued solutions of some compressible fluid models. *Nonlinearity* 28 (2015), 3873–3890.
- [29] K. Hopf. Singularities in L^1 -supercritical Fokker–Planck equations: A qualitative analysis. arXiv:2107.08531. (Accepted for publication in *Ann. Inst. H. Poincaré C, Anal. Non Lin.*)
- [30] M. Jacobs. Non-mixing solutions to the multispecies porous media equation. Submitted for publication, 2022. arXiv:2208.01792.
- [31] A. Jüngel. *Entropy Methods for Diffusive Partial Differential Equations*. Springer Briefs Math., Springer, 2016.
- [32] A. Jüngel, S. Portisch, and A. Zurek. Nonlocal cross-diffusion systems for multi-species populations and networks. *Nonlin. Anal.* 219 (2022), 112800, 26 pages.
- [33] A. Jüngel and I. Stelzer. Existence analysis of Maxwell–Stefan systems for multicomponent mixtures. *SIAM J. Math. Anal.* 45 (2013), 2421–2440.
- [34] A. Jüngel and A. Zurek. A finite-volume scheme for a cross-diffusion model arising from interacting many-particle population systems. In: R. Klöfkorn, E. Keilegavlen, F. Radu, and J. Fuhrmann (eds.), *Finite Volumes for Complex Applications IX*, pp. 223–231. Springer, Cham, 2020.
- [35] A. Jüngel and A. Zurek. A convergent structure-preserving finite-volume scheme for the Shigesada–Kawasaki–Teramoto population system. *SIAM J. Numer. Anal.* 59 (2021), 2286–2309.
- [36] S. Kawashima and Y. Shizuta. On the normal form of the symmetric hyperbolic–parabolic systems associated with the conservation laws. *Tohoku Math. J.* 40 (1988), 449–464.

- [37] P. Laurençot and B. Matioc. Weak–strong uniqueness for a class of degenerate parabolic cross-diffusion systems. *Arch. Math. (Brno)* 59 (2023), 201–213.
- [38] P.-L. Lions. *Mathematical topics in fluid dynamics. Vol. 1: Incompressible models*. Oxford Science Publication, Oxford, 1996.
- [39] T. Lorenzi, A. Lorz, and B. Perthame. On interfaces between cell populations with different mobilities. *Kinet. Relat. Models* 10 (2017), 299–311.
- [40] F. Nabet. Convergence of a finite-volume scheme for the Cahn–Hilliard equation with dynamic boundary conditions. *IMA J. Numer. Anal.* 36 (2016), 1898–1942.
- [41] A. Oulhaj. Numerical analysis of a finite volume scheme for a seawater intrusion model with cross-diffusion in an unconfined aquifer. *Numer. Meth. Partial Diff. Eqs.* 34 (2018), 857–880.
- [42] P. Pedregal. *Parametrized Measures and Variational Principles*. Birkhäuser, Basel, 1997.
- [43] C. Rao. Diversity and dissimilarity coefficients: a unified approach. *Theor. Popul. Biol.* 21 (1982), 24–43.
- [44] L. Tartar. Compensated compactness and applications to partial differential equations. In: R. Knops (ed.), *Nonlinear Analysis and Mechanics: Heriot–Watt Symposium*, Res. Notes Math. 39, pp. 136–212. Pitman, Boston, 1979.

KATHARINA HOPF, WEIERSTRASS INSTITUTE FOR APPLIED ANALYSIS AND STOCHASTICS (WIAS),
MOHRENSTR. 39, 10117 BERLIN, GERMANY

Email address: hopf@wias-berlin.de

ANSGAR JÜNGEL, INSTITUTE OF ANALYSIS AND SCIENTIFIC COMPUTING, TECHNISCHE UNIVERSITÄT
WIEN, WIEDNER HAUPTSTRASSE 8–10, 1040 WIEN, AUSTRIA

Email address: juengel@tuwien.ac.at