

## problem sheet 6

discussion: week of Monday, 15.11.2021

- 6.1.** a) Write a program with signature  $y = \text{composite\_gauss}(n, L, q)$  that realizes a composite Gauss rule for integration over  $(0, 1)$ . The composite Gauss rule uses  $n$  points for each of the  $L$  subintervals that are given by

$$(0, q^{L-1}), (q^{L-1}, q^{L-2}), (q^{L-2}, q^{L-3}) \dots, (q, 1)$$

Check your program with  $f(x) = x^m$ ,  $m = 0, 1, 2$ . *Hint:* Gauss points and weights can be obtained by `numpy.polynomial.legendre.leggauss` or `gauleg.m` (see homepage).

- b) Use your routine `composite_gauss` for  $n = L = 1, \dots, 20$  and the three choices  $q \in \{0.5, 0.15, 0.05\}$  and the integrand

$$f(x) = x^{0.1} \log x.$$

(The exact integral is  $\int_0^1 f(x) dx = -1/1.1^2 \approx -0.82644$ .) Plot semilogarithmically (`semilogy`) the quadrature error versus  $n$  for these 3 values of  $q$ . Which choice of  $q$  is the best one?

- c) Fit (using `polyfit`) the error curves to the law  $Ce^{-bn}$ .

- 6.2.** Give an explicit error bound (in dependence on  $n$ ) for the Gaussian quadrature error

$$\left| \int_{-1}^1 f(x) dx - Q_n^{Gauss}(f) \right| \quad \text{with } f(x) = (4 - x^2)^{-1}.$$

- 6.3.** (transformation techniques) we seek a quadrature formula for

$$\int_1^\infty f(x) dx.$$

Consider the specific case  $f(x) = \log x / (x^\pi)$  with

$$\int_1^\infty \frac{\log x}{x^\pi} dx = \frac{1}{\pi^2 - 2\pi + 1}.$$

- a) One possibility is to transform the integral to an integration over  $(0, 1)$  using a suitable substitution. Formulate such a transformation. The transformed problem can then be treated with the quadrature formula of Problem 6.1.
- b) Another option is the substitution  $x = e^y$ . One obtains an integral of the form

$$\int_{y=0}^\infty F(y) dy,$$

where the integrand decays rapidly so that the integral  $\int_{y=0}^\infty F(y) dy$  can be approximated well by  $\int_{y=0}^L F(y) dy$ . Again, the integral can be computed with a composite Gauss rule with  $n$  points per subinterval where the  $L$  subintervals are given by

$$(0, Lq^{L-1}), (Lq^{L-1}, Lq^{L-2}), \dots, (Lq, L)$$

Generate the composite quadrature rule using your program of Problem 6.1.

- c) Plot the error using `semilogy` for both methods with  $n = L = 1, \dots, 20$ . Choose  $q = 0.15$ .

- 6.4.** Consider quadrature rules  $Q^{2D}$  on the square  $S = [0, 1]^2$ .

- a) Show: the midpoint rule  $Q(F) = F(0.5, 0.5)$  is exact for polynomials of the form  $F(x, y) = a + bx + cy$ .

- b) Given  $p \in \mathbb{N}_0$ , give a quadrature formula  $Q^{2D}$  that is exact for polynomials of the form  $F(x, y) = \sum_{i,j=0}^p a_{ij} x^i y^j$ .

**6.5.** Consider the function

$$\varphi(x) = \sqrt{x+1} - \sqrt{x}$$

- a) Is the evaluation of  $\varphi$  well-conditioned for large  $x$ ? Consider relative conditioning.
- b) Formulate a stable numerical realization of  $\varphi$  (*Hint:* You may use that a stable realization of  $\sqrt{\cdot}$  is available.)