

numerical differentiation with one-sided difference quotients

$$D(h) = \frac{f(x_0 + h) - f(x_0)}{h}$$

h	$m = 0$	$m = 1$	$m = 2$	$m = 3$
h_0	$D(h_0) = D_{00}$	D_{01}	D_{02}	D_{03}
h_1	$D(h_1) = D_{10}$	D_{11}	D_{12}	D_{13}
h_2	$D(h_2) = D_{20}$	D_{21}	D_{22}	D_{23}
h_3	$D(h_3) = D_{30}$	D_{31}	D_{32}	D_{33}
h_4	$D(h_4) = D_{40}$	D_{41}	D_{42}	\vdots
h_5	$D(h_5) = D_{50}$	D_{51}	\vdots	
h_6	$D(h_6) = D_{60}$	\vdots		
\vdots	\vdots			

$$D_{i0} = D(h_i)$$

$$D_{ij} = D_{(i+1)(j-1)} - \frac{h_{i+j}}{h_{i+j} - h_i} [D_{(i+1)(j-1)} - D_{i(j-1)}], \quad j \geq 1$$

example: $f(x) = e^x$ and $x_0 = 0$ with one-sided difference quotients

Neville scheme:

h	$m = 0$	$m = 1$	$m = 2$	$m = 3$
1	1.71828182845905	0.87660325434147	1.00747997135508	0.99982039920503
2^{-1}	1.29744254140026	0.97476079210167	1.00077784572378	0.99999046433634
2^{-2}	1.13610166675097	0.99427358231826	1.00008888700977	0.99999944973780
2^{-3}	1.06518762453461	0.99863506083689	1.00001062939680	0.99999996693993
2^{-4}	1.03191134268575	0.99966673725682	1.00000129974704	0.99999999797395
2^{-5}	1.01578903997129	0.99991765912448	1.00000016069559	0.99999999987449
2^{-6}	1.00785334954789	0.99997953530281	1.00000001997713	
2^{-7}	1.00391644242535	0.99999489880855		
2^{-8}	1.00195567061695			

example: $f(x) = e^x$ and $x_0 = 0$ with one-sided difference quotients

error and reduction factor:								
h	$m = 0$	ρ	$m = 1$	ρ	$m = 2$	ρ	$m = 3$	ρ
1	7.18 ₋₁	—	-1.23 ₋₁	—	7.48 ₋₃	—	-1.78 ₋₄	—
2^{-1}	2.97 ₋₁	0.41	-2.52 ₋₂	0.20	7.78 ₋₄	0.10	-9.54 ₋₆	0.053
2^{-2}	1.36 ₋₁	0.46	-5.73 ₋₃	0.23	8.89 ₋₅	0.11	-5.50 ₋₇	0.058
2^{-3}	6.52 ₋₂	0.48	-1.37 ₋₃	0.24	1.06 ₋₅	0.12	-3.31 ₋₈	0.060
2^{-4}	3.19 ₋₂	0.49	-3.33 ₋₄	0.24	1.30 ₋₆	0.12	-2.03 ₋₉	0.061
2^{-5}	1.58 ₋₂	0.49	-8.23 ₋₅	0.25	1.61 ₋₇	0.12	-1.26 ₋₁₀	0.062
2^{-6}	7.85 ₋₃	0.50	-2.05 ₋₅	0.25	1.99 ₋₈	0.12		
2^{-7}	3.92 ₋₃	0.50	-5.10 ₋₆	0.25				
2^{-8}	1.96 ₋₃	0.50						
expected:	$O(h)$		$O(h^2)$		$O(h^3)$		$O(h^4)$	
		0.5		0.25		0.125		0.0625