Session of Focus Materialchemie – Wednesday, 10.01.2024 16:00 – @ Seminarraum Lehar 02 (TU-Wien, Getreidemarkt 9, BC, OG. 02, room A46) – join us on ZOOM (ID: 983 0066 2349)

## (Nano)spectroscopic signatures of intricate relations between 2D layers and their substrate

## Otakar Frank

J. Heyrovský Institute of Physical Chemistry, Czech Academy of Sciences, Dolejškova 2155/3, 182 23 Prague, Czech Republic

otakar.frank@jh-inst.cas.cz

Common spectroscopic investigation of two-dimensional materials and their van der Waals (vdW) heterostructures mostly relies either on diffraction-limited microRaman or photoluminescence (PL). However, these methods do not properly capture local variations caused by, e.g., nanometre-sized heterogeneities stemming from contamination trapped between the layers or complex strain and charge patterns formed by strong out-of-plane interactions.

Tip-enhanced spectroscopy methods enable the access to information on the local lattice deformation and also on the interaction between the individual layers composing the heterostructure. What may appear as peak splitting in micro-Raman or PL spectra of transition metal dichalcogenides (TMDC) on metal substrates or of vdW heterobilayers, can, in fact, often come from mixing up signals from various regions within the laser spot, including new or discretely shifted peaks. In other cases, however, peak splitting can indicate lifting the degeneracy of the phonon, due to, for example, uniaxial deformation. Spectroscopic fingerprints, both on micro- and nanoscale, of variously interacting vdW layers will be discussed, including TMDCs on metals [1-3] and TMDC heterobilayers [4-6].

## References:

- [1] Velicky et al. J. Phys. Chem. Lett. 11, 6112 (2020)
- [2] Velicky et al. Adv. Mater. Interfaces 7, 2001324 (2020)
- [3] Rodriguez et al. Phys. Rev. B 105, 195413 (2022)
- [4] Rodriguez et al. 2D Mater. 8, 025028 (2021)
- [5] Rodriguez et al. J. Phys. Chem. Lett. 13, 5854 (2022)
- [6] Rodriguez et al., ACS Nano 17, 7787 (2023)